

20 DEC 1991

BEST AVAILABLE COPY

Mr. David Paradise, Jr.
The Andersons Lawn Products Division
Green Free Flow Fertilizer
P.O. Box 115
Maumee Ohio 43507

Dear Mr. Paradise:

Subject: Eco Time Fertilizer 20-4-10 With EcoStar
EPA Registration No. 5198-75
Re: Label Amendment
Your submission dated November 20, 1991

Your labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide, and Fertilizer Act as amended is acceptable provided that you make the label changes under the "Environmental Hazard" Section of the label replace the statement "do not apply directly to water or wetland" with the statement "do not apply directly to water to areas where surface water is present or to intertidal areas below the means high water mark." Please submit five (5) copies of your final printed labels incorporating this amendment. A stamped copy is enclosed for your records.

Sincerely,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (4750FC)

enclosure		CONCURRENCES					
SYMBOL	H7505C						
SURNAME	Matys						
DATE	12/18/91						

2 of 4

(FRONT PANEL)

Tee Time[®] FERTILIZER 20-4-10 with Ronstar[®]

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN, wash with soap and water.
IF IN EYES, flush with plenty of water; get medical attention.

See Back Panel for Additional Precautionary Statements

Active ingredient:
Oxadiazon (2-tert-butyl-4-(2,4-dichloro-5-
isopropoxyphenyl)-6-1,3,4-oxadiazin-6-one) 1.38%
Inert ingredients 98.62%
Total

Net Weight 50 Lbs. (22.7 kg)
Treats Up To 16,666 Sq. Ft.

Ronstar - Trademark of Rhone-Poulenc
Tee Time - Trademark of The Andersons

EPA Reg. No. 9198-75
EPA Est. 9198-OH-1

ACCEPTED
with COMMENTS
in EPA Letter #10-10-91

20 DEC 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, this product is registered under FIFRA No. 9198-75

9198-75

BEST AVAILABLE COPY

Net Weight 50 Lbs. (22.7 kg) Tee Time® Fertilizer 20-4-10 with Ronstar® Treats Up To 16,666 Sq. Ft.

Guaranteed Analysis:	
Total Nitrogen (N)	20%
1.6% Ammoniacal Nitrogen	
12.4% Urea Nitrogen*	
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	10%
Sulfur (S)	2.0%
*Primary plant foods derived from Ammoniated Phosphates, Urea, Sulfur Coated Urea and Murates of Potash.	
**8.0% Slowly Available Urea Nitrogen from Sulfur Coated Urea	

PRECAUTIONARY STATEMENTS WARNING

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Avoid contamination of feed and foodstuffs. Keep opened containers away from children and animals. Wear protective clothing including long-sleeved shirt and gloves when handling. Wash thoroughly after handling. Wash clothing and equipment after handling this product.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not contaminate water by disposing of equipment washings. Do not apply directly to water.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION: This product is a selective pre-emergent herbicide for control of many annual grasses and broadleaf weeds in turf. This product controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

Do not apply to wet turf as discoloration may occur.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

This product will control the following weeds when used as directed:

Grassy Weeds
 Crabgrass
 Goosegrass
 Green and Yellow Foxtail
 Sandbur
 Annual Bluegrass
 Barnyardgrass

Broadleaf Weeds
 Oxalis
 Spurge
 Veronica Speedwell
 Common Purslane
 Florida Pursley
 Common Groundsel

Carpetgrass
 Pigweed
 Stinging Nettle

BEST AVAILABLE COPY

ORNAMENTAL TURF INCLUDING GOLF COURSE FAIRWAYS, TEES AND PARKS - Established perennial Bluegrass, Bermudagrass, Perennial Ryegrass, Tall Fescue, Zoysia and St. Augustinegrass turf:

For most economical use, rates of this product should be based on anticipated weed population, time of application and length of growing season. A well-maintained turf with low weed infestations will require less herbicide for effective control than will a thin turf with low fertility, minimum maintenance and heavy traffic. Geographic areas with longer growing seasons require higher rates. When cultural practices require application of this product well in advance of weed seed germination, rates should be adjusted upward to compensate for normal breakdown of the active ingredient.

Rake turf area free of leaves, mow, if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results.

If rain is not expected shortly after application, water turf thoroughly to increase effectiveness. When using rates below 180 pounds of product per acre, thoroughly watering assures more even distribution of the herbicide and avoids gaps in coverage.

Northern Turf: Crabgrass and other grasses and listed broadleaf weeds germinating in the spring: Apply at 3.0 to 6.7 lbs. per 1000 sq. ft. to control weeds 1 to 2 weeks prior to germination, usually during March and April.

Poa annua (Annual Bluegrass): Apply at 5.0 lbs. per 1000 sq. ft. in two consecutive rounds 6-8 weeks apart in late summer to fall, or in early spring, starting 1-2 weeks before expected weed seed germination, usually from August until winter, then just before spring thaw. Will not control weeds present at time of application. (Second of these two rounds may be reduced to 3.3 lbs. per 1000 sq. ft.)

Late germinating annual grasses and broadleaf weeds such as Prostrate Spurge: Apply at 3.0 to 6.7 lbs. per 1000 sq. ft. in two consecutive rounds 6-8 weeks apart in late April through July, beginning 1-2 weeks before expected weed seed germination.

Southern Turf: Goosegrass and Crabgrass: Apply at 3.0 to 6.7 lbs. per 1000 sq. ft. in two consecutive rounds 6-8 weeks apart (optionally reduced to 3.3 lbs. per 1000 sq. ft. in the second of these rounds) to control weeds, starting 1-2 weeks prior to germination, usually during January through March.

Poa annua (Annual Bluegrass): Apply at 5.0 lbs. per 1000 sq. ft. in two consecutive rounds 6-8 weeks apart (optionally reduced to 3.3 lbs. per 1000 sq. ft. in the second of these rounds) in late summer to fall, starting 1-2 weeks before expected weed seed germination, usually from mid-August through mid-November. Will not control weeds present at time of application.

Annual weeds as listed: Apply at 3.0 to 6.7 lbs. per 1000 sq. ft. in two consecutive rounds 6-8 weeks apart in fall, winter and spring, starting 1-2 weeks before expected weed seed germination, usually from mid-August through November for winter annuals, and January through mid-March for summer annuals.

Use the higher rate where weed pressure is expected to be greater, the lower where turf has been in a yearly maintenance program which has generally kept weeds in check.

If rain is not expected within 24 hours of the application, sprinkle the treated turf with 1/4 to 1/2 inch of water for maximum weed control.

For St. Augustinegrass turf, do not exceed the rate of 175 pounds per acre (2.4 pounds active ingredient). This product may cause temporary discoloration of Bermudagrass and St. Augustinegrass. This is normally outgrown within 2-3 weeks.

Red Fescue and Bentgrass are not tolerant of this product. The use of this product on turf containing Red Fescue or Bentgrass should be limited to situations where they are undesirable grasses. Do not apply to Dichondra, Centipedegrass or newly seeded areas.

Because of long residual activity, seeding into treated areas should be delayed until 4 months after treatment. Avoid contact with flowers and shrubs. This product should not be used on putting greens because the varying cultural practices used.

Apply with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly.

4 of 4

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides or seeds. In the event of a spill, sweep up minimizing dust and use or dispose of according to directions. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or, incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and soil conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

K1 - (ATTBLCSB - BACK)
DATE: 11/27/91

ATT206R15

BEST AVAILABLE COPY