



Tee TimeTM Fertilizer 32-3-5

ACCEPTED
JUN 30 1986
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 9198-76

with **Oftanol**[®]

the professional's
partnerTM



COMMERCIAL USE ONLY


ACTIVE INGREDIENTS
1-Methylethyl 2-(ethoxy(1-
methylethylamino)phosphon-
thio)oxybenzoate 15%
INERT INGREDIENTS 85%
100.0%

U.S. Patent No. 3,621,082
Canadian Patent No. 825,198
EPA Reg. No. 9198-76
EPA Est. No. 9198-OH-1

OFTANOL[®] is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen

STOP READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR STATEMENT OF PRACTICAL
TREATMENT AND ADDITIONAL PRECAUTIONARY STATEMENTS

Net Weight 50 Lbs. (22.7 Kg) Treats Up To 16,650 Sq. Ft.



Net Weight 50 Lbs. (22.7 Kg)
Treats Up To 16,650 Sq. Ft.

Tee Time Fertilizer 32-3-5

with **Oftanol**

GUARANTEED ANALYSIS

Total Nitrogen (N)	32%
1.2% Ammoniacal Nitrogen	
24.3% Urea Nitrogen	
8.5% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	5%
Sulfur (S)	4.9%
Primary Plant Foods Derived from Urea, Murate of Potash, Ammoniated Phosphates, and Sulfur Coated Urea. Sulfur derived from Sulfur Coated Urea. Potential Acidity 1500 pounds Calcium Carbonate equivalent per ton.	

PRECAUTIONARY STATEMENTS Hazards To Humans and Domestic Animals CAUTION

Harmful if swallowed or absorbed through the skin. May cause eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea. **IN CASE OF POISONING,** call a physician or Poisoning Control Center. Have patient lie down and keep quiet.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, VOMITING SHOULD BE INDUCED. Administer water freely and induce vomiting by giving one dose (½ oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to unconscious person. Get medical attention immediately. **IF ON SKIN,** wash skin immediately with soap and warm water. If irritation occurs, get medical attention. **IF EYES ARE CONTAMINATED,** wash with plenty of water. Get medical attention. **IF INHALED,** remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

TO PHYSICIAN: ANTIDOTE: Atropine sulfate has a slight antidotal effect and should be administered in therapeutic doses. Repeat as necessary to the point of tolerance.

Active ingredient inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling. **IMPORTANT:** Read these entire Directions for Use and Conditions of Sale before using Tee Time Fertilizer with Oftanol. **CONDITIONS OF SALE:** THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS AND/OR INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO OTHER CROPS, ANIMALS, MAN OR THE ENVIRONMENT. THE ANDERSONS OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF THE ANDERSONS AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

RECOMMENDED APPLICATIONS

The use of this product at the rate of 3 lbs. per 1,000 sq. ft. will provide 0.96 lb. of nitrogen per 1,000 sq. ft. At the same time the application will provide control of the below listed pests. The timing of the applications vary dependent upon the insect which is being controlled. Follow the directions as specified below for control of the listed turf grass pests. Consult your local turf, State Agricultural Experiment Station, or State Extension Specialist for more specific information regarding the timing of applications.

White Grub larvae (such as Japanese Beetle, Black Turf Grass Aeneas, Northern Masked Chaffer, European Chaffer, June and May Beetles).

PREVENTIVE APPLICATIONS

To control the above listed pests apply Tee Time Fertilizer with OFTANOL at the rate of 3 lbs./1,000 sq. ft. in the Spring (April thru mid May) and immediately apply sufficient water to wet the soil 1 to 1½ inches deep without causing runoff.

CURATIVE APPLICATIONS

If visible damage or high Grub populations have been observed apply 3 lbs./1,000 sq. ft. and immediately apply sufficient water to wet the soil 1 to 1½ inches deep without causing runoff.

EMerald Larvae: Apply 3 lbs./1,000 sq. ft. when adult activity is observed in the Spring usually late March to mid May. Water in as for preventive White Grub control.

Hyperodes Weevil Larvae: Apply at the rate of 3 lbs. per 1,000 sq. ft. in the Spring between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15 and water in as for preventive White Grub control.

Mole Crickets: Apply at the rate of 3 lbs./1,000 sq. ft. preventively or when damage is observed. Apply ½ to 1 inch of water immediately after application if rainfall does not occur.

Area Fire Ant Control: Apply 3 lbs./1,000 sq. ft. uniformly over area to be treated for residual control. Water immediately after application using ½ to 1 inch of water. If established fire ant mounds are present use the mound drunch treatment prior to area application. If a mound drunch treatment is made only one area treatment may be made.

Established Fire Ant Mounds—Drunch Treatments: Wet Conditions: Distribute 0.5 oz. evenly over 8 sq. ft. and immediately sprinkle gently with 3 gallons of water. **Dry Conditions:** Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one Mound Drunch Treatment.

Soil Weeviler Larvae and Chinch Bugs: Apply 1.5 lbs. or 3 lbs. per 1,000 sq. ft. (use higher rate for severe infestations) when insect damage is first observed. Apply ½ to 1 inch of water immediately after treatment. Re-apply as necessary for pest re-infestation (see restrictions).

APPLICATION TECHNIQUES

The application of Tee Time with OFTANOL can be made using a drop spreader. When using a drop spreader the application can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount of Tee Time Fertilizer with OFTANOL over the area, then immediately apply the remaining half at right angles or crosswise. The full recommended rate must be used for maximum benefit.

When applying Tee Time Fertilizer with OFTANOL, use appropriate marking systems and make application when the wind is calm to prevent missed areas.

SPREADER SETTINGS

Approximate spreader settings to apply Tee Time fertilizer with OFTANOL at the required rate are listed below. They are calculated to deliver the correct amount based on new equipment and ideal conditions. This serves as a guide only. Calibration of each spreader is necessary to apply an accurate rate.

SUGGESTED SPREADER SETTINGS

Spreader	Spread Width	1.5 Lbs. Per 1,000 Sq. Ft.	3 Lbs. Per 1,000 Sq. Ft.
Cyclone (B1-80)	10-12 Ft.	4 1/4"	5 1/4"
Scotts Professional Broadcast (R-74)	10-12 Ft.	C.D.	D 1/2 E 1/4
Lely (Type W)	30 Ft.	3 1/4"	5

Do not mow treated turf before rainfall or irrigation has been applied as per recommended applications. Keep children and pets off treated areas until dry. Do not use sponges from treated areas for feed or forage or allow food animals to graze treated areas.

RESTRICTIONS

The maximum amount of Tee Time Fertilizer with OFTANOL which can be used is 8 lbs. per 1,000 sq. ft. per year. Applications at the rate of 1.5 lbs. per 1,000 sq. ft. must have a 15 day interval and applications at the rate of 3 lbs. per 1,000 sq. ft. must have a 30 day interval between applications. A maximum of 1.45 ounces of active ingredient per 1,000 sq. ft. per acre per year can be applied regardless of formulation.

Manufactured by

the professional's partner



Lawn Fertilizer Division
P.O. Box 119
Maumee, Ohio 43537