

PM 116 9198-76 1075

(FRONT PANEL)

TEE TIME
FERTILIZER
32-3-5

WITH OFTANOLR

the professional's partner™

THE ANDERSONS

COMMERCIAL USE ONLY

ACCEPTED
JUN 12 1990
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registration number 9198-76
EPA Reg. No. 9198-76

ACTIVE INGREDIENTS:

1-Methylethyl 2-((ethoxy((1-methylethyl)amino)phosphorothioyl)oxy)benzoate	1.5%
INERT INGREDIENTS	98.5%
	TOTAL 100.0%

OFTANOLR is a Reg. TM of the Patent Company of Farbenfabriken Bayer GmbH, Leverkusen

U.S. Patent No. 3,621,882
Canadian Patent No. 825,198

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

CAUTION


SEE BACK PANEL FOR STATEMENT OF PRACTICAL TREATMENT AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT 50 LBS. (22.7 Kg)

TREATS UP TO 16,650 SQ.FT.

EPA REG. NO. 9198-76

EPA EST. NO. 9198-OH-1



NET WEIGHT 50 LBS. (22.7 Kg)

TREATS UP TO 16,650 SQ.FT.

TEE TIME FERTILIZER

32-3-5

WITH OFTANOL

GUARANTEED ANALYSIS

Total Nitrogen (N)	32%
1.2% Ammoniacal Nitrogen	
24.3% Urea Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P2O2)	3%
Soluble Potash (K2O)	5%
Sulfur (S)	4.9%

Primary Plant Foods Derived from Urea, Muriate of Potash, Ammoniated Phosphates, and Sulfur Coated Urea. Sulfur derived from Sulfur Coated Urea.

Potential Acidity 1500 pounds Calcium Carbonate equivalent per ton.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENTS OF PRACTICAL TREATMENT


If swallowed: Call a physician or Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If in eyes: flush with plenty of water. If on skin, wash with soap and hot water.

If inhaled: remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.



DIRECTIONS FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use and Conditions of Sale before using Tee Time Fertilizer with Oftanol.

RECOMMENDED APPLICATIONS

The use of this product at the rate of 3 lbs. per 1,000 sq.ft. will provide 0.96 lb. of nitrogen per 1,000 sq.ft. At the same time it's application will provide control of the below listed pests. The timing of the applications vary dependent upon the insect which is being controlled. Follow the directions as specified below for control of the listed turf grass pests. Consult your local turf, State Agricultural Experiment Station, or State Extension Specialists for more specific information regarding the timing of applications.

White Grub larvae (such as Japanese Beetle, Black Turf Grass Alsenius, Northern Masked Chafer, European Chafer, June and May Beetle).

PREVENTATIVE APPLICATIONS

To control the above listed pests apply Tee Time Fertilizer with OFTANOL at the rate of 3 lbs./1,000 sq.ft. in the Spring (April thru mid May) and immediately apply sufficient water to wet the soil 1 to 1 1/2 inches deep without causing runoff.

CURATIVE APPLICATIONS

If visible damage to high Grub populations have been observed apply 3 lbs./1,000 sq.ft. and immediately apply sufficient water to wet the soil 1 to 1 1/2 inches deep without causing runoff.

Billbug Larvae: Apply 3 lbs./1,000 sq.ft. when adult activity is observed in the Spring usually late March to mid May. Water in as for preventative White Grub control.

Hyperodes Weevil Larvae: Apply at the rate of 3 lbs. per 1,000 sq.ft. in the Spring between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15 and water in as for preventative White Grub control.

Mole Crickets: Apply at the rate of 3 lbs./1,000 sq.ft. preventively or when damage is observed. Apply 1/2 to 1 inches of water immediately after application if rainfall does not occur.

Area Fire Ant Control: Apply 3 lbs./1,000 sq.ft. uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present, use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

Established Fire Ant Mounds - Drench Treatments: Wet Conditions: Distribute 0.5 oz. evenly over 9 sq.ft. and immediately sprinkle gently with 3 gallons of water. **Dry Conditions:** Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one Mound Drench Treatment.

Sod Webworm Larvae and Chinch Bugs: Apply 1.5 lbs. or 3 lbs. per 1,000 sq.ft. (use higher rate for severe infestations) when insect damage is first observed. Apply 1/4 to 1/2 inches of water immediately after treatment. Reapply as necessary for pest reinfestation (see restrictions).

APPLICATION TECHNIQUES

This application of Tee Time with OFTANOL can be made using a drop spreader. When using a drop spreader the application can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount of Tee Time Fertilizer with OFTANOL over the area, then immediately apply the remaining half at right angles or crosswise. The full recommended rate must be used for maximum benefit.

When applying Tee Time Fertilizer with OFTANOL, use appropriate marking systems and make application when the wind is calm to prevent missed areas.

SUGGESTED SPREADER SETTINGS

Approximate spreader setting to apply Tee Time Fertilizer with OFTANOL at the required rate are listed below. They are calculated to deliver the correct amount based on new equipment and ideal conditions. This serves as a guide only. Calibration of each spreader is necessary to apply an accurate rate.

(Available at final printing)

RESTRICTIONS

The maximum amount of Tee Time Fertilizer with OFTANOL which can be used is 6 lbs. per 1,000 sq.ft. per year. Applications at the rate of 1.5 lbs. per 1,000 sq.ft. must have a 15 day interval and applications at the rate of 3 lbs. per 1,000 sq.ft. must have a 30 day interval between applications. A maximum of 1.45 oz. of active ingredient per 1,000 sq.ft. per acre per year can be applied regardless of formulation.

Do not mow treated turf before rainfall or irrigation has been applied as per recommended applications. Keep children and pets off treated areas until dry. Do not use clippings from treated areas for feed or forage or allow food animals to graze treated areas.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage or disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

MANUFACTURED BY:

the professional's partner
THE ANDERSONS
LAWN FERTILIZER DIVISION
P.O. BOX 119
MAUMEE, OH 43537