

AUG 31 1992

5417114 300  
18

The Andersons Lawn Fertilizer Division  
d/b/a/Free Flc Fertilizer  
P.O. Box 119  
Maumee, OH 43537

Gentlemen:

Subject: Anderson's Tee-Time 32-3-5 with Oftanol  
EPA Registration No. 9198-76  
Your Labeling Submitted April 8, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the revised labeling:
  - a. The conflict which currently exists on your registered label for the subject product must be resolved. Your front panel includes the statement, "Commercial Use Only" while your "Storage and Disposal" instructions are headed by the statements "For homeowner use only" and "For Professional/Commercial Use only." If it is your intention that the product be used around the home by the homeowner as well as on commercial sites by the professional or commercial uses, the conflicting front panel statement, "Commercial Use Only" must be deleted.

If it is your intention that the product be used only on commercial sites by the commercial user, your label statement, "Commercial Use Only" must be qualified to indicated that it is only a recommendation and is not an EPA-imposed restriction.

63613:I:WP50:Mautz:WP14-3:KEVRIC:08/11/92:09/06/92

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

- b. Update the term, "wetlands" in your "Environmental Hazards" statements to read as follows:

"Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark."

- c. Delete the phrase, "For best results" appearing in your instructions to water in after treatment. Reinstate instructions to water in immediately after treatment as specified in your label accepted June 12, 1990.
- d. Refer to the enclosed copy of PR Notice 83-3 for the appropriate professional/commercial pesticide disposal instructions for your product. Note that the subject pesticide is not included on the "RCRA" F List" and thus is not considered a toxic waste.

- 2. Submit five(5) copies of your final printed label before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Robert A. Forrest  
Product Manager (14)  
Insecticide-Rodenticide Branch  
Registration Division (H7504C)

(FRONT PANEL)

TEE TIME

FERTILIZER

32-3-5

WITH OPTANOL®

COMMERCIAL USE ONLY

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

AUG 31 1992

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
9198-76

ACTIVE INGREDIENTS:

Methylethyl 2-((ethoxy((1-methylethyl)amino)phosphorothioyl)oxy)benzoate .....	1.5%
INERT INGREDIENTS .....	98.5%
TOTAL	100.0%

OPTANOL® is a Reg. TM of the Patent Company of Farbenfabriken Bayer GmbH, Leverkusen

U.S. Patent No. 3,621,082  
Canadian Patent No. 825,198

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR STATEMENT OF PRACTICAL TREATMENT  
AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT \_\_\_\_ LBS.

TREATS UP TO \_\_\_\_\_ SQ. FT.

(BACK PANEL)

NET WEIGHT \_\_\_ LBS. (22.7 Kg)

TREATS UP TO \_\_\_\_\_ SQ.FT.

TEE TIME FERTILIZER

32-3-5

WITH OFTANOL

GUARANTEED ANALYSIS

Total Nitrogen (N) .....	32%
1.2% Ammoniacal Nitrogen	
24.3% Urea Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P2O2) .....	3%
Soluble Potash (K2O) .....	5%
Sulfur (S) .....	4.9%

Primary Plant Foods Derived from Urea, Muriate of Potash, Ammoniated Phosphates, and Sulfur Coated Urea. Sulfur derived from Sulfur Coated Urea.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENTS OF PRACTICAL TREATMENT

- IF SWALLOWED:** Call a physician or Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- IF ON SKIN:** Wash with soap and hot water.
- IF IN EYES:** Flush with plenty of water.
- IF INHALED:** Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.



### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and other wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

### RECOMMENDED APPLICATIONS

Use this product at the rate of 3 lbs. per 1,000 sq.ft. The timing of the applications vary dependent upon the insect which is being controlled. Follow the directions as specified below for control of the listed turf grass pests. Consult your local turf, State Agricultural Experiment Station, or State Extension Specialists for more specific information regarding the timing of applications.

White Grub larvae (such as Japanese Beetle, Black Turf Grass Alsenius, Northern Masked Chafer, European Chafer, June and May Beetle).

### PREVENTATIVE APPLICATIONS

To control the above listed pests apply this product at the rate of 3 lbs./1,000 sq.ft. in the Spring (April thru mid May). For best results, apply sufficient water to wet the soil 1 to 1½ inches deep without causing runoff.

6 of 8

**CURATIVE APPLICATIONS**

If visible damage of high Grub populations have been observed apply 3 lbs./1,000 sq.ft. For best results, apply sufficient water to wet the soil 1 to 1 1/2 inches deep without causing runoff.

**Billbug Larvae:** Apply 3 lbs./1,000 sq.ft. when adult activity is observed in the Spring usually late March to mid May. Water in as for preventative White Grub control.

**Hyperodes Weevil Larvae:** Apply at the rate of 3 lbs. per 1,000 sq.ft. in the Spring between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15 and water in as for preventative White Grub control.

**Grass Crickets:** Apply at the rate of 3 lbs./1,000 sq.ft. preventively or when damage is observed. For best results, apply 1/2 to 1 inches of water after application if rainfall does not occur.

**Area Fire Ant Control:** Apply 3 lbs./1,000 sq.ft. uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present, use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

**Established Fire Ant Mounds - Drench Treatments:** Wet Conditions: Distribute 0.5 oz. evenly over 9 sq.ft. and immediately sprinkle gently with 3 gallons of water. Dry Conditions: Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one Mound Drench Treatment.

**Sod Webworm Larvae and Chinch Bugs:** Apply 1.5 lbs. or 3 lbs. per 1,000 sq.ft. (use higher rate for severe infestations) when insect damage is first observed. For best results, apply 1/2 to 1 inches of water after treatment. Reapply as necessary for pest reinfestation (see restrictions).

**SUGGESTED SPREADER SETTINGS**

(Available at Final Printing)

The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applies and roughness of terrain. Calculate the size of the area to be treated and do not over or under apply. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary, in order to assure correct rate of application. Close spreader when filling, stopping or turning. Do not hand spread.

**BEST AVAILABLE COPY**

7 of 8

### RESTRICTIONS

The maximum amount of this product which can be used is 6 lbs. per 1,000 sq. ft. per year. Applications at the rate of 1.5 lbs. per 1,000 sq.ft. must have a 15 day interval and applications at the rate of 3 lbs. per 1,000 sq. ft. must have a 30 day interval between applications. A maximum of 1.45 oz. of active ingredient per 1,000 sq.ft. per acre per year can be applied regardless of formulation.

Do not mow treated turf before rainfall or irrigation has been applied as per recommended applications. Keep children and pets off treated areas until dry. Do not use clippings from treated areas for feed or forage or allow food animals to graze treated areas.

### STORAGE AND DISPOSAL

#### FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage or disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

#### FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

**WARRANTY**

**MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. **FOLLOW DIRECTIONS CAREFULLY.** Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

**MANUFACTURED BY:** the professional's partner  
THE ANDERSONS  
LAWN FERTILIZER DIVISION  
P.O. BOX 119  
MAUMEE, OH 43537

**Date Printed:** 04/07/92  
**Date Approved:**  
**Supersedes:** 06/12/90