

Reg # 9198-79

PM-23

198

MAR 18 1994

Mr. Daniel Paradiso, Jr.
The Andersons Lawn Fertilizer Division
DEA Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537

Dear Mr. Paradiso:

**Subject: The Andersons Tee Time Fertilizer with 1.15% Team
EPA Registration No. 9198-79
Application Dated March 25, 1994, RE: PR Notice
93-3, WPS Exclusionary Statement, Labeling
Format Revision and Minor Changes and Split Labeling**

The proposed amendments to the registration of the subject pesticide product have been reviewed and found to be acceptable for registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended provided that you:

1. The wetlands statement required by PR Notice 93-3 and 93-8 must appear in the Environmental Hazards section of the "Homeowner Use Only" labeling.
2. On page 4 of 8, "Commercial Use Only" labeling, the change "apply 3.0 to 4 pounds" must be revised to read:

"apply 3.0 to 4.0 pounds per thousand square feet (1.5 to 2.0 pounds active.....)".
3. Submit five(5) printed copies of the final printed label before releasing the product for shipment under the subject labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Stamped copies of the labels are enclosed for your records.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

enclosure (2)

EWilson:Diskette:ABC11:05-18-94

BEST AVAILABLE COPY

29 8

(FRONT PANEL)

HOMEOWNER USE ONLY

THE ANDERSONS

TEE TIME

FERTILIZER

WITH 1.15% TEAM

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass.

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-alpha, alpha, alpha-trifluoro-2,6-dinitro-p-toluidine	0.77%
Trifluralin: alpha, alpha, alpha-trifluoro-2,6-dinitro-N, N-dipropyl-p-toluidine	0.38%
INERT INGREDIENTS	98.85%

KEEP OUT OF THE REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 50 LBS. (22.7 kg)

TREATS UP TO 15,625 SQ. FT.


ACCEPTED
with COMMENTS
In EPA Letter Dated

MAY 18 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9198-79

EPA Reg. No. 9198-79



398

(BACK PANEL)

HOMEOWNER USE ONLY

NET WEIGHT 50 LBS. (22.7 kg)

TREATS UP TO 15,625 SQ.FT.

THE ANDERSONS

TEE TIME FERTILIZER

WITH 1.15% TEAM

GUARANTEED ANALYSIS

Total Nitrogen (N)	25%
1.2% Ammoniacal Nitrogen	
17.3% Urea Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P2O5)	3%
Soluble Potash (K2O)	8%
Sulfur (S)	4.8%

Primary Plant Foods Derived from: Ammoniated Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea. Calcium Carbonate Equivalent per ton.


PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

WARNING

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.



BEST AVAILABLE COPY

498

(BACK PANEL)

HOMEOWNER USE ONLY

STATEMENT OF PRACTICAL TREATMENT

FIRST AID

IF IN EYES: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Call a physician.

IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: If breathing difficulty occurs, move individual to fresh air and get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply to water. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product has a preemergence herbicide which is applied to establish turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass, silver crabgrass or crowfoot, barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds. When applied as directed, the following established turfgrasses are tolerant:

- | | | | |
|--------------------|----------------|--------------|--------------------|
| Kentucky bluegrass | Centipedegrass | Zoysiagrass | St. Augustinegrass |
| Perennial ryegrass | Fescue | Bermudagrass | Bahiagrass |

BEST AVAILABLE COPY

518

(BACK PANEL)

HOMEOWNER USE ONLY

This product will not harm most nearby established ornamental trees and shrubs.

APPLICATION DIRECTIONS

Rates, Frequency of Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass
 - A. Northern (cool season) turfgrasses
For preemergence control, apply 3.2 to 4 pounds per thousand square feet in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 to 6 pounds per thousand square feet in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of this product from one half to the full recommended rate may be made 6 to 10 weeks after initial treatment for continued control.


2. Late spring, early summer germinating annual grasses - goosegrass. Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should, therefore, be timed accordingly.

BEST AVAILABLE COPY

(BACK PANEL)

HOMEOWNER USE ONLY

- A. Northern (cool season) turfgrass
For preemergence control, apply 4 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 to 6 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
3. Northern and southern turfgrasses
In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications from one half to the full recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.
4. Poa annua (annual bluegrass)
In areas of heavy infestation or annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.
- A. Northern (cool season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 6 pounds per thousand square feet in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application from 3 to 6 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.



BEST AVAILABLE COPY

798

(BACK PANEL)

HOMEOWNER USE ONLY

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization can be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

APPLICATION EQUIPMENT

Apply evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequency to be sure equipment is working properly and distributing uniformly.

SUGGESTED SPREADER SETTINGS

(Available at final printing)

USE PRECAUTIONS

Apply to the turf prior to the germination of weed grasses since this product will not control established weeds. This product may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 1.72 pounds per 1000 sq. ft. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application may thin the overseeded grasses. Because of varying cultural practices, this product should not be used on putting greens. Do not use on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.



BEST AVAILABLE COPY

(BACK PANEL)

HOMEOWNER USE ONLY

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of applications, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by: The Andersons
Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537

Date Printed: 01/24/94

Date Approved:

Supersedes:

Label Change:

