

PM 23

9198-79

4-30-98

1048



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number
233842

Application for Pesticide - Section I

1. Company/Product Number 9198-79	2. EPA Product Manager Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Tee Time Fertilizer with 1.15% Team	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Andersons, Lawn Fertilizer Division d/b/a Free Flow Fertilizer P.O. Box 119 Maumee, OH 43537 <input type="checkbox"/> Check here if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name

Section II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter	NOTIFICATION APR 30 1998
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - explain below	

Explanation: Use additional page(s) if necessary. (For section I and II).

Notification of Minor Label Change per PR Notice 95-2: Minor label changes: added optional marketing text and spreading information on both homeowner and commercial labels

This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling of the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes * <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes," Unit Package wgt.	No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
		If "Yes," Unit Package wgt.	No. per container	<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions <input type="checkbox"/> On label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in which Label Is Affixed To Product		<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled	<input type="checkbox"/> Other (_____)		

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Dawn L. Walters	Title Regulatory Administrator	Telephone No. (Include Area Code) (429) 891-2781
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		Date Application Received (Stamped)
2. Signature <i>Dawn L. Walters</i>	3. Title Regulatory Administrator	
4. Typed Name Dawn L. Walters	5. Date 4/17/98	

2 of 8

NOTIFICATION

APR 30 1998

9198-79C
Page 1 of 7

(FRONT PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass.

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-alpha,alpha,alpha-trifluoro-2,6-dinitro-p-toluidine 0.77%

Trifluralin: alpha,alpha,alpha-trifluoro-2,6-dinitro-N, N-dipropyl-p-toluidine 0.38%

INERT INGREDIENTS 98.85%

Total 100.00%

KEEP OUT OF THE REACH OF CHILDREN

CAUTION


SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 50 LBS. (22.7 kg)

TREATS UP TO 15,625 SQ.FT.

EPA Reg. No. 9198-79

EPA Est. 9198-OH-1



(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

NET WEIGHT 50 LBS. (22.7 kg)

TREATS UP TO 15,625 SQ.FT.

THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM

GUARANTEED ANALYSIS

Total Nitrogen (N)	25%
1.2% Ammoniacal Nitrogen	
17.3% Urea Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
Available Phosphate Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	8%
Sulfur (S)	4.8%
4.8% Free Sulfur(S)	

Primary Plant Foods Derived from: Ammonium Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

CAUTION

Causes eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust and contact with skin, eyes, or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt, and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation develops.

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation develops.

IF INHALED: If discomfort occurs, remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, provide cardiopulmonary resuscitation assistance (mouth-to-mouth) and call a physician.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product has a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass, silver crabgrass or crowfoot, barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds. When applied as directed, the following established turfgrasses are tolerant:

- | | | | |
|--------------------|----------------|--------------|--------------------|
| Kentucky bluegrass | Centipedegrass | Zoysiagrass | St. Augustinegrass |
| Perennial ryegrass | Fescue | Bermudagrass | Bahiagrass |


This product will not harm most nearby established ornamental trees and shrubs.

APPLICATION DIRECTIONS

Rates, Frequency of Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply approximately one to two weeks prior to the germination of annual grass weeds. At summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.



(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

- 1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass
 - A. Northern (cool season) turfgrasses
For preemergence control, apply 3 to 4 pounds per thousand square feet (1.5 to 2 pounds active ingredient per acre) in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 to 6 pounds per thousand square feet (2 to 3 pounds active ingredient per acre) in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of this product from one half to the full recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

- 2. Late spring, early summer germinating annual grasses - goosegrass
Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should, therefore, be timed accordingly.
 - A. Northern (cool season) turfgrass
For preemergence control, apply 4 pounds per thousand square feet (2 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 to 6 pounds per thousand square feet (2 to 3 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

- 3. Northern and southern turfgrasses
In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications from one half to the full recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

- 4. Poa annua (annual bluegrass)
In areas of heavy infestation or annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable



(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

- A. Northern (cool season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet (2 pounds active ingredient per acre) in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.
- B. Southern (warm season) turfgrasses
For preemergence control, apply 6 pounds per thousand square feet (3 pounds active ingredient per acre) in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application from 3 to 6 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization can be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.


APPLICATION EQUIPMENT

Apply evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly.

SUGGESTED SPREADER SETTINGS

(Available at final printing)

Important: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.



(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

USE PRECAUTIONS

Apply to the turf prior to the germination of weed grasses since this product will not control established weeds. This product may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 75 pounds per acre (1.5 pounds active ingredient). This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application may thin the overseeded grasses. Because of varying cultural practices, this product should not be used on putting greens. Do not use on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

MANUFACTURERS STATEMENT

This product conforms to the ingredient statement on the label. Manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage and handling of this product.

THIS PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED. IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND.

Manufactured by: The Andersons Lawn Fertilizer Division
d/b/a Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537



(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Optional Spreader Directions:

NOTE: These (or Tee Time) spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

It is important to follow the directions as indicated on this quality lawn care product as overapplication may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Refer to the table and picture below to determine the size of your lawn. Close spreader hopper and adjust the spreader setting using the suggested spreader settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn. Refer to "Path A" when using a broadcast spreader and "Path B" when using a drop spreader. Close the spreader hopper whenever spreader stops or turns. After application, water lawn thoroughly.

_____	Length	(Diagram of property)
x _____	Width	
= _____	Total Sq. Ft.	
- _____	Non-Lawn Areas	
= _____	Total Sq. Ft. Coverage You Need	

(Diagram of Path A)

(Diagram of Path B)

Optional Marketing Text:

Contains Team* for pre-emergence control of crabgrass and most other annual weed grasses.

Date Printed: 4/17/98
 Date Approved: 02/05/96
 Supersedes: 05/18/94
 Label Change: notification: correction of complete company name/add optional spreader directions
 Filename: 9198-79c.wpf

