

JAN - 7 2000

Ms. Dawn L. Walters
Regulatory Administrator
The Andersons, Inc.
Lawn Fertilizer Division
P.O. Box 112
Maumee, Ohio 43537

Dear Ms. Walters:

subject: The Andersons Tee Time Fertilizer with 1.15% Team[®]
EPA registration no. 9198-79

Your 6 month response to the Trifluralin Reregistration Eligibility Document (RED), Your Letter Dated December 2, 1999 and Amended Labeling In Response To Agency's Labeling Review and Letter Dated August 11, 1999

Your responses to the Trifluralin RED for the subject pesticide product registrations under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, have been reviewed. As these products contain two active ingredients and one of them, benefin, has not been issued an EPA RED; the subject products are presently not reregisterable. Under this circumstance this Agency requires that you revise these pesticide product labels to conform with the finding under the issued RED, in this case the Trifluralin RED. The proposed labeling has been reviewed and found acceptable under FIFRA, provided that you:

1. Revise the ingredient statement of the "Homeowner Use Only" labeling by including the respective CAS number for benefin and trifluralin.
2. On both the "Homeowner Use Only" and the Commercial Use Only" labels align the percent composition to the decimal place in the "Guaranteed Analysis" statement on page 2 of 6.
3. On the "Homeowner Use Only" label align to the decimal place the composition given in the ingredient statement.
4. Delete printer notes.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

-2-

section 5(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505C)

Enclosures (2)

E.wilson: Diskette ABC00: 01-07-00

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

(FRONT PANEL)
COMMERCIAL USE ONLY
(for internal use only, will not appear on final label)

THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM*

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipede grass, St. Augustine grass and Zoysiagrass.

ACTIVE INGREDIENTS:

Benefin: N-butyl-N-ethyl-alpha,alpha,alpha-trifluoro-2,6-dinitro-p-toluidine	0.77%
Trifluralin: alpha,alpha,alpha-trifluoro-2,6-dinitro-N, N-dipropyl-p-toluidine	0.38%
INERT INGREDIENTS	98.85%
Total	100.00%

*Trademark of Dow AgroSciences LLC

KEEP OUT OF THE REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 50 LBS. (22.7 kg)
TREATS UP TO 15,625 SQ.FT.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JAN - 7 2000
**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

9198-79

EPA Reg. No. 9198-79
EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code

(BACK PANEL)
COMMERCIAL USE ONLY
(for internal use only, will not appear on final label)

NET WEIGHT 50 LBS. (22.7 kg)
TREATS UP TO 15.625 SQ.FT.

THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM

GUARANTEED ANALYSIS

Total Nitrogen (N)	25%
1.2% Ammoniacal Nitrogen	
17.3% Urea Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
Available Phosphate Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	8%
Sulfur (S)	4.8%
4.8% Free Sulfur(S)	

Primary Plant Foods Derived from: Ammonium Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea.

PRECAUTIONARY STATEMENTS

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told by a poison control center or doctor.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 15 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • <i>If person is not breathing, call 911 or an ambulance, the give artificial respiration, preferably mouth to mouth, if possible.</i> • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning: Causes substantial but temporary eye irritation. Harmful if swallowed or absorbed through skin, or inhaled. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before reuse.

(BACK PANEL)
COMMERCIAL USE ONLY
(for internal use only, will not appear on final label)

Personal Protective Equipment (PPE)

Applicator and other handlers must wear:

- long-sleeved shirt and long pants
- socks and shoes
- chemical resistant gloves (such as water-proof)
- goggles or face shield

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations: Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Keep container closed and away from food, feedstuffs, and domestic water supplies.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This product has a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass, silver crabgrass or crowfoot, barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds. When applied as directed, the following established turfgrasses are tolerant:

Kentucky bluegrass	Centipedegrass	Zoysiagrass	St. Augustinegrass
Perennial ryegrass	Fescue	Bermudagrass	Bahiagrass

(BACK PANEL)
COMMERCIAL USE ONLY
(for internal use only, will not appear on final label)

Reseeding

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization can be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

Application Equipment

Apply evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequency to be sure equipment is working properly and distributing uniformly.

Suggested Spreader Settings

{Note: spreaders and settings vary due to market conditions and availability. Spreaders listed on final product label will be dependent on market availability and subregistrants.}

USE PRECAUTIONS AND RESTRICTIONS

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Apply to the turf prior to the germination of weed grasses since this product will not control established weeds. This product may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 75 pounds per acre (1.5 pounds active ingredient). This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application may thin the overseeded grasses. Because of varying cultural practices, this product should not be used on putting greens. Do not use on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

Entry Restrictions: Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

(BACK PANEL)
COMMERCIAL USE ONLY
(for internal use only, will not appear on final label)

APPLICATION INSTRUCTIONS

Rates, Frequency of Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass
 - A. Northern (cool season) turfgrasses
For preemergence control, apply 3 to 4 pounds per thousand square feet (1.5 to 2 pounds active ingredient per acre) in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 to 6 pounds per thousand square feet (2 to 3 pounds active ingredient per acre) in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of this product from one half to the full recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

2. Late spring, early summer germinating annual grasses - goosegrass
Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should, therefore, be timed accordingly.
 - A. Northern (cool season) turfgrass
For preemergence control, apply 4 pounds per thousand square feet (2 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 to 6 pounds per thousand square feet (2 to 3 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
3. Northern and southern turfgrasses

(BACK PANEL)
COMMERCIAL USE ONLY
(for internal use only, will not appear on final label)

In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications from one half to the full recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

4. Poa annua (annual bluegrass)

In areas of heavy infestation or annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

A. Northern (cool season) turfgrasses

For preemergence control, apply 4 pounds per thousand square feet (2 pounds active ingredient per acre) in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.

B. Southern (warm season) turfgrasses

For preemergence control, apply 6 pounds per thousand square feet (3 pounds active ingredient per acre) in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application from 3 to 6 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

(AP112085/RVred120199)

Manufactured by:
The Andersons Lawn Fertilizer Division
d/b/a Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537

Date Printed: 12/2/99
Date Approved:
Supersedes: 02/5/96
Label Change: RED requirements
Filename: 9198-79C.doc

9/14

(FRONT PANEL)
HOMEOWNER USE ONLY
(for internal use only, will not appear on final label)

THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM*

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass.

ACTIVE INGREDIENTS:

Benefin:.....	0.77%
Trifluralin:.....	0.38%
INERT INGREDIENTS	<u>98.85%</u>
Total	100.00%

*Trademark of Dow AgroSciences LLC

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JAN - 7 2000
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
9198-79

KEEP OUT OF THE REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 50 LBS. (22.7 kg)

TREATS UP TO 15,625 SQ.FT.

EPA Reg. No. 9198-79
EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code

(BACK PANEL)
HOMEOWNER USE ONLY
(for internal use only, will not appear on final label)

NET WEIGHT 50 LBS. (22.7 kg)
TREATS UP TO 15,625 SQ.FT.

THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM

GUARANTEED ANALYSIS

Total Nitrogen (N)	25%
1.2% Ammoniacal Nitrogen	
17.3% Urea Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
Available Phosphate (P2O5)	3%
Soluble Potash (K2O)	8%
Sulfur (S)	4.8%
4.8% Free Sulfur(S)	

Primary Plant Foods Derived from: Ammonium Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea.

PRECAUTIONARY STATEMENTS

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told by a poison control center or doctor.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 15 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning: Causes substantial but temporary eye irritation. Harmful if swallowed or absorbed through skin, or inhaled. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before reuse.

- **Personal Protective Equipment (PPE):** Wear goggles or face shield.

(BACK PANEL)
HOMEOWNER USE ONLY
(for internal use only, will not appear on final label)

User Safety Recommendations: Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
Pesticide Storage: Keep container closed and away from food, feedstuffs and domestic water supplies.
Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

GENERAL INFORMATION

This product has a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass, silver crabgrass or crowfoot, barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds. When applied as directed, the following established turfgrasses are tolerant:

- | | | | |
|--------------------|----------------|--------------|--------------------|
| Kentucky bluegrass | Centipedegrass | Zoysiagrass | St. Augustinegrass |
| Perennial ryegrass | Fescue | Bermudagrass | Bahiagrass |

Reseeding

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization can be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

Application Equipment

Apply evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequency to be sure equipment is working properly and distributing uniformly.

(BACK PANEL)
HOMEOWNER USE ONLY
(for internal use only, will not appear on final label)

Suggested Spreader Settings

(Available at final printing)

{Note: spreaders and settings vary due to market conditions and availability. Spreader listed on final product label will be dependent on market availability and subregistrants.}

GENERAL PRECAUTIONS AND RESTRICTIONS

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Apply to the turf prior to the germination of weed grasses since this product will not control established weeds. This product may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 1.72 pounds per 1000 sq. ft. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application may thin the overseeded grasses. Because of varying cultural practices, this product should not be used on putting greens. Do not use on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

Entry Restrictions: Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

APPLICATION INSTRUCTIONS

Rates, Frequency of Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass
 - A. Northern (cool season) turfgrasses
For preemergence control, apply 3 to 4 pounds per thousand square feet in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.
 - B. Southern (warm season) turfgrasses

(BACK PANEL)
HOMEOWNER USE ONLY
(for internal use only, will not appear on final label)

For preemergence control, apply 4 to 6 pounds per thousand square feet in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of this product from one half to the full recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

- 2. Late spring, early summer germinating annual grasses - goosegrass
Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should, therefore, be timed accordingly.
 - A. Northern (cool season) turfgrass
For preemergence control, apply 4 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 to 6 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

- 3. Northern and southern turfgrasses
In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications from one half to the full recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

- 4. Poa annua (annual bluegrass)
In areas of heavy infestation or annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.
 - A. Northern (cool season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 6 pounds per thousand square feet in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application from 3 to 6 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

(BACK PANEL)
HOMEOWNER USE ONLY
(for internal use only, will not appear on final label)

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.
(AP112085/Rvred12199)

Manufactured by:
The Andersons Lawn Fertilizer Division
d/b/a Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537

Date Printed: 12/1/99
Date Approved:
Supersedes: 02/05/96
Label Change: RED requirements
Filename: 9198-79H.doc