

9198-79

01/16/2008

1/8



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration

(under FIFRA, as amended)

EPA Reg. Number:

Date of Issuance:

9198-79

1-16-08

Term of Issuance:

Name of Pesticide Product:

The Andersons Tee Time  
Fertilizer with 1.15%  
TEAM

Name and Address of Registrant (include ZIP Code):

The Andersons, Inc.  
P.O. Box 119  
Maumee, Ohio 43537

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.


On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

**This product is conditionally reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:**

1. The order of the First Aid section should be changed to "If in Eyes, If Swallowed, If on Skin or on Clothing, and If Inhaled".
2. Per PR Notice 2000-1, a toll free telephone number for emergency medical information should be added to the label.
3. Change the Hazards to Humans and Domestic Animals statements to "Causes substantial but temporary eye injury. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated contact may cause allergic reactions in some individuals."
4. To the User Safety Recommendation the text in bold must be added "Users should remove clothing/**PPE**..."
5. To the Warranty section add "to the extent consistent with applicable law" in front of "no

1-16-08

other express" and "Manufacturer disclaims".	
<p>Signature of Approving Official:</p>  <p>Erik Kraft, Acting PM 23 Herbicide Branch, Registration Division (7505P)</p>	<p>Date:</p> <p>1-16-08</p>

**You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). A stamped copy of labeling is enclosed for your records.**

3/8

(FRONT PANEL)  
COMMERCIAL USE ONLY  
(for internal use only, will not appear on final label)

# THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM\*

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass.

*Optional marketing claims:*

- Combination granular fertilizer and Team\* preemergent herbicide for control of crabgrass and other annual weed grasses as listed.
- Consistently sized particles reduce segregation potential.
- Low nitrogen, high potassium formula for root development and improved drought tolerance.
- Team\* combines Balan\* and Treflan\* for effective preemergent control
- Combination fertilizer and Team\* preemergent herbicide
- Provides preemergent control of crabgrass and other listed annual grasses.
- High potassium formula
- Low nitrogen formula minimizes surge growth.

**ACTIVE INGREDIENTS:**

Benfen: N-butyl-N-ethyl-alpha,alpha,alpha-trifluoro-2,6-dinitro-p-toluidine ..... 0.77%  
 Trifluralin: alpha,alpha,alpha-trifluoro-2,6-dinitro-N, N-dipropyl-p-toluidine ..... 0.38%

**INERT INGREDIENTS** ..... 98.85%  
 Total ..... 100.00%

\*Trademark of Dow AgroSciences LLC

**KEEP OUT OF THE REACH OF CHILDREN**

**WARNING**

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS**

NET WEIGHT XX LBS. (XX.X kg)

*Optional:*

TREATS UP TO XX,XXX SQ.FT.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

1-16-08

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

9198-79

EPA Reg. No. 9198-79

EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code

4/8

(BACK PANEL)  
COMMERCIAL USE ONLY  
(for internal use only, will not appear on final label)

NET WEIGHT XX LBS. (XX.X kg)  
Optional:  
TREATS UP TO XX,XXX SQ.FT.

### THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM

<b>Guaranteed Analysis</b>	
Total Nitrogen (N) .....	XX%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%
Plant nutrients derived from: <i>For products being sold into Washington state:</i> Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.regulatory-info-ap.com/">http://www.regulatory-info-ap.com/</a>	

#### PRECAUTIONARY STATEMENTS

##### FIRST AID

<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told by a poison control center or doctor.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 15 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**Note to Physician:** Probable mucosal damage may contraindicate the use of gastric lavage.

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**Warning:** Causes substantial but temporary eye irritation. Harmful if swallowed or absorbed through skin, or inhaled. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before reuse.

#### Personal Protective Equipment (PPE)

All loaders, applicator and other handlers must wear:

- long-sleeved shirt and long pants
- socks and shoes
- chemical resistant gloves (such as water-proof)
- goggles or face shield

5/8

(BACK PANEL)  
COMMERCIAL USE ONLY  
*(for internal use only, will not appear on final label)*

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations:** Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**  
Do not contaminate water, food or feed by storage and disposal.  
**Pesticide Storage:** Keep container closed and away from food, feedstuffs, and domestic water supplies.  
**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

**GENERAL INFORMATION**

This product has a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass, silver crabgrass or crowfoot, barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds. When applied as directed, the following established turfgrasses are tolerant:

- |                    |                |              |                    |
|--------------------|----------------|--------------|--------------------|
| Kentucky bluegrass | Centipedegrass | Zoysiagrass  | St. Augustinegrass |
| Perennial ryegrass | Fescue         | Bermudagrass | Bahiagrass         |

**Reseeding**

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization can be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

6/8

(BACK PANEL)  
COMMERCIAL USE ONLY  
*(for internal use only, will not appear on final label)*

**Application Equipment**

Apply evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequency to be sure equipment is working properly and distributing uniformly.

**USE PRECAUTIONS AND RESTRICTIONS**

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Apply to the turf prior to the germination of weed grasses since this product will not control established weeds. This product may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 75 pounds per acre (1.5 pounds active ingredient). This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application may thin the overseeded grasses. Because of varying cultural practices, this product should not be used on putting greens. Do not use on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

**Entry Restrictions:** Do not enter or allow other people (or pets) to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow other people (or pets) to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow other people (or pets) to enter the treated area until the surface is dry after the watering-in.

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons (or pets), either directly or through drift. Only protected handlers may be in the area during application.

**APPLICATION INSTRUCTIONS**

**Rates, Frequency of Timing of Application**

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

This product must be watered in as soon as possible after application. Watering-in must be performed by the commercial applicator or the commercial applicator must provide the following instructions to the resident or owner in writing:

- This product must be watered in as soon as possible.
- Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

(BACK PANEL)  
COMMERCIAL USE ONLY  
(for internal use only, will not appear on final label)

**Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass**

*Northern (cool season) turfgrasses:* For preemergence control, apply 3 to 4 pounds per thousand square feet in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

*Southern (warm season) turfgrasses:* For preemergence control, apply 4 pounds per thousand square feet in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of this product from one half to the full recommended rate may be made 8 to 10 weeks after initial treatment for continued control. Do not exceed the Maximum Application Rates listed below.

**Late spring, early summer germinating annual grasses - goosegrass**

Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should, therefore, be timed accordingly.

*Northern (cool season) turfgrasses:* For preemergence control, apply 4 pounds per thousand square feet (2 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

*Southern (warm season) turfgrasses:* For preemergence control, apply 4 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

**Northern and southern turfgrasses**

In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications from one half to the full recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species. Do not exceed the Maximum Application Rates listed below.

***Poa annua* (annual bluegrass)**

In areas of heavy infestation or annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

*Northern (cool season) turfgrasses:* For preemergence control, apply 4 pounds per thousand square feet (2 pounds active ingredient per acre) in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.

*Southern (warm season) turfgrasses:* For preemergence control, apply 4 pounds per thousand square feet in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application from 3 to 4 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.

**Maximum Application Rates**

***Cool season turfgrasses:***

Maximum of 8.9 lbs. per 1,000 sq. ft. (390 lbs. product per acre, 3 lbs. benefin active ingredient per acre) per application.

Maximum of 2 applications per year.

Maximum of 17.9 lbs. per 1,000 sq. ft. (779 lbs. product per acre, 6 lbs. benefin active ingredient per acre) per year.

8/8

(BACK PANEL)  
COMMERCIAL USE ONLY  
(for internal use only, will not appear on final label)

**Warm season turfgrasses:**

Maximum of 4.5 lbs. per 1,000 sq. ft. (195 lbs. product per acre, 1.5 lbs. benefin active ingredient per acre) per application.

Maximum of 2 applications per year.

Maximum of 8.9 lbs. per 1,000 sq. ft. (390 lbs. product per acre, 3 lbs. benefin active ingredient per acre) per year.

**IMPORTANT: READ BEFORE USE:**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER and LIMITATION of LIABILITY:**

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS:**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVred100406)

Manufactured by:  
The Andersons Lawn Fertilizer Division  
d/b/a Free Flow Fertilizer  
P.O. Box 119  
Maumee, OH 43537

<b>Suggested Spreader Settings (Available at final printing)</b>				
These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.				
XX lbs. treats X,XXX sq. ft. (XX acre) at the 131 lbs. product/acre - LOW RATE (3.0 lbs./1,000 sq. ft.)				
XX lbs. treats X,XXX sq. ft. (XX acre) at the 174 lbs. product/acre - HIGH RATE (4.0 lbs./1,000 sq. ft.)				
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LOW RATE	HIGH RATE
Active ingredient/acre (benefin, trifluralin)			(1.0, 0.50)	(1.34, 0.66)
AA	X mph	XX ft/	X	X
BB	X mph	XX ft	X	X

Note to EPA: This list of spreaders and settings will vary dependent on market availability of spreaders

Date Printed	10/4/06
Date Approved	
Supersedes	6/17/03
Label change	Benefin RED
Filename	9198-79C
Label code	RVred100406