

9198-79

11/29/2011

1099



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

NOV 29 2011

Ms. Debbie Ziehr  
The Andersons, Inc.  
P.O. Box 119  
Maumee, Ohio 43537

Subject: Notification: Revised Environmental Hazards per PR Notice 2008-1  
The Andersons Tee Time Fertilizer with 1.15% Team  
EPA Reg. No. 9198-79  
Your Application Dated September 26, 2011, as Amended by Email November 29, 2011

Dear Ms. Ziehr:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2008-1 for the subject product. The Registration Division (RD) has conducted a review of this request for applicability under PRN 2008-1 and finds that the label changes requested fall within the scope of the PR Notice. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at (703) 305-1243 or Susan Stanton of my staff at (703) 305-5218.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Program

2049

Please read instructions on reverse before completing form.

Form Approved OMB No. 2070-0080, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 9198-79	2. EPA Product Manager Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Tee Time Fertilizer with 1.15% Team	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div. Inc. P.O. Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

**NOTIFICATION**  
NOV 28 2011

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of Label change per PR Notice 98-10: See cover letter for details.

This notification is consistent with the guidance of PR Notice 98-10 and the requirements of EPA's regulations at 40 CFR 153.10 and 40 CFR 152.46 and not other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 98-10 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Plastic	
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 25 - 1,000 lbs.		5. Location of Label Directions <input checked="" type="checkbox"/> On label	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Debbie Ziehr	Title Regulatory Administrator	Telephone No. (Include Area Code) 419-891-6671
2. Signature 		6. Date Application Received (Stamped)
3. Title Regulatory Administrator		
4. Typed Name Debbie Ziehr	5. Date 9/26/11	

**Certification**  
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

(FRONT PANEL)  
HOMEOWNER USE ONLY  
*(for internal use only, will not appear on final label)*

# THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM\*

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass.

### ACTIVE INGREDIENTS:

Benefin	CAS No. 1861-40-1	0.77%
Trifluralin	CAS No. 1582-09-8	0.38%
<b>INERT INGREDIENTS</b>		<u>98.85%</u>
Total		100.00%

**NOTIFICATION**  
**NOV 29 2011**

\*Trademark of Dow AgroSciences LLC

**KEEP OUT OF THE REACH OF CHILDREN**

### WARNING

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS**

NET WEIGHT XX LBS. (22.7 kg)  
TREATS UP TO XX,XXX SQ.FT.

- Controls smooth & hairy crabgrass, goosegrass, barnyardgrass & yellow & green foxtail
- Extended greening from controlled release nitrogen
- Encourages quick spring greenup
- For thick, green, crabgrass-free lawns
- Controls crabgrass and other annual grasses all season long
- Formulated for regions with naturally high phosphate levels
- XX Nitrogen – essential for plant growth and a thick, green lawn
- XX Phosphorous – Phosphorous free
- XX Potassium – to help grass resist the elements and stay strong
- Prevents crabgrass and other annual weed grasses as listed
- Provides your lawn with controlled-release nitrogen for extended feeding
- Early Spring application
- Apply with a broadcast or drop spreader
- Eliminates crabgrass while fertilizing your lawn
- Controlled release nitrogen provides for extended feeding

EPA Reg. No. 9198-79  
EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A  
Underlined letter is first letter used in run code

(BACK PANEL)  
HOMEOWNER USE ONLY  
(for internal use only, will not appear on final label)

### THE ANDERSONS TEE TIME FERTILIZER WITH 1.15% TEAM

<b>Guaranteed Analysis</b>	
Total Nitrogen (N) .....	XX%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%
Plant nutrients derived from:	
<i>For products being sold into Washington state:</i>	
Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.regulatory-info-ap.com/">http://www.regulatory-info-ap.com/</a>	

*For products being sold into Kentucky:*

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.) pounds of nitrogen per 1,000 square feet (Optional P2O5 and K2O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

#### PRECAUTIONARY STATEMENTS

##### FIRST AID

<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told by a poison control center or doctor.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 15 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**Note to Physician:** Probable mucosal damage may contraindicate the use of gastric lavage.

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**Warning:** Causes substantial but temporary eye irritation. Harmful if swallowed or absorbed through skin, or inhaled. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash before reuse.

- **Personal Protective Equipment (PPE):** Wear goggles or face shield.

**User Safety Recommendations:** Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. ~~Do not apply directly to water.~~ Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. ~~Do not contaminate water when disposing of equipment washwaters.~~ To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**  
Do not contaminate water, food, or feed by storage and disposal.  
**Pesticide Storage:** Keep container closed and away from food, feedstuffs and domestic water supplies.  
**DISPOSAL:**  
If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.  
If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**GENERAL INFORMATION**

This product has a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (Poa annua), smooth and hairy crabgrass, goosegrass, silver crabgrass or crowfoot, barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds. When applied as directed, the following established turfgrasses are tolerant:

- |                    |                |              |                    |
|--------------------|----------------|--------------|--------------------|
| Kentucky bluegrass | Centipedegrass | Zoysiagrass  | St. Augustinegrass |
| Perennial ryegrass | Fescue         | Bermudagrass | Bahiagrass         |

**Reseeding**

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization can be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

**Application Equipment**

Apply evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly.

**Suggested Spreader Settings**

(Available at final printing)

Note: It is important to follow the directions on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness or area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. Do not hand spread.

**Broadcast Spreader Settings**

{diagram}{diagram}

For best results, apply in a circular path starting on the outside and working toward the center (see Broadcast Spreader Path diagram at left). A little overlap of product will ensure complete coverage.

	3.2# Rate
Earthway 2000A	10 - 11
Precision SB50	7 1/2 - 8 1/2
Republic EZ540/74540	2 1/2
Scotts Speedygreen	4 1/2
Scotts Easygreen	27 - 28

**Drop Spreader Settings**

{diagram}{diagram}

First apply 2 border strips along the edge of each end of the lawn (see Drop Spreader Path diagram at left). Then apply product along the length of the lawn, overlapping slightly and shutting of the spreader before turning as you reach the edge of the border strips.

	3.2# Rate
Earthway 7350	4 1/2 - 5 1/2
Precision FH22	8 1/2 - 9
Republic EZ530/75350	5 1/2 - 8 1/2
Scotts PF4	5 - 6
Scotts Accugreen	5 1/2 - 6

{Note: spreaders and settings vary due to market conditions and availability. Spreaders listed on final product label will be dependent on market availability and subregistrants.}

**GENERAL PRECAUTIONS AND RESTRICTIONS**

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Apply to the turf prior to the germination of weed grasses since this product will not control established weeds. This product may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 1.72 pounds per 1000 sq. ft. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application may thin the overseeded grasses. Because of varying cultural practices, this product should not be used on putting greens. Do not use on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

**Entry Restrictions:** Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

**Application Restrictions:** Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

## APPLICATION INSTRUCTIONS

### Rates, Frequency of Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass
  - A. Northern (cool season) turfgrasses  
For preemergence control, apply 3 to 4 pounds per thousand square feet in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.
  - B. Southern (warm season) turfgrasses  
For preemergence control, apply 4 to 6 pounds per thousand square feet in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of this product from one half to the full recommended rate may be made 8 to 10 weeks after initial treatment for continued control.
2. Late spring, early summer germinating annual grasses - goosegrass  
Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should, therefore, be timed accordingly.
  - A. Northern (cool season) turfgrass  
For preemergence control, apply 4 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
  - B. Southern (warm season) turfgrasses  
For preemergence control, apply 4 to 6 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
3. Northern and southern turfgrasses  
In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications from one half to the full recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.
4. *Poa annua* (annual bluegrass)  
In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

- A. Northern (cool season) turfgrasses  
For preemergence control, apply 4 pounds per thousand square feet in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.
- B. Southern (warm season) turfgrasses  
For preemergence control, apply 6 pounds per thousand square feet in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application from 3 to 6 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.

**IMPORTANT: READ BEFORE USE:**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER and LIMITATION of LIABILITY:**

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS:**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Visit our website at [www.andersonsstore.com](http://www.andersonsstore.com)

Manufactured by:  
The Andersons Lawn Fertilizer Division  
d/b/a Free Flow Fertilizer  
P.O. Box 119  
Maumee, OH 43537

Optional marketing information:  
(see next page)

Date Printed	09/23/11
Date Approved	1/16/08
Supersedes	1/7/00
Label change	Notification - Environment Hazards
Filename	9198-79H.doc
Label code	RVnot092311





# 4-Season Cycle

## Fertilizer Application Guide

### U.S. Climate Zones:



**Turf Varieties by Zone:**  
 ■ Bluegrass, Ryegrass, Fine & Tall Fescue  
 ■ Bermudagrass, Zoysiagrass  
 ■ St. Augustine  
 ■ Centipedegrass, Zoysiagrass, Carpetgrass, Bahiagrass, St. Augustine  
 Abnormal climatic conditions can affect appropriate timing.

Turf Zone & Variety	February	March	April	May	June	July	August	September	October	November	December
<b>NORTHERN ZONE</b> Cool Season Grasses		29-0-3 Crabgrass Preventer		29-0-3 Weed & Feed	29-0-3 Weed & Feed	25-0-5 Fertilizer + Insect Control	25-0-5 Fertilizer + Insect Control	24-0-12 Winterizer			
<b>TRANSITIONAL ZONE</b>		27-0-5 Crabgrass Preventer		29-0-3 Weed & Feed	29-0-3 Weed & Feed	25-0-5 Fertilizer + Insect Control	25-0-5 Fertilizer + Insect Control	24-0-12 Winterizer			
<b>SOUTHERN ZONE</b> Warm Season Grasses		27-0-5 Crabgrass Preventer		29-0-3 Weed & Feed	29-0-3 Weed & Feed	25-0-5 Fertilizer + Insect Control	25-0-5 Fertilizer + Insect Control	24-0-12 Winterizer			

  

<b>Early Spring</b> Crabgrass Preventer + Lawn Fertilizer 27-0-5	<b>Late Spring</b> Weed & Feed + Lawn Fertilizer 29-0-3	<b>Summer</b> Insect Control + Lawn Fertilizer 25-0-5	<b>Fall</b> Winterizer + Lawn Fertilizer 24-0-12	<b>Four Seasons</b> Lawn Fertilizer 30-0-3
--	---	---	--	--