

1 of 4
pm25
9198-81



Tee TimeTM Fertilizer with 1.15 Balan[®] 20-4-10

RECEIVED
JAN 28 1987
9198-81

GUARANTEED ANALYSIS

Total Nitrogen (N).....	20%
1.1% Ammoniacal Nitrogen	
17.9% Urea Nitrogen	
8.0% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	10%
Sulfur (S).....	4.5%

Primary Plant Foods Derived from: Ammoniated Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea. Sulfur derived from Sulfur Coated Urea.
Potential Acidity 653 pounds Calcium Carbonate Equivalent per ton.

ACTIVE INGREDIENT:	
N-butyl N-ethyl-s,s,a-trifluoro-2,6-dinitro-p-toluidine.....	1.15%
INERT INGREDIENTS: (relative to herbicidal content).....	98.85%
TOTAL.....	100.00%

¹Balan[®] —the registered trademark of Elanco Products Co.

2614

the professional's
partner™




KEEP OUT OF THE REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Net Weight 50 Lbs. (22.75 kg)
Treats Up to 16,700 Sq. Ft.

EPA Reg. No. 9198-81

EPA Est. No. 9198-OH-1



Net Weight 50 Lbs. (22.75 kg)
Treats Up to 16,700 Sq. Ft.

Tee Time™ Fertilizer with 1.15 Balan®

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
Hazards To Humans and Domestic Animals

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly after handling. If inhaled: If breathing discomfort occurs, move to fresh air. Get medical attention if breathing difficulty continues.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 2 glasses water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

IF IN EYES: Flush eyes with plenty of water and get prompt medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store in original container only. In case of spill contain material and dispose as waste. Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Tee Time Fertilizer with 1.15 Balan is a selective pre-emergence herbicide for the control of crabgrass and most other annual weed grasses on lawns, golf courses and ornamental turfs in established.

Bluegrass (perennial)	Fescue	Bermuda grass	St. Augustine
Ryegrass (perennial)	Bahia grass	Centipede grass	Zoysia grass

This product controls the listed annual weeds by killing their seeds as they germinate. Apply Tee Time Fertilizer with 1.15 Balan before these weeds germinate since it does not control established weeds. Successful pre-emergence control of the annual weeds listed requires proper timing of application. All annual grasses do not germinate at the same time. Time of germination may also vary from year to year because of soil temperature and rainfall.

- FOR OPTIMUM RESULTS**
- Any cultural practices which disturb the soil such as aeration or verticillaging should be performed prior to application of this product.
 - Apply Tee Time Fertilizer with 1.15 Balan, one to two weeks prior to the anticipated germination of the targeted weed species based on growing conditions for the particular year.
 - In those locations where both spring germinating annual grasses (crabgrasses, foxtails, barnyard grasses) and summer germinating annual grasses (primarily goosegrass) are anticipated, when the normal period of germination is extended because of favorable climatic conditions, in southern turf grass areas with longer growing seasons, or when cultural practices require application of Tee Time Fertilizer with 1.15 Balan well in advance of weed seed germination dates, two applications at the recommended rate, spaced 8 to 10 weeks apart may be required for season long control.
 - Apply with equipment designed for application of granular pesticides. Calibrate the applicator prior to use according to manufacturer's instructions. Check frequently to applicator prior to use according to manufacturer's instructions. Check frequently to be sure equipment is working properly and distributing granules uniformly.
 - Uniform application is essential for satisfactory weed control. A more uniform application can be made by spreading half the required amount over the area and then application, water thoroughly. This assures more even distribution of the herbicide and avoids gaps in coverage.
 - A well maintained turf with low weed infestations will require less herbicide for effective control than will a thin turf with low fertility, minimum maintenance and heavy traffic. Tee Time Fertilizer with 1.15 Balan should be part of an overall turf maintenance program.

APPLICATION RATES:

40F4

Northern (Cool Season) Turf grasses	Lbs/A	Lbs/M	Lbs a/A	Anticipated Days of Control
Spring germinating grassy weeds	130	3	1.5	60 90
(crabgrass, foxtails, barnyard grass)	174	4	2.0	60 120
Early summer germinating grassy weeds (goosegrass)	174	4	2.0	60 100
Spring and summer germinating grassy weeds	174	4	2.0 followed by up to 2.0 in a second application 8 to 10 weeks after 1st application	120 150

Southern (warm season) Turf Grasses	Lbs/A	Lbs/M	Lbs a/A	Anticipated Days of Control
Spring germinating grassy weeds	174 to 261	4 to 6	2 to 3	60 120
Early summer germinating grassy weeds	261	6	3	60 100
Spring and summer germinating grassy weeds	174 to 261	4 to 6	2 to 3 followed by up to 3 in a second application 8 to 10 weeks after 1st application	120 160

Special instructions for annual bluegrass (Poa Annua) control:

	Lbs/A	Lbs/M	Lbs a/A
Northern Turf Apply	174	4	2.0 in early fall followed by 2.0 applied early spring
	174	4	
	Lbs/A	Lbs/M	Lbs a/A
Southern Turf Apply	260	6	3.0 in late summer followed by 3.0 10 to 12 weeks later
	260	6	

RESEEDING/SPRIGGING

Reseeding should be delayed for at least 6 weeks after applying Tee Time Fertilizer with 1.15 Balan. When using at the highest rates, reseeding should be delayed such as irrigation and fertilization be followed. For best results, the amount of seed used should be increased and care should be taken to insure contact of the seed with the soil.

This product should not be applied in the spring to turf grasses planted the previous fall. Do not apply to newly sprigged areas of Bermuda Grass, St. Augustine Grass or Centipede Grass until these turfs are well established. In Bermuda Grass areas that have been over seeded with winter grasses, a spring application of Tee Time Fertilizer with 1.15 Balan may thin the over seeded grasses.

USE PRECAUTIONS

This product may thin established annual bluegrass (Poa Annua) turf and fine-leaved fescues at rates above 1.5 pounds of active ingredient per acre. Do not apply to putting greens or Dichondra.

Manufactured by

the professional's partner™



Lawn Fertilizer Division
 P.O. Box 119
 Maumee, Ohio 43537