

Tee Time

Contains
1.5% Oftanol[®]
Insecticide

the professional's
partner

The
Andersons

ACTIVE INGREDIENT:

1-Methylethyl 2-(ethoxy((1-methylethyl)amino)phosphinothioylloxy)benzoate 1.5%

INERT INGREDIENTS 98.5%
100.0%

"OFTANOL is a registered trademark of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen."
U.S. Patent No. 3,621,082
Canadian Patent No. 825,198

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR STATEMENTS OF PRACTICAL TREATMENT AND OTHER PRECAUTIONARY STATEMENTS

NET WEIGHT 25 lbs. (11.3 kg)
TREATS up to 16,650 Sq. Ft.

ACCEPTED
JUL 8 1986
USE OF THIS PRODUCT IS LIMITED TO THE CONTROL OF PESTS OF ORNAMENTAL PLANTS AND HOUSE PLANTS.
EPA Reg. No. 9196-08

ANDERSONS TEE-TIME WITH 1.5% OFTANOL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or the Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush with plenty of water. IF ON SKIN: Wash with soap and hot water.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned stay out of smoke.

Store in a cool, dry place in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Do not walk through spilled material. Dispose of pesticide as directed above.

ANDERSONS TEE TIME WITH 1.5% OFTANOL

SPREADER SETTINGS

Approximate spreader settings to apply OFTANOL 1.5% Granular insecticide at the required rate are listed below. They are calculated to deliver the correct amount based on new equipment and ideal conditions. This serves as a guide only. Calibration of each spreader is necessary to apply an accurate rate.

SUGGESTED SPREADER SETTINGS

<u>Spreader</u>	<u>Speed</u>	<u>Spread Width</u>	<u>1.5 lbs./1,000 sq. ft.</u>	<u>3 lbs./1,000 sq. ft.</u>
Cyclone	3 MPH	10-12 ft.	4-4½	4½-4 ¾
Scotts Broadcast	3 MPH	10-12 ft.	B½-D	D½-F
Scotts Drop	-	-	5-5 ¾	6½-7
Lely	5 MPH	39 ft.	3	4

RECOMMENDED APPLICATIONS ON LAWNS

SOIL INSECTS:

Apply OFTANOL 1.5% Granular at the rate of 3 lbs. per 1,000 sq. ft. for the following soil insects: Whitegrub larvae (such as Japanese Beetle, Black Turfgrass Ataenius, Chafers, Phyllophaga spp., Green June Beetle, Oriental Beetle, Asiatic Garden Beetle) and Mole Crickets. Larval stage of Billbug and Hyperodes Weevil.

Be sure to read "APPLICATION EQUIPMENT" section of this label. Do not apply more than 6 lbs. of OFTANOL 1.5% Granular per 1,000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1,000 sq. ft. regardless of formulation used. Minimum interval between applications must be 30 days.

For White Grub and Billbug Larval Control: Water turf area thoroughly within 24 hours of application, using sufficient water to wet soil 1 to 1½ inches deep.

For Hyperodes Weevil Control: Apply between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15, and water-in as for white grub larval control.

For Mole Cricket Control: Water immediately after application using ½ to 1 inch of water

Consult your local, state Agricultural Experiment Station, or State Extension Service Specialist for more specific information regarding timing of application.

ANDERSONS TEE-TIME WITH 1.5% OFTANOL

DIRECTIONS FOR USE

It is in violation of the Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use and Conditions of Sale before using OFTANOL 1.5% Granular insecticide.

CONDITION OF SALE: The directions on this label were determined through research to be the directions for correct use of this product. This product has been tested for a range of weather conditions similar to those weather conditions that are ordinary and customary in the geographic area where the product is used. Insufficient control of pests may result from the occurrence of extraordinary or unusual weather, or from failure to follow label directions. In addition, failure to follow label directions may cause injury to the crop to which the product is applied, other crops, animals, man or the environment. The Andersons and the seller offer, and the buyer accepts and uses this product subject to the conditions that extraordinary or unusual weather, or failure to follow label directions are beyond the control of The Andersons and are, therefore, the responsibility of the buyer. Oftanol 1.5% granular insecticide is recommended for use on lawns, and other ornamental and recreational turf grass areas.

APPLICATION EQUIPMENT

Apply OFTANOL 1.5% Granular uniformly over the turf area with suitable granular turf application equipment. Both drop-type and rotary-type spreaders may be used to apply this product. Avoid the use of spreaders which would apply the material in narrow concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

APPLICATION TECHNIQUES

OFTANOL 1.5% Granular application using a pull-type drop spreader can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount over the area, and then immediately apply the remaining half at right angles or crosswise. The full recommended rate must be used for maximum benefit. When applying OFTANOL 1.5% Granular, use appropriate marking systems and make application when the wind is calm to prevent missed areas.

ANDERSONS TEE-TIME WITH 1.5% OFTANOL

FOR FIRE ANTS:

Area Fire Ant Control: Apply 3 lbs./1,000 sq. ft. uniformly over area to be treated for residual control. Water immediately after application using $\frac{1}{2}$ to 1 inch of water. If established fire and mounds are present use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

Established Fire Ant Mounds: Drench Treatments: Wet Conditions:
Distribute 0.5 oz. evenly over 9 sq. ft. and immediately sprinkle gently with 3 gallons of water.

Dry Conditions: Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one mound drench treatment.

SURFACE INSECTS:

For Chinch Bugs and Sod Webworm Larvae: Make application at the rate of $1\frac{1}{2}$ or 3 pounds per 1,000 sq. ft.. Apply $\frac{1}{2}$ to 1 inch of water within 24 hours of treatment. Re-apply as necessary for pest re-infestations. Do not apply more than 6 lbs. per 1,000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1,000 sq. ft. regardless of formulation used. Minimum interval between applications is 15 days for 1.5 lb. rate and 30 days for the 3 lb. rate.

Consult your local turf, State Agricultural Experiment Station, or State Extension Service Specialist for more specific information regarding timing of application.

RESTRICTIONS

Do not mow treated turf area before rainfall or irrigation has been applied as per RECOMMENDED APPLICATIONS. Do not graze treated areas or use clippings from treated areas for feed or forage. Keep children and pets off treated area until dry.

Manufactured by:

The Andersons
Maumee, Ohio 43537