

Tee Time™

1.5% **Oftanol**
insecticide

the professional's
partner™

The Andersons

ACTIVE INGREDIENT:
1-Methylethyl 2-((ethoxy((1-
methylethyl)amino)phosphino-
thio)loxy)benzoate

15%

INERT INGREDIENTS

98.5%
100.0%

FOR COMMERCIAL USE ONLY

"OFTANOL is a registered trademark of the
Parent Company of Farbenfabriken Bayer
GmbH, Leverkusen"

US Patent No. 3,621,082
Canadian Patent No. 825,198

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Statements of Practical Treatment
and Other Precautionary Statements

Net Weight 25 Lbs. (11.3 Kg)
Treats Up To 16,650 Sq. Ft.

EPA Reg. No. 9198-88

EPA Est. No. 9198-OH 1

ACCEPTED

DEC 10 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 9198-88

1.5% Olfanol

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful, if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of the reach of children.

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
IF IN EYES: Flush with plenty of water. **IF ON SKIN:** Wash with soap and hot water.
PHYSICIAN INFORMATION: 1.5% Olfanol is a cholinesterase inhibitor. Atropine and 2-PAM have been shown to be effective antidotes.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.
Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children.
Handle and open containers in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Do not wash through spilled material. Dispose of pesticide as directed above.

DIRECTIONS FOR USE

It is a violation of the Federal law to use this product in a manner inconsistent with its labeling.
IMPORTANT: Read these entire Directions for Use and Conditions of Sale before using OLFANOL 1.5% Granular Insecticide.
CONDITIONS OF SALE: The directions on this label were determined through research to be the directions for correct use of this product. This product has been tested for a range of weather conditions similar to those weather conditions that are ordinary and customary in the geographic area where the product is used. Insufficient control of pests may result from the occurrence of extraordinary or unusual weather or from failure to follow label directions. In addition, failure to follow label directions may cause injury to the crop to which the product is applied, other crops, animals, man or the environment. The Andersons and the seller offer and the buyer accepts and uses this product subject to the conditions that extraordinary or unusual weather, or failure to follow label directions are beyond the control of The Andersons and are, therefore, the responsibility of the buyer.
OLFANOL 1.5% Granular Insecticide is recommended for use on lawns, and other ornamental and recreational turf grass areas.

APPLICATION EQUIPMENT

Apply OLFANOL 1.5% Granular uniformly over the turf area with suitable granular turf application equipment. Both drop-type and rotary-type spreaders may be used to apply OLFANOL 1.5% Granular. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

APPLICATION TECHNIQUES

OLFANOL 1.5% Granular application using a pull-type drop spreader can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount of OLFANOL 1.5% Granular over the area, and then immediately apply the remaining half at right angles or crosswise. The full recommended rate of OLFANOL 1.5% Granular must be used for maximum benefit.

When applying OLFANOL 1.5% Granular, use appropriate marking systems and make application when the wind is calm to prevent missed areas.

SPREADER SETTINGS

Approximate spreader settings to apply OLFANOL 1.5% Granular insecticide at the required rate are listed below. They are calculated to deliver the correct amount based on new equipment and ideal conditions. This serves as a guide only. Calibration of each spreader is necessary to apply an accurate rate.

SUGGESTED SPREADER SETTINGS

Spreader	Speed	Spread Width	Settings	
			15 Lbs / 1000 Sq Ft	3 Lbs / 1000 Sq Ft
Cycloze B1-84	3 MPH	10-12 ft	4-4 1/2	4 1/2-4 1/2
Scotts Broadcast R-7X	3 MPH	10-12 ft	8 1/2-9	8 1/2-9
Scotts Drop PF2	—	—	5-6 1/2	6 1/2-7
Lily Type W	5 MPH	36 ft	3	4

RECOMMENDED APPLICATIONS ON LAWNS

SOIL INSECTS:

Apply OLFANOL 1.5% Granular at the rate of 3 lbs. per 1000 sq. ft. for the following soil insects: White grub larvae (such as Japanese Beetle, Black Turfgrass Ateuchus, Chalcids, Phyllophaga spp., Green June Beetle, Oriental Beetle, Asiatic Garden Beetle) and Mole Crickets. Larval stage of Billbug and Hyperandrus Wood.

Be Sure to Read "APPLICATION EQUIPMENT" Section of this label. Do not apply more than 6 lbs. of OLFANOL 1.5% Granular per 1000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1000 sq. ft. regardless of formulation used. Minimum interval between applications must be 30 days.

For White Grubs and Billbug Larval Control: Water turf area thoroughly within 24 hours of application, using sufficient water to wet soil 1 to 1 1/2 inches deep.

For Hyperandrus Wood Control: Apply between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15, and water-in as for white grub larval control.

For Mole Cricket Control: Water immediately after application using 1/2 to 1 inch of water.

Consult your local turf, state Agricultural Experiment Station, or state Extension Service Specialists for more specific information regarding timing of application.

FOR FIRE ANTS:

Area Fire Ant Control: Apply 3 lbs. / 1000 sq. ft. uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

Established Fire Ant Mounds, Drench Treatments, Wet Conditions: Distribute 0.5 oz. evenly over 8 sq. ft. and immediately sprinkle gently with 3 gallons of water.

Dry Conditions: Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment.

Make only one mound drench treatment.

SURFACE INSECTS:

For CRICKET BUGS and Sod Webworm Larvae: Make application at the rate of 1 1/2 or 2 pounds per 1000 sq. ft. Apply 1/2 to 1 inch of water within 24 hours of treatment. Reapply as necessary for pest re-infestations. Do not apply more than 6 lbs. of OLFANOL 1.5% Granular per 1000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1000 sq. ft. regardless of formulation used. Minimum interval between applications is 15 days for 1.5 lb. rate and 30 days for the 3 lb. rate.

Consult your local turf, state Agricultural Experiment Station, or state Extension Service Specialists for more specific information regarding timing of application.

RESTRICTIONS

Do not mow treated turf area before rainfall or irrigation has been applied as per RECOMMENDED APPLICATIONS. Do not graze treated areas or use clippings from treated areas for 14 or longer. Keep children and pets off treated area until dry.

Manufactured by
**the professional's
partner™**



Lawn Fertilizer Division
P.O. Box 119

Wapakoneta, OH 45389

© Copyright 1983, 1 A