

FEB 8 1991

S390107

The Andersons Lawn & Fertilizer Division
P.O. Box 119
Maumee, OH 43537

Gentlemen:

Subject: Anderson's Tee-Time With 1.5% Oftanol
EPA Registration No. 9198-88
Your Labeling Submitted May 1, 1990

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the labeling change listed below before you release the product for shipment bearing the revised labeling.
 - The statement "For Commercial Use Only" appearing on the front panel conflicts with the statement "For Homeowner Use Only" appearing under the "Storage and Disposal" heading on the back panel. This conflict must be resolved by deleting one of these statements.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

64972:I:Mautz:L16-8:KEVRIC:02/07/91:03/05/91:CL:WO:DD

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

MAM
Marilyn A. Mants
Acting Product Manager (16)
Insecticide-Rodenticide Branch
Registration Division (E7505C)

Enclosure

THE ANDERSONS

FEB 8 1991

TEE TIME

1.5% OFTANOLR

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

9198-88

ACTIVE INGREDIENTS:

1-Methylethyl 2-((ethoxy((1-methylethyl)amino)phosphorothioyl)oxy)benzoate	1.5%
INERT INGREDIENTS	98.5%
TOTAL	100.0%

OFTANOLR is a Reg. TM of the Patent Company of Farbwerke Bayer GmbH, Leverkusen

U.S. Patent No. 3,621,082
Canadian Patent No. 825,198

FOR COMMERCIAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

CAUTION


SEE BACK PANEL FOR STATEMENT OF PRACTICAL TREATMENT AND OTHER PRECAUTIONARY STATEMENTS

NET WEIGHT 25 LBS. (11.3 Kg)

TREATS UP TO 16,650 SQ.FT.

EPA REG. NO. 9198-88

EPA EST. NO. 9198-OH-1



(BACK PANEL)

THE ANDERSONS

TEE TIME

1.5% OFTANOL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Give a glass or two of water, and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If in eyes: flush with plenty of water. If on skin: wash with soap and hot water.

If inhaled: remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS


This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use before using The Andersons Tee Time with 1.5% Oftanol.

OFTANOL 1.5% Granular Insecticide is recommended for use on lawns and other ornamental and recreational turf grass areas.



APPLICATION EQUIPMENT

Apply 1.5% Granular uniformly over the turf area with suitable granular turf application equipment. Both drop-type and rotary-type spreaders may be used to apply OFTANOL 5% Granular. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

APPLICATION TECHNIQUES

OFTANOL 1.5% Granular application using a pull-type drop spreader can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount of OFTANOL 1.5% Granular over the area, and then immediately apply the remaining half at right angles or crosswise. The full recommended rate of OFTANOL 1.5% Granular must be used for maximum benefit.

When applying OFTANOL 1.5% Granular, use appropriate marking systems and make application when the wind is calm to prevent missed areas.

SPREADER SETTINGS

Approximate spreader setting to apply Tee Time Fertilizer with OFTANOL at the required rate are listed below. They are calculated to deliver the correct amount based on new equipment and ideal conditions. This serves as a guide only. Calibration of each spreader is necessary to apply an accurate rate.

SUGGESTED SPREADER SETTINGS

(Available at final printing)


RECOMMENDED APPLICATIONS ON LAWNS

SOIL INSECTS:

Apply OFTANOL 1.5% Granular at the rate of 3 lbs. per 1,000 sq.ft. for the following insects: Whitegrub larvae (such as Japanese Beetle, Black Turfgrass Attenius, Chafers, Phyllophaga spp., Green June Beetle, Oriental Beetle, Asiatic Garden Beetle) and Mo. Crickets, Larval stage of Billbug and Hyperodes Weevil.

Be Sure to Read APPLICATION EQUIPMENT Section of This Label. Do not apply more than 6 lbs. of OFTANOL 1.5% Granular per 1,000 sq.ft. per year or an equivalent amount of 1.45 oz. active ingredient per 1,000 sq.ft. regardless of formulation used. Minimum interval between applications must be 30 days.

For White Grub and Billbug Larvae Control: Water turf area thoroughly within 24 hours of application, using sufficient water to wet soil to 1/2 inches deep.



Hyperodes Weevil Control: Apply between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15 and water in as for preventative White Grub control.

For Mole Cricket Control: Water immediately after application using 1/2 to 1 inch water.

Consult your local turf, state Agricultural Experiment Station, or Extension Service Specialists for more specific information regarding timing of application.

FOR FIRE ANTS:

Area Fire Ant Control: Apply 3 lbs./1,000 sq.ft. uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present, use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

Established Fire Ant Mounds - Drench Treatments: Wet Conditions: Distribute 0.5 oz. evenly over 9 sq.ft. and immediately sprinkle gently with 3 gallons of water. Dry Conditions: Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one Mound Drench Treatment.


SURFACE INSECTS:

For CHINCH BUGS and Sod Webworm Larvae: Make application at the rate of 1 1/2 or 3 lbs. per 1,000 sq.ft. Apply 1/2 to 1 inches of water within 24 hours after treatment. Reapply as necessary for pest reinfestation. Do not apply more than 6 lbs. of OFTANOL 1.5% Granular per 1,000 sq.ft. per year or an equivalent amount of 1.45 oz. active ingredient per 1,000 sq.ft. regardless of formulation used. Minimum interval between applications is 15 days for 1.5 lb. rate and 30 days for the 3 lbs. rate.

Consult your local turf, state Agricultural Experiment Station, or Extension Service Specialist for more specific information regarding timing of application.

RESTRICTIONS

Do not mow treated turf area before rainfall or irrigation has been applied as per RECOMMENDED APPLICATIONS. Do not graze treated areas or use clippings from treated areas for food or forage. Keep children and pets off treated area until dry.



STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage or disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.


WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

MANUFACTURED BY:

the professional's partner
THE ANDERSONS
LAWN FERTILIZER DIVISION
P.O. BOX 119
MAUMEE, OH 43537



:9198-88R
05/01/90