

(FRONT PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

THE ANDERSONS TEE TIME with 1.5% OFTANOL

ACTIVE INGREDIENTS:

1-Methylethyl 2-((ethoxy((1-methylethyl)amino)phosphorothioyl)oxy)benzoate 1.5%

INERT INGREDIENTS 98.5%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT __ LBS.

TREATS __ SQ.FT.

ACCEPTED
JUN 24 1998
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9198-88

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

NET WEIGHT _ LBS.

TREATS ___ SQ.FT.

THE ANDERSONS TEE TIME WITH 1.5% OFTANOL

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush with plenty of water.

IF ON SKIN: Wash with soap and hot water.

IF INHALED: Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes

This product is recommended for use on lawns and other ornamental and recreational turf grass areas.

RECOMMENDED APPLICATIONS

ON LAWNS:

Apply this product at the rate of 3 lbs. per 1,000 sq.ft. for the following insects: Whitegrub larvae (such as Japanese Beetle, Black Turfgrass Ataenius, Chafers, Phytophaga spp., Green June Beetle, Oriental Beetle, Asiatic Garden Beetle) and Mole Crickets, Larval stage of Billbug and Hyperodes Weevil.

White Grub and Billbug Larvae Control: For best results Water turf area immediately after application, using sufficient water to wet soil 1 to 1 1/2 inches deep.

Hyperodes Weevil Control: Apply between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15 and water in as for preventative White Grub control.

Mole Cricket Control: For best results Water immediately after application using 1/2 to 1 inch water.

Consult your local turf, state Agricultural Experiment Station, or Extension Service Specialists for more specific information regarding timing of application.

FOR FIRE ANTS:

Area Fire Ant Control: Apply 3 lbs./1,000 sq.ft. uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present, use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

Established Fire Ant Mounds - Drench Treatments: Wet Conditions: Distribute 0.5 oz. evenly over 9 sq.ft. and immediately sprinkle gently with 3 gallons of water. Dry Conditions: Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one Mound Drench Treatment.

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

FOR SURFACE INSECTS:

For Chinch Bugs and Sod Webworm Larvae: Make application at the rate of 1 1/2 or 3 lbs. per 1,000 sq. ft. **For best results** Apply 1/4 to 1 inches of water immediately after application. Reapply as necessary for pest reinfestation. Minimum interval between applications is 15 days for 1.5 lb. rate and 30 days for the 3 lbs. rate.

Consult your local turf, state Agricultural Experiment Station, or Extension Service Specialists for more specific information regarding timing of application.

SUGGESTED SPREADER SETTINGS

(Available at final printing)

NOTE: These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate. Do not hand spread.

RESTRICTIONS

This product shall be watered in by the applicator with sufficient water as listed on the directions.

Do not mow treated turf area before rainfall or irrigation has been applied. **Do not collect clippings at first mowing after application.** Do not graze treated areas or use clippings from treated areas for food or forage. Keep children and pets off treated area until dry.

Do not apply more than 6 lbs. of this product per 1,000 sq.ft. per year or an equivalent amount of 1.45 oz. active ingredient per 1,000 sq.ft. regardless of formulation used. Minimum interval between applications must be 30 days for the 3 lbs. rate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticides are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

MANUFACTURERS STATEMENT

This product conforms to the ingredient statement on the label. Manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage and handling of this product.

THIS PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED. IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND.

**Manufactured by:
The Andersons Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537**

Date Printed: 06/02/98
Date Approved: 02/22/95
Supersedes: 08/31/92
Label Change: correct wps exclusionary/delete 'for best results' in watering statement/add restrictions/add spreader directions/update mfg statement
Filename: 9198-88C

6/10

(FRONT PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

THE ANDERSONS TEE TIME with 1.5% OFTANOL

ACTIVE INGREDIENTS:

1-Methylethyl 2-((ethoxy((1-methylethyl)amino)phosphorothioyl)oxy)benzoate 1.5%

INERT INGREDIENTS 98.5%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT _ LBS.

TREATS _ SQ.FT.

ACCEPTED
JUN 24 1998
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9198-88

(BACK PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

NET WEIGHT _ LBS.

TREATS ___ SQ.FT.

THE ANDERSONS TEE TIME WITH 1.5% OFTANOL

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush with plenty of water.

IF ON SKIN: Wash with soap and hot water.

IF INHALED: Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, and wildlife. Do not apply to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

This product is recommended for use on home lawns and other ornamental and recreational turf grass areas around the home.

(BACK PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

RECOMMENDED APPLICATIONS ON LAWNS

Apply this product at the rate of 3 lbs. per 1,000 sq.ft. for the following insects: Whitegrub larvae (such as Japanese Beetle, Black Turfgrass Ataenius, Chafer, Phytophaga spp., Green June Beetle, Oriental Beetle, Asiatic Garden Beetle) and Mole Crickets, Larval stage of Billbug and Hyperodes Weevil.

White Grub and Billbug Larvae Control: For best results Water turf area immediately after application. Using sufficient water to wet soil 1 to 1 1/2 inches deep.

Hyperodes Weevil Control: Apply between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15 and water in as for preventative White Grub control.

Mole Cricket Control: For best results Water immediately after application using 1/2 to 1 inch water.

Consult your local turf, state Agricultural Experiment Station, or Extension Service Specialists for more specific information regarding timing of application.

FOR FIRE ANTS:

Area Fire Ant Control: Apply 3 lbs./1,000 sq.ft. uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present, use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

Established Fire Ant Mounds - Drench Treatments: Wet Conditions: Distribute 0.5 oz. evenly over 9 sq.ft. and immediately sprinkle gently with 3 gallons of water. Dry Conditions: Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one Mound Drench Treatment.

SURFACE INSECTS:

For Chinch Bugs and Sod Webworm Larvae: Make application at the rate of 1 1/2 or 3 lbs. per 1,000 sq.ft. For best results Apply 1/2 to 1 inches of water immediately after application. Reapply as necessary for pest reinfestation. Minimum interval between applications is 15 days for 1.5 lb. rate and 30 days for the 3 lbs. rate.

Consult your local turf, state Agricultural Experiment Station, or Extension Service Specialists for more specific information regarding timing of application.

9/10

(BACK PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

SUGGESTED SPREADER SETTINGS

(Available at final printing)

NOTE: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of terrain. Calculate the size of the area to be treated and do not over or under apply. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary, in order to assure correct rate of application. Close spreader when filling, stopping or turning. Do not hand spread.

RESTRICTIONS

Do not mow treated turf area before ~~rainfall~~ or irrigation has been applied as per RECOMMENDED APPLICATIONS. Do not graze treated areas or use clippings from treated areas for food or forage. Keep children and pets off treated area until dry.

Do not apply more than 6 lbs. of this product per 1,000 sq.ft. per year or an equivalent amount of 1.45 oz. active ingredient per 1,000 sq.ft. regardless of formulation used. Minimum interval between applications must be 30 days for the 3 lbs. rate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

(BACK PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

MANUFACTURERS STATEMENT

This product conforms to the ingredient statement on the label. Manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage and handling of this product.

THIS PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED. IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND.

Manufactured by: The Andersons Lawn Fertilizer Division
d/b/a Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537

Date Printed: 06/01/98
Date Approved: 02/22/95
Supersedes: 08/31/92
Label Change: delete 'for best results' on watering statement/update mfg statement
Filename: 9198-88H