

1016
Pm05-
4198-90

Tee-Time 5-10-30 with Balan and Surflan, Formula I, Net Weight 50 pounds

Tee-Time 5-10-30 with Balan and Surflan, Formula I, is a selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

Bahiagrass	Centipedegrass	Zoysiagrass
Bermudagrass	St. Augustinegrass	Tall Fescue

Active Ingredients:


N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine57%
oryzalin: 3,5-dinitro-N4,N4-dipropylsulfanilamide57%
Inert ingredients	98.86%

Contains .57 pounds active ingredients per 50 pound bag

Balan(R) and Surflan(R) are trademarks of Elanco Products.

**Keep Out of Reach of Children
CAUTION**

EPA Reg. No. 9198-



Directions for Use Read All Directions Carefully Before Applying

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Formula I is a granular preemergence herbicide which is applied to established Southern warm season turf (excluding greens) for the control of annual grasses and some broadleaf weeds. Formula I works by controlling weeds as the seeds germinate.

ANNUAL GRASSES CONTROLLED BY FORMULA I


- Summer Annuals**
- Barnyardgrass (watergrass)
- Crabgrass
 - (Large Crabgrass)
 - (Smooth Crabgrass)
- Crowfootgrass
- Foxtails
 - (Bottlegrass)
 - (Bristlegrass)
 - (Giant Foxtail)
 - (Green Foxtail)
 - (Pigeongrass)
 - (Robust Foxtail)
 - (Yellow Foxtail)
- Goosegrass
 - (Silver Crabgrass)
- Johnsongrass
 - (form seed only)
- Ryegrass, annual
- Sandbur, field

- Winter Annuals**
- Bluegrass, annual

ANNUAL BROADLEAF WEEDS CONTROLLED BY FORMULA I

- Summer Annuals**
- Carpetweed
- Knotweed, prostrate
- Purslane, common

- Winter Annuals**
- Chickweed, common
- Henbit



ANNUAL BROADLEAF WEEDS PARTIALLY CONTROLLED BY FORMULA I

- Groundsel, common
- Spurge, prostrate
- Woodsorrel, yellow

GENERAL DIRECTIONS

RATE, FREQUENCY AND TIMING OF APPLICATION

Successful preemergence control of the annual weeds listed on this label requires proper timing of application.

Apply Formula I prior to the germination of annual weeds. All summer annual weeds do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual weeds listed on this label, weed control results may be erratic or poor. In the absence of timely rainfall, irrigation can be used to move Formula I into the soil surface.

All cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Formula I.

APPLICATION DIRECTIONS

Formula I can be applied in the Spring for summer annual grass and broadleaf weed control, and in the fall for Poa annua and winter annual broadleaf weed control.

1. **SPRING GERMINATING SUMMER ANNUAL GRASSES AND BROADLEAF WEEDS.**
For preemergence control, apply 4-6 pounds per 1000 square feet (2-3 pounds active ingredient per acre) in the late winter or early spring prior to the onset of conditions favorable for annual weed grass germination. For control of goosegrass, apply 6 pounds per 1000 square feet (3 pounds active ingredient per acre) in the late winter or early spring.
2. **POA ANNUA (ANNUAL BLUEGRASS) AND BROADLEAF WINTER ANNUALS.**
For preemergence control, apply 6 pounds per 1000 square feet (3 pounds active ingredient per acre) in the late summer or early fall prior to the expected germination of annual bluegrass. Do not apply Formula I on areas where fall overseeding will occur. In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

For application to smaller areas, apply the following amounts of Formula I.

Formula I per acre
 175 pounds
 263 pounds

Formula I per 1000 sq. ft.
 4 pounds
 6 pounds

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying Formula I. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding

results after using Formula I the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using Formula I at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

APPLICATION EQUIPMENT

Apply Formula I evenly over the turfgrass area without skips or misses. A more uniform application can be made by spreading half the required amount over the area and then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated band. For best results use equipment designed to apply granular herbicides and insecticides. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

APPROPRIATE SPREADER BETTING AS A GUIDE FOR CALIBRATION

Note: These settings are provided as suggested starting points in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations.

Spreader Type	Spreader Setting Necessary To Apply Pounds Formulated Product Per 1000 sq. ft.	
	4 lbs./1000 sq. ft.	6 lbs./1000 sq. ft.
Greenview Drop	9 1/2	10 1/2
Central Broadcast	4 1/2	4 3/4
Cyclone Broadcast	4	5
Lely	4	5
Ortho Drop	6	6 1/2
Ortho Broadcast	1	1 1/2
Scotts Drop	3 1/2	4 1/2
Scotts Broadcast	L 1/2	M 1/2
Scotts Professional Broadcast	H 1/2	I 1/2
Sears Drop	8 1/2	9 1/2
Sears Broadcast	4 1/2	4 3/4
Central Drop	9 1/2	10 1/2

PRECAUTIONS

Apply Formula I to the turf prior to the germination of weed grasses since Formula I will not control established weeds. Formula I may thin established annual bluegrass (*Poa annua*) turfs and the fine-leaved fescues at rates above 175 pounds per acre (2 pounds active ingredient). Formula I should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been over-seeded with winter grasses, a spring application of Formula I may thin the overseeded grasses. Formula I should not be used on dichondra. Formula I should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until these turfs are well established.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or eyes. Use eye protection and protective clothing such as overalls, a long-sleeved shirt, and impermeable gloves when handling this product.

Statement of Practical Treatment

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation exists.

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If inhaled: If discomfort occurs, remove individual to fresh air. Get medical attention if breathing difficulty occurs.

Environmental Hazards: Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL


Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE: The directions on this label were determined through research to be the directions for correct use of this product. This product has been tested for a range of weather conditions similar to those weather conditions that are ordinary and customary in the geographic area where the product is used. Insufficient control of pests may result from the occurrence of extraordinary or unusual weather, or from failure to follow label directions. In addition, failure to follow label directions may cause injury to the crop to which the product is applied, other crops, animals, man or the environment. The Andersons and the seller offer, and the buyer accepts and uses this product subject to the conditions that extraordinary or unusual weather, or failure to follow label directions are beyond the control of The Andersons and are, therefore, the responsibility of the buyer.

The Andersons
Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537



(6)