

pm25

9198-91

1714

9198-91C
Page 1 of 7


(FRONT PANEL)

COMMERCIAL USE ONLY

THE ANDERSONS

TEE TIME

PLUS 0.57% TEAM



A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass.

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-alpha,alpha,alpha-trifluoro-2,6-dinitro-p-toluidine 0.38%

Trifluralin: alpha,alpha,alpha-trifluoro-2,6-dinitro-N,
N-dipropyl-p-toluidine 0.19%

INERT INGREDIENTS 99.43%

KEEP OUT OF THE REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 50 LBS. (22.7 kg)

TREATS UP TO 12,500 SQ.FT.

EPA Reg. No. 9198-91

EPA Est. 9198-QH-1

(BACK PANEL)

COMMERCIAL USE ONLY

NET WEIGHT 50 LBS. (22.7 kg)

Treats up to 12,500 Sq.Ft.

THE ANDERSONS

TEE TIME

PLUS 0.57% TEAM

GUARANTEED ANALYSIS

| | |
|--|------|
| Total Nitrogen (N) | 25% |
| 1.2% Ammoniacal Nitrogen | |
| 17.3% Urea Nitrogen | |
| 6.5% Coated Slow Release Urea Nitrogen | |
| Available Phosphoric Acid (P2O5) | 3% |
| Soluble Potash (K2O) | 8% |
| Sulfur (S) | 4.8% |

Primary Plant Foods Derived from: Ammoniated Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

WARNING

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

BEST AVAILABLE

(BACK PANEL)

COMMERCIAL USE ONLY

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Call a physician.
- IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.
- IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.
- IF INHALED: If breathing difficulty occurs, move individual to fresh air and get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product has been specially formulated to provide a second kill during the growing season. This additional one pound of active ingredient per acre should be used as a second application following another formulation providing two pounds of active ingredient per acre. This product will control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass, silver crabgrass or crowfoot, barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. When this product is applied as directed, the following established turfgrasses are tolerant:

EST. 10/1/80

(BACK PANEL)

COMMERCIAL USE ONLY

| | | | |
|--------------------|----------------|--------------|--------------------|
| Kentucky bluegrass | Centipedegrass | Zoysiagrass | St. Augustinegrass |
| Perennial ryegrass | Fescue | Bermudagrass | Bahiagrass |

This product will not harm most nearby established ornamental trees and shrubs.

APPLICATION DIRECTIONS

RATES, FREQUENCY OF TIMING OF APPLICATION

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass
 - A. Northern (cool season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product (1 pound active ingredient per acre) eight to ten weeks after early spring application of this product (2 pounds active ingredient per acre).
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product (1 pound active ingredient per acre) eight to ten weeks after the late winter or early spring application of this product (3 pounds active ingredient per acre).
2. Late spring, early summer germinating annual grasses - goosegrass
Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications of this product should, therefore, be timed accordingly.

BEST AVAILABLE G

(BACK PANEL)

COMMERCIAL USE ONLY

- A. Northern (cool season) turfgrass
For preemergence control, apply 4 pounds per thousand square feet of this product (1 pound active ingredient per acre) eight to ten weeks after the late spring or early summer application of this product (2 pounds active ingredient per acre).
- B. Southern (warm season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product (1 pound active ingredient per acre) eight to ten weeks after the late spring or early summer application of this product (3 pounds active ingredient per acre).
3. Northern and southern turfgrasses
In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications of this product spaced eight to ten weeks apart may be made to provide full season control of these annual grass species.
4. Poa annua (annual bluegrass)
In areas of heavy infestation or annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.
- A. Northern (cool season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product (1 pound active ingredient per acre) eight to ten weeks after the late summer or early fall applications of this product (2 pounds active ingredient per acre).
- B. Southern (warm season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product (1 pound active ingredient per acre) eight to ten weeks after the late summer or early fall applications of this product (3 pounds active ingredient per acre).

BEST AVAILABLE

(BACK PANEL)

COMMERCIAL USE ONLY

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization can be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

USE PRECAUTIONS

Apply this product to the turf prior to the germination of weed grasses since it will not control established weeds. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of this product should not be used on putting greens. This product should not be used on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

SUGGESTED SPREADER SETTINGS

(Available at final printing)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE

(BACK PANEL)

COMMERCIAL USE ONLY

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of applications, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by: The Andersons
Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537

Date Printed: 01/24/94
Date Approved:
Supersedes:
Label Change:


8/14

(FRONT PANEL)
HOMEOWNER USE ONLY

THE ANDERSONS

TEE TIME

PLUS 0.57% TEAM



A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass.

| | | |
|--|--|---------------|
| ACTIVE INGREDIENTS: | | |
| N-butyln-ethyl-alpha, alpha, alpha-trifluoro-2,6-dinitro-p-toluidine | | 0.38% |
| Trifluralin: alpha, alpha, alpha-trifluoro-2,6-dinitro-N, N-dipropyl-p-toluidine | | 0.19% |
| INERT INGREDIENTS | | 99.43% |

KEEP OUT OF THE REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 50 LBS. (22.7 kg)

TREATS UP TO 12,500 SQ.FT.

EPA Reg. No. 9198-91

EPA Reg. No. 9198-OH-1

BEST AVAILABLE

9814

(BACK PANEL)

HOMEOWNER USE ONLY

NET WEIGHT 50 LBS. (22.7 kg)

Treats up to 12,500 Sq.Ft.

THE ANDERSONS

TEE TIME

PLUS 0.57% TEAM

GUARANTEED ANALYSIS

| | |
|--|------|
| Total Nitrogen (N) | 25% |
| 1.2% Ammoniacal Nitrogen | |
| 17.3% Urea Nitrogen | |
| 6.5% Coated Slow Release Urea Nitrogen | |
| Available Phosphoric Acid (P2O5) | 3% |
| Soluble Potash (K2O) | 8% |
| Sulfur (S) | 4.8% |

Primary Plant Foods Derived from: Ammoniated Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

WARNING

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

BEST AVAILABLE

(BACK PANEL)

HOMEOWNER USE ONLY

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Call a physician.
- IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.
- IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.
- IF INHALED: If breathing difficulty occurs, move individual to fresh air and get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply to water. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product has been specially formulated to provide a second kill during the growing season. This additional one pound of active ingredient per acre should be used as a second application following another formulation providing two pounds of active ingredient per acre. This product will control annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass, silver crabgrass or crowfoot, barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. When this product is applied as directed, the following established turfgrasses are tolerant:

BEST

(BACK PANEL)

HOMEOWNER USE ONLY

| | | | |
|--------------------|----------------|--------------|--------------------|
| Kentucky bluegrass | Centipedegrass | Zoysiagrass | St. Augustinegrass |
| Perennial ryegrass | Fescue | Bermudagrass | Bahiagrass |

This product will not harm most nearby established ornamental trees and shrubs.

APPLICATION DIRECTIONS

RATES, FREQUENCY OF TIMING OF APPLICATION

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Tee Time Plus Team.

1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass
 - A. Northern (cool season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product eight to ten weeks after early spring application of this product.
 - B. Southern (warm season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product eight to ten weeks after the late winter or early spring application of this product.
2. Late spring, early summer germinating annual grasses - goosegrass
Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications of this product should, therefore, be timed accordingly.
 - A. Northern (cool season) turfgrass
For preemergence control, apply 4 pounds per thousand square feet of this product eight to ten weeks after the late spring or early summer application of this product.

BEST AVAILABLE COPY

(BACK PANEL)

HOMEOWNER USE ONLY

- B. Southern (warm season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product eight to ten weeks after the late spring or early summer application of this product.
3. Northern and southern turfgrasses
In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications of this product spaced eight to ten weeks apart may be made to provide full season control of these annual grass species.
4. Poa annua (annual bluegrass)
In areas of heavy infestation or annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.
- A. Northern (cool season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product eight to ten weeks after the late summer or early fall applications of this product.
- B. Southern (warm season) turfgrasses
For preemergence control, apply 4 pounds per thousand square feet of this product eight to ten weeks after the late summer or early fall applications of this product.

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization can be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 15 weeks after application.

BEST AVAILABLE COPY

13 of 14

9198-91H
Page 6 of 7

(BACK PANEL)

HOMEOWNER USE ONLY

USE PRECAUTIONS

Apply this product to the turf prior to the germination of weed grasses since this product will not control established weeds. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of this product should not be used on putting greens. This product should not be used on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

SUGGESTED SPREADER SETTINGS

(Available at final printing)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

14 of 14

(BACK PANEL)

HOMEOWNER USE ONLY

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of applications, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by: The Andersons
Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537

Date Printed: 01/25/94
Date Approved:
Supersedes:
Label Change: