

10F6
Pm 25
9198-94

Front Label

Andersons Tee-Time 20-4-10 Plus Team(R)

(LOGO)

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established: Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysiagrass.

Active Ingredients:

N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine62%
trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine30%
Inert Ingredients	99.08%

TEAM(R) ---- the trademark for Elanco Products benefin plus trifluralin

Keep Out of Reach of Children

Warning

See Back Panel For Additional Precautionary Statements

Net Weight 50 lbs. (22.7 kg)

Treats Up To 10,000 Sq.Ft.


EPA REG. NO. 9198-

EPA Est. 9198-OH-1

ACCEPTED

AUG - 4 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9198-94



Back Panel

Net Weight 50 lbs. (22.7 kg)

Treats up to 10,000 Sq.Ft.

Andersons Tee-Time 20-4-10 Plus Team(R)

Guaranteed Analysis

Total Nitrogen (N)	20%
1.6% Ammoniacal Nitrogen	
11.9% Urea Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O)	10%
Sulfur	4.8%

Plant Nutrients Derived From: Ammoniated Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea. Potential Acidity: 54 lbs. Calcium Carbonate Equivalent per ton.

**Precautionary Statements
Keep Out of Reach of Children
Hazards To Humans And Domestic Animals
Warning**

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Environmental Hazards

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

Statement of Practical Treatments:

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.

If swallowed: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Tee-Time Plus Team(R) has a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (Poa annua), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass), and green and yellow foxtail. Tee-Time Plus Team(R) controls these annual weeds by killing their seeds as they germinate. Apply Tee Time Plus Team(R) before these annual weeds begin to germinate since it does not control established weeds.

When Tee-Time Plus Team(R) is applied as directed, the following established turfgrasses are tolerant:

Kentucky bluegrass	Fescue	St. Augustinegrass
Perennial ryegrass	Zoysiagrass	Bahiagrass
Centipedegrass	Bermudagrass	

Tee-Time Plus Team(R) will not harm most nearby established ornamental trees and shrubs.

Application Directions

Rates, Frequency and Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply Tee Time Plus Team(R) approximately one or two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application of Tee-Time Plus Team(R).

1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass.
 - A. Northern (cool season) turfgrasses
For preemergence control, apply 5 pounds per thousand square feet of the Tee-Time Plus Team(R) (2 pounds active ingredient per acre) in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

- B. Southern (warm season) turfgrasses
For preemergence control, apply 7.5 pounds per thousand square feet of the Tee-Time Plus Team(R) (3 pounds active ingredient per acre) in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of Tee Time Plus Team(R) from one half to the full recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

- 2. Late spring, early summer germinating annual grasses - goosegrass. Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications of Tee-Time Plus Team(R) should, therefore, be timed accordingly.

- A. Northern (cool season) turfgrass
For preemergence control, apply 5 pounds per thousand square feet of Tee-Time Plus Team(R) (2 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
- B. Southern (warm season) turfgrass
For preemergence control, apply 7.5 pounds per thousand square feet of Tee-Time Plus Team(R) (3 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

3. Northern and Southern turfgrasses

In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications of from one half to the full recommended rate 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

4. Poa annua (annual bluegrass)

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

- A. Northern (cool season) turfgrasses
For preemergence control, apply 5 pounds per thousand square feet of Tee-Time Plus Team(R) (2 pounds active ingredient per acre) in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.

B. Southern (warm season) turfgrasses
 For preemergence control, apply 7.5 pounds per thousand square feet of Tee-Time Plus Team(R) (3 pounds active ingredient per acre) in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application of Tee Time Plus Team(R) from 3.75 to 7.5 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.

Reseeding

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying Tee-Time Plus Team(R). When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using Tee-Time Plus Team(R) the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as Rogers Aero Seeder) should be employed. When using Tee-Time Plus Team(R) at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

Application Equipment

Apply Tee-Time Plus Team(R) evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area and then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly.

Suggested Spreader Settings

(This section will be completed for final labeling)

Use Precautions

Apply Tee-Time Plus Team(R) to the turf prior to the germination of weed grasses since Tee-Time Plus Team(R) will not control established weeds. Tee-Time Plus Team(R) should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of Tee-Time Plus Team(R) should not be used on putting greens. Tee-Time Plus Team(R) should not be used on dichondra. Tee-Time Plus Team(R) should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

Storage and Disposal

Storage: Storage in original container only. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Do not contaminate water, food, or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

NOTICE

"Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided with proper equipment in good working order and under normal conditions of use. Buyer assumes the risk on any contrary use. Seller makes no other express or implied warranty including the warranty of fitness for a particular purpose, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made."

Manufactured By: The Andersons
Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537

