

Tee Time Fertilizer 20-4-10 with Team

The bag is printed with the product name and with either brown, red or green field on the top and bottom. Logo is reversed out of the color field or printed in black over the color.

9198 94

the professional's partner




A selective preemergence herbicide and fertilizer product for the control of crabgrass and other annual weed grasses on lawns and golf courses, new established Bluegrass, Kentucky bluegrass, perennial ryegrass, fescue, Bahiagrass, Bermudagrass, Centipedegrass, and Augustgrass and crabgrass.

Active Ingredients:

2-(Chloroethyl)benzyl-alphahydroxyethylcarbamate should be 1.62%
Tetraethylammonium chloride, 1.4%, and Methylene bisacrylate
as emulsifying agents.

Trade Name: The Andersons Tee Time Fertilizer



12 pt.
14 pt.

Net Weight 50 Lbs. (22.7 kg)
Treats up to 10,000 Sq. Ft.

EPA Reg. No. 9198-94

EPA Est. 9198-OH-1

100 J. K. LAMBERT

**Net Weight 50 Lbs. (22.7 kg.)
Treats up to 10,000 Sq. Ft.**

Tee Time Fertilizer **20-4-19** with Team

GUARANTEED ANALYSIS

1. The following are:
 (a) All members of the family.
 (b) All persons in the household.
 (c) All persons in the same dwelling.
 (d) All persons in the same room.
 (e) All persons in the same house.
 (f) All persons in the same building.
 (g) All persons in the same block.
 (h) All persons in the same city.
 (i) All persons in the same state.
 (j) All persons in the same country.

PRECAUTIONARY STATEMENTS
KEEP OUT OF REACH OF CHILDREN
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

STATEMENT OF PRACTICAL TREATMENT

STATEMENT OF PRACTICAL TREATMENT
If eyes - Wash eyes clean and wash with a weakly saline cream of tartar for 15 minutes. Call a physician.
If swallowed - An adult who has ingested a large quantity of milk, egg white, gelatin solution, or these are not available, large quantities of water. Avoid alcohol.
If on skin - Wash with gentle soap and water. If medical attention persists.

MANAGEMENT AND DESIGN

This pesticide is flammable and may be dangerous to water or wetland life if spilled, leaked, or mishandled. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring waters.

DIRECTIONS FOR USE

DIRECTIONS FOR USE

142-10-1000

The following eight species of grasses are present:

1. 10/15/2015 - 09:30 a.m. - 09:45 a.m. - Open the door to a room where a man is sitting at a desk.

APPLICATION DIRECTIONS

Rates, Frequency and Timing of Appearance

5.1. General framework for the analysis of the CBR (5) class metric, based on the above two main sources of uncertainty.

For best results, mix the granules with the recommended water volume just before use. At 100ml/ha, 10g of granules should germinate at the same time. If greater than 10% seed has not germinated within 7 days, reapply.

2. In areas where the soil is cold and relatively moist, the time of the year needs to be extended to fall, winter, and spring.

3. Sowing germinating summer annual grasses - ryegrass, fescues and hairy-meadowgrass.

A. Ryegrass - 10 kg/ha per hectare

B. Fescue (either - 10 kg/ha per hectare) or hairy-meadowgrass (the latter is more expensive) in the spring (one to two weeks prior to the onset of frost) one kg/ha per hectare.

C. Hairy-meadowgrass

D. Winter-warm-season ryegrass

The winter-warm-season ryegrass 1.5 kg/ha per hectare per square meter. The sowing agreement per hectare in the late winter or early spring (one to two weeks prior to the onset of cold temperatures) 1.5 kg/ha per hectare.

E. Winter-warm grass germination

In areas of the United States where germination of the annual seedling has stopped, continues for extended periods of time, a second application (one half to the full recommended rate) may be made 8 to 10 weeks after initial sowing for maximum control.

4. Sowing early-summer germinating annual grasses - ryegrass.

A. Ryegrass and summer annual grasses - ryegrass, some ryegrass varieties in 8 weeks after sowing, fescues and other summer annual grass species. Applications should therefore be

B. Sowing

(See back page 1)

A. Fall Seeding - Annual Grasses

For emergence control, apply 5 pounds per thousand square feet at the rate of 15 pounds per thousand square feet applied two weeks prior to the expected germination of grasses.

B. Fall Seeding - Warm Season Grasses

For emergence control, apply 15 pounds per thousand square feet applied two weeks prior to the expected germination of grasses.

C. Northern and Southern Turfgrasses

Grasses other than those areas where preemergence control is desired, apply 5 pounds per thousand square feet applied two weeks prior to the expected germination of grasses.

D. Pre-emergence Annual Grasses

For annual grasses, apply 5 pounds per thousand square feet applied two weeks prior to the expected germination of grasses and two weeks prior to the application of seed.

E. Fall Seeding - Warm Season Turfgrasses

For warm season turfgrasses, apply 5 pounds per thousand square feet applied two weeks prior to the expected germination of grasses.

F. Fall Seeding - Warm Season Turfgrasses

For warm season turfgrasses, apply 15 pounds per thousand square feet applied two weeks prior to the expected germination of grasses.

RESEEDING

Reseeding is not recommended for warm season annual weed seeds. If reseeding is desired, it should be delayed until at least 6 weeks after applying. When reseeding, if seed is to be applied by hand, use a hand spreader to broadcast seed evenly over the area and then apply a light mist of water to keep the seed from blowing away. If seed is to be applied by machine, spread seed evenly over the area and then apply a light mist of water to keep the seed from blowing away.

APPLICATION EQUIPMENT

All equipment used for spreading seed without seed spreaders. A hand spreader is recommended for warm season annual weed seeds. Broadcast seed evenly over the area and then apply a light mist of water to keep the seed from blowing away.

SUGGESTED SPREADER SETTINGS

Spreader Type	Speed	Spread Width	Settings	
			4 lbs./1000	8 lbs./1000
Hand Spreader	10 ft/min	10' 12"	5	7
USDA TESTED Broadcast Spreader	10 ft/min	10' 12"	0.6	0.3
USDA TESTED Broadcast Spreader	10 ft/min	24' 9"	5	7

USE PRECAUTIONS

Always use caution when applying weed control products to lawns or ornamental areas. This product should be applied in the spring. Do not apply to lawns planted to chrysanthemums, dianthus, or any other plants that have been sprayed with a herbicide containing dicamba. Do not apply on orchids. This product should not be applied to new lawns or areas of bare ground. Avoid application to sensitive grasses. All statements made concerning this product are based on facts known to us.

STORAGE AND DISPOSAL

Storage: Store in original containers only. In case of eye contact, wash eyes immediately with water.

Disposal: Disperse or dilute waste liquid to meet disposal requirements of local, state, and federal regulations. Do not pour unused waste down storm drains or into containers for disposal. Do not spray onto any surface which may be washed into a storm drain. Wash containers and rinsing equipment with water and return to the manufacturer or distributor. Do not reuse containers. Do not contaminate water supplies.

The manufacturer makes no warranties regarding the use of this product. The user is responsible for determining the use of this product in accordance with all statements made concerning this product. All statements made concerning this product are based on facts known to us.

NOTICE

The manufacturer makes no warranties regarding the use of this product. The user is responsible for determining the use of this product in accordance with all statements made concerning this product. All statements made concerning this product are based on facts known to us.

Manufactured by
**the professional's
 partner®**



Lawn Fertilizer Division
 P.O. Box 119
 Maumee, Ohio 43537

© Copyright 1987 T.A.
 0100-04

