

Tee Time[®] Fertilizer 20-4-10 with Team[®]

JUN 1 1988
9198-94

the professional's
partner



A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual, weed grasses on lawns and golf courses in established Bluegrass (perennial), Ryegrass (perennial), Fescue (Bahia grass, bermudagrass, Centipedegrass, St. Augustine grass and Zoysiagrass).

Active Ingredients
 N-buty-N-ethyl-alpha, alpha-chloro-N-(2-Ethoxyethyl)amine 0.62%
 Thiuram, alpha, alpha-chloro-N-(2-Ethoxyethyl)amine 0.30%
 Inert Ingredients 99.08%

TEAM — the trademark for Fisons' products bearing this trademark.

KEEP OUT OF THE REACH OF CHILDREN
WARNING
 See Back Panel for Additional Precautions

Net Weight 50 Lbs. (22.7 kg)
 Treats up to 10,000 Sq. Ft.

- B. Southern warm season turfgrasses
For preemergence control, apply 7.5 pounds per thousand square feet (3 pounds active ingredient per acre) in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass germination.
In areas of the United States where germination of the annual weed grasses is extended for extended periods of time, a second application from one half to the full recommended rate may be made 8 to 10 weeks after initial treatment for continued control.
2. Late spring-early summer germinating annual grasses—goosegrass
A. Insects classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks after crabgrass, foxtails and other summer annual grass species. Applications should, therefore, be timed accordingly.
A. Northern cool season turfgrass
For preemergence control, apply 3.75 to 5 pounds per thousand square feet (1.5 to 2 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
B. Southern warm season turfgrass
For preemergence control, apply 7.5 pounds per thousand square feet (3 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
3. Northern and Southern turfgrasses
In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications of from one half to the full recommended rate 8 to 10 weeks apart may be made to provide full season control of these annual grass species.
4. Post annual (annual bluegrass)
In areas of heavy infestation of annual bluegrass, its elimination will typically result in thin turfgrass areas. Irrigation and soil incorporated reseedling will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.
A. Northern cool season turfgrasses
For preemergence control, apply 3.75 to 5 pounds per thousand square feet (1.5 to 2 pounds active ingredient per acre) in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.
B. Southern warm season turfgrasses
For preemergence control, apply 7.5 pounds per thousand square feet (3 pounds active ingredient per acre) in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application from 3.75 to 7.5 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.

RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as Rogers Aero Seeder) should be employed. When using at the highest recommended rate, reseeding should be delayed 10 to 16 weeks after application.

APPLICATION EQUIPMENT

Apply evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area and then applying the remaining half at right angles. Both chip type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly.

SUGGESTED SPREADER SETTINGS

Spreader Type	Speed	Spread Width	Settings	
			4 lbs./1000	6 lbs./1000
Cyclone	3 MPH	10-12 ft.	E	7
Scotts Broadcast Broadcast	3 MPH	10-12 ft.	D-E	F-G
Leif Type M	5 MPH	29 ft.	E	7

USE PRECAUTIONS

Apply to the turf prior to the germination of weed grasses, since Tee-Time with Teq™ will not control established weeds. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application should not be used on putting green. Do not apply on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

STORAGE AND DISPOSAL

- Storage:** Store in original container only. In case of spill, contain material and dispose as waste.
- Pesticide Disposal:** Do not contaminate water, food or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

The manufacturer makes no warranties, express or implied, concerning this product or its use which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

NOTICE

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided with proper equipment in good working order under normal conditions of use. Buyer assumes the risk on any contrary use. Seller makes no other express or implied warranty, including the warranty of fitness for a particular purpose, and no agent of seller is authorized to do so, except in writing with a specific reference to this warranty. In no event shall seller's liability for any breach of warranty exceed the purchase price of the material, as to which a claim is made.

Manufactured by
**the professional's
partner®**



Lawn Fertilizer Division
P.O. Box 119
Maumee, Ohio 43537

© Copyright 1987 T.A.
9-85-84