

Front Panel  
Commercial Use only  
(for internal use only-will not appear on label)

# THE ANDERSONS TEE TIME WITH 0.92% TEAM\*

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established: Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysiagrass.

### ACTIVE INGREDIENTS:

Benfenin: N-butyl-N-ethyl-alpha,alpha,alpha,-trifluoro-2,6-dinitro-p-toluidine .....	0.62%
Trifluralin: alpha,alpha,alpha-trifluoro-2,6-dinitro-N, N-dipropyl-p-toluidine .....	0.30%

**INERT INGREDIENTS** ..... 99.08%

Total .. 100.00%


\*Trademark of Dow AgroSciences LLC

**KEEP OUT OF REACH OF CHILDREN  
WARNING  
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS**

NET WEIGHT 50 LBS. (22.7 kg)

TREATS UP TO 10,000 SQ.FT.

EPA Reg. No. 9198-94  
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A  
Underlined letter is first letter used in run code on bag



(BACK PANEL)  
COMMERCIAL USE ONLY  
*(for internal use only-will not appear on final label)*

NET WEIGHT 50 LBS. (22.7 kg)  
TREATS UP TO 10,000 SQ.FT.

## THE ANDERSONS TEE TIME WITH 0.92% TEAM

<b>Guaranteed Analysis</b>	
Total Nitrogen (N) .....	XX%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%
Plant nutrients derived from: <i>For products being sold into Washington state:</i>	
Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.regulatory-info-ap.com/">http://www.regulatory-info-ap.com/</a>	

### PRECAUTIONARY STATEMENTS

FIRST AID	
If in eyes	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing</li> <li>Call a poison control center or doctor for treatment advice</li> </ul>
If swallowed	<ul style="list-style-type: none"> <li>Call a poison control center or doctor for treatment advice</li> <li>Have person sip a glass of water if able to swallow</li> <li>Do not induce vomiting unless told to by a poison control center or doctor</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>Take off contaminated clothing</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>Call a poison control center or doctor for treatment advice</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>Move person to fresh air</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible</li> <li>Call a poison control center or doctor for treatment advice</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN:</b> Probable mucosal damage may contraindicate the use of gastric lavage.	

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin, or inhaled. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Remove contaminated clothing and wash clothing before reuse.

### Personal Protective Equipment (PPE)

Applicator and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical resistant gloves (such as water proof)
- Goggles or face shield

(BACK PANEL)  
COMMERCIAL USE ONLY  
*(for internal use only-will not appear on final label)*

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations:** Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to freshwater marine and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**  
Do not contaminate water, food or feed by storage and disposal.  
**PESTICIDE STORAGE:** Keep container closed and away from food, feedstuffs, and domestic water supplies.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

**GENERAL INFORMATION**

This product has a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass), and green and yellow foxtail. This product controls these annual weeds by killing their seeds as they germinate. Apply this product before these annual weeds begin to germinate since it does not control established weeds.

When this product is applied as directed, the following established turfgrasses are tolerant:

- |                    |            |                    |                    |
|--------------------|------------|--------------------|--------------------|
| Kentucky bluegrass | Fescue     | St. Augustinegrass | Perennial ryegrass |
| Zoysiagrass        | Bahiagrass | Centipede          | Bermudagrass       |

(BACK PANEL)  
COMMERCIAL USE ONLY  
(for internal use only-will not appear on final label)

**RESEEDING**

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying this product. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as Rogers Aero Seeder) should be employed. When using this product at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

**SUGGESTED SPREADER SETTINGS**

Spreader	Setting	Pattern	Width
Cyclone Broadcast B1-84	4 1/2		4
Earthway Broadcast 2400	17		10
Lely W	6	III-B	52
Lesco Broadcast	G 1/2	0.25	10
Prizelawn Broadcast CB85	4 1/2	C	9
Scotts Broadcast R8A	L 1/2	3	11
Spyker	4.3		4
Vicon 402	37		40

**GENERAL PRECAUTIONS AND RESTRICTIONS**

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Apply this product to the turf prior to the germination of weed grasses since it will not control established weeds. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of this product should not be used on putting greens. Not for use on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

**Entry Restrictions:** Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow other to enter the treated area until the surface is dry after the watering-in.

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

**APPLICATION INSTRUCTIONS**

**RATES, FREQUENCY AND TIMING OF APPLICATION**

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply this product approximately one or two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results

(BACK PANEL)  
COMMERCIAL USE ONLY  
*(for internal use only-will not appear on final label)*

may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

- 1. Spring germinating summer annual grasses, crabgrass, foxtails and barnyardgrass.
  - A. Northern (cool season) turfgrasses  
For preemergence control, apply 3.75 - 5 pounds per thousand square feet in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.
  - B. Southern (warm season) turfgrasses  
For preemergence control, apply 7.5 pounds per thousand square feet in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of this product from one half to the full recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

- 2. Late spring, early summer germinating annual grasses - goosegrass.  
Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications of this product should, therefore, be timed accordingly.
  - A. Northern (cool season) turfgrass  
For preemergence control, apply 3.75 - 5 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
  - B. Southern (warm season) turfgrass  
For preemergence control, apply 7.5 pounds per thousand square feet in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

- 3. Northern and Southern turfgrasses  
  
In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications of from one half to the full recommended rate 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

- 4. *Poa annua* (annual bluegrass)  
In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseedling will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

6/6

(BACK PANEL)  
COMMERCIAL USE ONLY  
*(for internal use only-will not appear on final label)*

- A. Northern (cool season) turfgrasses  
For preemergence control, apply 3.75 - 5 pounds per thousand square feet in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.
  
- B. Southern (warm season) turfgrasses  
For preemergence control, apply 7.5 pounds per thousand square feet in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application of this product from 3.75 to 7.5 pounds per thousand square feet can be made 10 to 12 weeks after the initial treatment.

**IMPORTANT: READ BEFORE USE:**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER and LIMITATION of LIABILITY:**

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS:**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.(RVamd111302)

Manufactured by: The Andersons Lawn Fertilizer Division  
d/b/a Free Flow Fertilizer  
PO Box 119  
Maumee, OH 43537

Date Printed	11/13/02
Date Approved	
Supersedes	9/7/00
Label change	Amendment - first aid, fert, warranty
Filename	9198-94C
Labelcode	RVamd111302