

MAR 20 1995

Mr. Daniel Paradiso, Jr.  
The Andersons Lawn Fertilizer Division  
DBA Free Flow Fertilizer  
P.O. Box 119  
Maumee, OH 43537

Dear Mr. Paradiso:

Subject: The Andersons Tee Time With Team/Dursban I  
EPA Registration No. 9198-98  
Application Dates March 25, 1994, Request To  
Revise Labeling for Homeowner and Commercial  
Uses, Compliance with PR Notices 93-3, 93-7 and  
93-11 and Label Format Revisions

The proposed amendments to the subject pesticide product labeling have been reviewed and are acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) as amended provided that you:

1. Add the exclusionary statement "not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes" to descope this product from the EPA Worker Protection Standard. This statement must be added to both the homeowner and commercial labels.
2. Submit one (1) copy of the final printed labels before releasing the product for shipment under the subject labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Joanne I. Miller  
Product Manager (23)  
Fungicide-herbicide Branch  
Registration Division (7505C)

enclosures

Wilson: ABC8: 03-20-95

2911  
-7

(FRONT PANEL)

COMMERCIAL USE ONLY

THE ANDERSONS TEE TIME WITH TEAM/DURSBAN I

(Logo)

ACTIVE INGREDIENTS:

N-butyl-N-ethyl--,-,--trifluoro-2,6-dinitro-p-toluidine . . . . .	0.77%
Trifluralin: alpha, alpha, alpha-trifluoro-2,6-dinitro- N,N-dipropyl-p-toluidine . . . . .	0.38%
Chlorpyrifos [0,0-diethyl 0-(2,4,6-trichloro- 2-pyridyl)phosphorothioate] . . . . .	0.57%
INERT INGREDIENTS . . . . .	98.28%
Total	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements

Net Weight 50 lbs. (22.7 kg.)

Treats Up to 12,500 Sq. Ft.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
MAR 20 1995

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

9198-98

EPA Reg. No 9198-98

EPA Est. 9198-OH-1

1162

3911  
y

(BACK PANEL)

COMMERCIAL USE ONLY

Net Weight 50 lbs. (22.7 kg.)

Treats Up To 12,500 Sq. Ft.

THE ANDERSONS TEE TIME WITH TEAM/DURSBAN I

GUARANTEED ANALYSIS

Total Nitrogen (N) .....	25%
1.2% Ammoniacal Nitrogen	
17.8% Urea Nitrogen	
6.0% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P2O5) .....	3%
Soluble Potash (K2O) .....	8%

Plant Nutrients Derived From: Ammoniated Phosphates, Urea, Sulfur Coated Urea, and Muriate of Potash.

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with eyes. If in eyes, flush with plenty of water. Harmful if swallowed, inhaled or absorbed through the skin. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.

1163

4 2 11

(BACK PANEL)

COMMERCIAL USE ONLY

- IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching finger to the back of throat. Never give anything by mouth to an unconscious or convulsing person.
- IF ON SKIN: Wash with soap and plenty of water. Get medical attention if irritation persists.
- IF INHALED: If breathing difficulty occurs, move individual to fresh air and get medical attention.
- NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote by injection only.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when cleaning or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is in violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a combination of growth promoting fertilizer ingredients, a preemergent herbicide, and a turf insecticide. It is designed to control smooth and hairy crabgrass, goosegrass (silver crabgrass or crowsfoot), barnyard grass (watergrass), green and yellow foxtail, and annual bluegrass (*Poa Annua*), as well as ants, armyworms, billbugs, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers, and sod webworms.

When applied as directed, the product can be used on the following turfs: Kentucky Bluegrass, perennial ryegrasses, centipedegrass, fescue, zoysiagrass, bermudagrass, St. Augustinegrass, and bahiagrass. The product will not harm nearby established ornamental trees and shrubs.

4

5 2 11

(BACK PANEL)

COMMERCIAL USE ONLY

#### APPLICATION DIRECTIONS

Apply at the rate of 4 pounds per 1,000 sq. ft. Because this product is generally applied very early in the insect cycle, it may be necessary to give a second treatment of insecticide late in the season.

#### USE PRECAUTIONS

Apply this product to the turf prior to the germination of weed grasses since it will not control established weeds. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of this product should not be used on putting greens. This product should not be used on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

#### RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying this product. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as Rogers Aero Seeder) should be employed. When using this product, reseeding should be delayed 12 to 16 weeks after application.

#### APPLICATION EQUIPMENT

Apply this product evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area and then applying the remaining half at the right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly.

69 11

(BACK PANEL)

COMMERCIAL USE ONLY

SUGGESTED SPREADER SETTINGS

(This section will be completed for final labeling)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured By: The Andersons  
Lawn Fertilizer Division  
P.O. Box 109  
Maumee, OH 43537

Date Printed: 01/24/94  
Date Approved:  
Supersedes:  
Label Change:

79 11

(FRONT PANEL)

HOMEOWNER USE ONLY

THE ANDERSONS TEE TIME WITH TEAM/DURSBAN I

(Logo)

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-- , - - -trifluoro-2,6-dinitro-p-toluidine . . . . .	0.77%
Trifluralin: alpha, alpha, alpha-trifluoro-2,6-dinitro- N,N-dipropyl-p-toluidine . . . . .	0.38%
Chlorpyrifos [0,0-diethyl 0-(2,4,6-trichloro- 2-pyridyl)phosphorothioate] . . . . .	0.57%
INERT INGREDIENTS . . . . .	98.28%
TOTAL. . . . .	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements

Net Weight 50 lbs. (22.7 kg.)

Treats Up to 12,500 Sq. Ft.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
MAR 20 1995

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

9198-98

EPA Reg. No. 9198-98

EPA Est. 9198-CH-1

8911

(BACK PANEL)

HOMEOWNER USE ONLY

Net Weight 50 lbs. (22.7 kg.)

Treats Up To 12,500 Sq. Ft.

THE ANDERSONS TEE TIME WITH TEAM/DURSBAN I

GUARANTEED ANALYSIS

Total Nitrogen (N) .....	25%
1.2% Ammoniacal Nitrogen	
17.8% Urea Nitrogen	
6.0% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid (P2O5) .....	3%
Soluble Potash (K2O) .....	8%

Plant Nutrients Derived From: Ammoniated Phosphates, Urea, Sulfur Coated Urea, and Muriate of Potash.

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with eyes. If in eyes, flush with plenty of water. Harmful if swallowed, inhaled or absorbed through the skin. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician



(BACK PANEL)

HOMEOWNER USE ONLY

- IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching finger to the back of throat. Never give anything by mouth to an unconscious or convulsing person.
- IF ON SKIN: Wash with soap and plenty of water. Get medical attention if irritation persists.
- IF INHALED: If breathing difficulty occurs, move individual to fresh air and get medical attention.
- NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote by injection only.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when cleaning or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is in violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a combination of growth promoting fertilizer ingredients, a preemergent herbicide, and a turf insecticide. It is designed to control smooth and hairy crabgrass, goosegrass (silver crabgrass or crowsfoot), barnyard grass (watergrass), green and yellow foxtail, and annual bluegrass (Poa Annua), as well as ants, armyworms, billbugs, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers, and sod webworms.

When applied as directed, the product can be used on the following turfs: Kentucky Bluegrass, perennial ryegrasses, centipedegrass, fescue, zoysiagrass, bermudagrass, St. Augustinegrass, and bahiagrass. The product will not harm nearby established ornamental trees and shrubs.

9

10 9 11  
y

(BACK PANEL)

HOMEOWNER USE ONLY

#### APPLICATION DIRECTIONS

Apply at the rate of 4 pounds per 1,000 sq.ft. Because this product is generally applied very early in the insect cycle, it may be necessary to give a second treatment of insecticide late in the season.

#### USE PRECAUTIONS

Apply this product to the turf prior to the germination of weed grasses since it will not control established weeds. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of this product should not be used on putting greens. Not for use on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

#### RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying this product. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as Rogers Aero Seeder) should be employed. When using this product, reseeding should be delayed 12 to 16 weeks after application.

#### APPLICATION EQUIPMENT

Apply evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area and then applying the remaining half at the right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly.

#### SUGGESTED SPREADER SETTINGS

(AVAILABLE AT FINAL PRINTING)

(BACK PANEL)

HOMEOWNER USE ONLY

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured By: The Andersons  
Lawn Fertilizer Division  
P.O. Box 119  
Maumee, OH 43537

Date Printed: 01/24/94  
Date Approved:  
Supersedes:  
Label Change: