

APR 15 1993

Mr. Daniel Paradiso
The Andersons Management Corporation
P.O. Box 119
Maumee, Ohio 43537

Dear Mr. Paradiso:

Subject: Updated and Expanded Labeling
The Andersons Tee Time Fertilizer with Sevin
EPA Registration Number 9198-106
Your Submission Dated January 18, 1993

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the following label changes before you release the product for shipment:

a. Change "poising" to "poisoning" on first page under Statement of Practical Treatment.

b. Add use rates for commercial application if you intend to market this product for that purpose.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

Submit five (5) copies of your final printed label before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide Rodenticide Branch
Registration Division (H7505C)

Enclosure

9198-106
Page 1 of 7

(FRONT PANEL)

THE ANDERSONS TEE TIME
FERTILIZER WITH SEVIN®

ACTIVE INGREDIENT:	
Carbaryl (1-naphtyl N-methylocarbamate)	6.2%
INERT INGREDIENTS	93.8%
Total 100.00%	

SEVIN® is a registered trademark of Rhone-Poulenc Nederland BV for carbaryl insecticides.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Net Weight _____ lbs.

Treats _____ Sq. Ft.


ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 15 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
9198-106

EPA Reg. No. 9198-106

EPA Est. 9198-OH-1



(BACK PANEL)

Net Weight _____ lbs.

Treats _____ Sq. Ft.

THE ANDERSONS TEE TIME

FERTILIZER WITH SEVIN®

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HARMFUL IF SWALLOWED OR ABSORBED THROUGH THE SKIN. OVEREXPOSURE MAY CAUSE: Salivation, watery eyes, pinpoint eye pupils, blurred vision, muscle tremors, difficult breathing, excessive sweating/ abdominal cramps, nausea, vomiting, diarrhea, weakness, headache. In severe cases, convulsion, unconsciousness and respiratory failure may occur. Signs and symptoms occur rapidly following overexposure to this product.

Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling this product. Change to clean clothing daily. Bathe and wash hair after each work day. Do not eat, drink or use tobacco while working with this product. Wash hands and face before eating, drinking or using tobacco. Keep out of reach of children and domestic animals.

STATEMENT OF PRACTICAL TREATMENT

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

- IF SWALLOWED: Never give anything by mouth to mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger.
- IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.
- IF ON SKIN: Wash thoroughly with soap and water.
- IF INHALED: Move from contaminated atmosphere and call a physician.

BEST AVAILABLE COPY

ANTIDOTE STATEMENT: ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended. See NOTE TO PHYSICIAN below.

NOTE TO PHYSICIAN: Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible. Gastric lavage may be used if this product has been swallowed. Carbaryl poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Atropine is antidotal. Narcotics and other sedatives should not be used. Further drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water and areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning equipment or disposal of wastes. Do not contaminate water when disposing of equipment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LAWNS, RECREATIONAL, AND ORNAMENTAL TURF AREAS: For control of turfgrass pests, apply granules uniformly to the turfgrass. For optimum control, follow directions for proper watering or irrigation after application. Do not allow public use of treated area during application.

INSECT	POUNDS PRODUCT 1000 SQ. FT.	SPECIFIC DIRECTIONS
Ants, Armyworms, Centipedes, Chiggers, Chinch Bugs, Crickets, Cutworms, Darkling Ground Beetles, Earwigs, Essex Skipper, Fall Armyworms, Fiery Skipper, Fleas, Grasshoppers, Green June Beetles, Green June Beetle Grubs, Lawn Moths (Sod Webworm), Leafhoppers, Lucerne Moth Larvae, Millipedes, Sowbugs, Spittlebugs, Springtails, Ticks, Yellowstriped Armyworms	2.2 TO 3.0	For general insect control apply granules uniformly when grass is dry. For best results lawn should be watered lightly after application. Repeat applications may be made as needed for control of new infestations.

BEST AVAILABLE COPY

POUNDS OF TEE TIME
FERTILIZER WITH
SEVIN (R) PER
1000 SQ. FT.

INSECT

SPECIFIC DIRECTIONS


White Grubs (such as Japanese Beetle Grubs, Chafer Beetles, Phyllophaga spp. grubs and Annual White Grubs, Bluegrass billbugs larvae, European crane-fly larvae

3.0

For white grub control, applications should be made when grubs are feeding near the soil surface, usually during late March through May or July to early September, or as recommended by local agricultural extension agents. For the best results the material should be watered in sufficiently to wash the insecticide into the zone of insect infestation. Repeat applications may be made as needed for control of new infestations.

NUISANCE PEST CONTROL IN AND AROUND FLOWER BEDS AND ORNAMENTAL PLANTINGS AND AROUND BUILDINGS

For control of ants, armyworms, ticks, cutworms, centipedes, crickets, earwigs, firebrats and silverfish, fleas, millipedes, sowbugs, scorpions and spiders; apply granules uniformly in a 6 foot band around the building. Apply granules at the rate of 5.8 ounces per 120 square feet (6 x 20 foot band) to shrub and flower beds, foundations, ornamental plantings, and lawn or soil areas immediately adjacent to the building. For best results, treated area should be lightly watered after application to remove the chemical from the granules. Repeat treatments as necessary to control new infestations.



BEST AVAILABLE COPY

FRUIT AND VEGETABLE CROPS (Not for Use in California)

For control of Ants, Centipedes, Crickets, Earwigs, Grasshoppers, Millipedes, Sowbugs (Pillbugs), Springtails, and Ticks, apply 0.75 lb. per 1000 ft of SEVIN® brand Granular evenly over the soil surface. For best results water lightly after application to release the chemical from the granules. Applications may be made at any time before, during or after the growing season. If applying during the growing season, observe indicated pre-harvest intervals. (PHI).

0 day PHI - Beans, Blueberries, Carrots, Corn, Cucumber, Eggplant, Melons, Okra, Peanuts, Pepper, Potato, Pumpkins, Squash, Sweet Potato, Tomato

1 day PHI - Asparagus, Strawberries

3 day PHI - Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Cowpeas, Garden Beet Root, Head Lettuce, Kohlrabi, Parsnips, Peas, Radishes, Rutabagas, Salsify Roots, Turnip Roots

7 day PHI - Blackberries, Dewberries, (including boysenberries and loganberries), Grapes, Raspberries

14 day PHI - Celery, Chinese Cabbage, Collards, Dandelion, Endive (Escarole), Garden beet Tops, Hanove Salad, Kale, Leaf Lettuce, Mustard Greens, Parsley, Salsify Tops, Spinach, Swiss Chard, Turnip Tops

IMPORTED FIRE ANT CONTROL

For use as a mound treatment to control Imported fire ants, apply 1.5 to 3.0 ounces by weight over and around each fire ant mound. Irrigate mound thoroughly immediately after treatment to remove the chemical from the granules and flush it down into the mound. A minimum of 1 gallon of water should be applied to each mound, but do not use more than 2 gallons on water as flooding of the mound may result. Apply water gently to avoid disturbing ant and causing migration before chemical can be effective. For best results, apply in the early morning or late evening hours when ants are least active. Treat new mounds as they appear.

CONTROL OF TICKS WHICH TRANSMIT LYME DISEASE

For control of juvenile and adult ixodes spp. ticks (Deer tick, Bear tick, and black-legged tick); and Amblyomma spp. ticks (Lone star tick). Apply granules at the rate of 2.5 to 3.0 pounds of formulated product per 1000 square feet.

Begin applications in late spring or early summer (May/June). Uniformly apply granules to entire lawn, perimeter wooded areas, and property boundaries including shrub beds, ornamental plantings, and wooded areas where exposure to ticks may occur. For best results, irrigate treated areas after application to remove chemical from granules.

BEST AVAILABLE COPY

SUGGESTED SPREADER SETTINGS
(AVAILABLE AT FINAL PRINTING)

IMPORTANT: THE ABOVE SPREADER SETTINGS ARE APPROXIMATE. DIFFERENCES MAY OCCUR DUE TO CONDITION OF SPREADER, SPEED AT WHICH APPLIED AND ROUGHNESS OF TERRAIN. CALCULATE THE SIZE OF THE AREA TO BE TREATED AND DO NOT OVER OR UNDER APPLY. ACCURATELY WEIGH A FEW POUNDS INTO YOUR SPREADER AND APPLY TO A MEASURED AREA. THEN ADJUST THE SPREADER SETTING, IF NECESSARY, IN ORDER TO ASSURE CORREC. RATE OF APPLICATION. CLOSE SPREADER WHEN FILLING, STOPPING OR TURNING, DO NOT HAND SPREAD.


STORAGE AND DISPOSAL

(THIS STATEMENT WILL BE USED IF SOLD IN THE HOMEOWNER MARKET)

Do not contaminate water, food, or feed by storage or disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

(THIS STATEMENT WILL BE USED IF SOLD IN THE COMMERCIAL MARKET)

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes can not be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, burning. If burned, stay out of smoke.



BEST AVAILABLE COPY

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop condition, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured By: The Andersons
Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537

Date Printed: 01/18/93
Date Approved:
Supersedes:
Label Change:

