

9198-106

03-24-2004

111

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-9

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

**Application for Pesticide - Section I**

1. Company/Product Number 9198-106	2. EPA Product Manager  PM#	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Tee Time Fertilizer with Seving		
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fertilizer Division, Inc. d/b/a Free Flow Fertilizer PO Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____  Product Name _____	

**Section - II**

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

**NOTIFICATION**  
MAR 24 2004

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of Minor Label change per PR Notices 2001-6 and 98-10 : optional front panel claims, guarantee

This notification is consistent with the provisions of PR Notices 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.


**Section - III**

<b>1. Material This Product Will Be Packaged In:</b>				<b>2. Type of Container</b>	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
<b>3. Location of Net Contents Information</b> <input type="checkbox"/> Label <input type="checkbox"/> Container		<b>4. Size(s) Retail Container</b>		<b>5. Location of Label Directions</b> <input type="checkbox"/> On Label	
<b>6. Manner in Which Label is Affixed to Product</b>			<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		
			<input type="checkbox"/> Other _____		

**Section - IV**

**1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)**

Name Dawn L. Walters	Title Regulatory Administrator	Telephone No. (include Area Code) 334-262-2766
-------------------------	-----------------------------------	---

<b>Certification</b>			6. Date Application Received  (Stamped)
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			
2. Signature 	3. Title Regulatory Administrator		
4. Typed Name Dawn L. Walters	5. Date 2/26/04		

e/11

(FRONT PANEL)

# THE ANDERSONS TEE TIME FERTILIZER WITH SEVIN®

**ACTIVE INGREDIENT:**

Carbaryl (1-naphtyl N-methylcarbamate) ..... 6.2%

**INERT INGREDIENTS** ..... 93.8%

Total 100.00%

**NOTIFICATION**  
MAR 24 2004

SEVIN® is a registered trademark of Aventis Crop Protection

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**SEE BACK PANEL FOR ADDITIONAL PRECUATIONS**

OR

**SEE BACK PANEL FOR PRECAUTIONARY STATEMENTS**

Net Weight \_\_\_\_\_ lbs.

Treats \_\_\_\_ Sq. Ft.

*Optional marketing statements:*

- Controls: White Grubs, Chinch Bugs, Sod Webworms, Ticks, Ants, Fleas & more
- Contains Poly-S® controlled release nitrogen for extended greening
- (For best results, use with other lawn care products in the Sam's Choice™ Fertilizer Program)
- Fertilizes lawns & controls insects in one step
- For lawns
- Soil & Turf insect control while fertilizing your lawn
- Controls lawn damaging surface and soil feeding insects while feeding your lawn with a premium blend of fertilizer that provides quick and extended greening.
- Soil & Turf Insect Control Plus Fertilizer
- For insect control on lawns
- Contains controlled release nitrogen for long-lasting feeding of your lawn

EPA Reg. No. 9198-106

EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code on bag

3/11

(BACK PANEL)

Net Weight \_\_\_\_\_ lbs.  
Treats \_\_\_\_ Sq. Ft.

### THE ANDERSONS TEE TIME FERTILIZER WITH SEVIN®

**Guaranteed Analysis**

Total Nitrogen (N) .....	XX%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%

Plant nutrients derived from:

*For products being sold into Washington state:*

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-ap.com/>

**KEEP OUT OF REACH OF CHILDREN****CAUTION****PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing before reuse. Do not eat, drink or use tobacco while working with this product.

**FIRST AID**

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN**. If poisoning is suspected in animals, contact a veterinarian.

If swallowed	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If on skin	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.	

**NOTE TO PHYSICIAN:**

Carbaryl is a N-methyl carbamate insecticide that inhibits cholinesterase. Atropine is antidotal. Do not use 2-PAM (pyridine-2-aldoxime methiodide), opiates or cholinesterase inhibiting drugs. Probable mucosal damage may contraindicate the use of gastric lavage. Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24

4/11

(BACK PANEL)

hours. To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen.

**ANTIDOTE STATEMENT: ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE.**

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are **NOT** recommended. See **NOTE TO PHYSICIAN** below.

**ENVIRONMENTAL HAZARDS**

***For Homeowner Use:***

This product is toxic to fish, birds and other wildlife. Bees exposed to direct treatment or residues on plants are highly susceptible. Do not apply directly to water. Do not contaminate water by disposing of equipment washings. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

***For Commercial Use:***

This product is toxic to fish, birds and other wildlife. Bees exposed to direct treatment or residues on plants are highly susceptible. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washings. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

**LAWN, RECREATIONAL, AND ORNAMENTAL TURF AREAS:**

For control of turfgrass pests, apply granules uniformly to the turfgrass. For optimum control, follow directions for proper watering or irrigation after application. Do not allow public use of treated area during application.

(BACK PANEL)

PEST		LBS OF PRODUCT PER 1,000 SQ. FT.	SPECIFIC DIRECTIONS
Ants Armyworms Centipedes Chiggers Chinch Bugs Crickets Cutworms Darkling Ground Beetles Earwigs Essex Skipper Fall Armyworms Fiery Skipper Fleas Grasshoppers	Green June Beetle Green June Beetle Grubs Lawn Moths (Sod Webworm) Leafhoppers Lucrene Moth Larvae Millipedes Sowbugs Spittlebugs Springtails Ticks Yellowstriped Armyworms	2.2 to 3.0	For general insect control, apply granules uniformly when grass is dry.  For best results, lawn should be watered lightly after application.  Repeat applications may be made as needed for control of new infestations.
White Grubs (such as Japanese Beetle Grubs, Chafer Beetles, <i>Phyllophaga</i> spp. Grubs and Annual White Grubs) Bluegrass billbugs larvae European crane fly larvae		3.0	For white grub control, applications should be made when grubs are feeding near the soil surface, usually during late March through May or July to early September, or as recommended by local agricultural extension agents.  For the best results the material should be watered in sufficiently to wash insecticide into the zone of insect infestation.  Repeat applications may be made as needed for control of new infestations.

**IMPORTED FIRE ANT CONTROL**

For use as a mound treatment to control Imported fire ants, apply 1.5 to 3.0 ounces by weight over and around each fire ant mound. Irrigate mound thoroughly immediately after treatment to remove the chemical from the granules and flush it down into the mound. A minimum of 1 gallon of water should be applied to each mound, but do not use more than 2 gallons on water as flooding of the mound may result. Apply water gently to avoid disturbing ant and causing migration before chemical can be effective. For best results, apply in the early morning or late evening hours when ants are least active. Treat new mounds as they appear.

**CONTROL OF TICKS WHICH TRANSMIT LYME DISEASE**

For control of juvenile and adult ixodes spp. ticks (Deer tick, Bear tick, and Black legged tick). Apply granules at the rate of 2.5 to 3.0 pounds of formulated product per 1000 square feet.

Begin applications in late spring or early summer (May/June). Uniformly apply granules to entire lawn, perimeter wooded areas, and property boundaries including shrub beds, ornamental plantings, and wooded areas where exposure to ticks may occur. For best results, irrigate treated areas after application to remove chemical from granules.

(BACK PANEL)

**NUISANCE PEST CONTROL IN AND AROUND FLOWER BEDS AND ORNAMENTAL PLANTINGS AND AROUND BUILDINGS**

For control of ants, armyworms, ticks, cutworms, centipedes, crickets, earwigs, firebrats and silverfish, fleas, millipedes, sowbugs, scorpions and spiders; apply granules uniformly in a 6 foot band around the building. Apply granules at the rate of 5.8 ounces per 120 square feet (6 x 20 foot band) to shrub and flower beds, foundations, ornamental plantings, and lawn or soil areas immediately adjacent to the building. For best results, treated area should be lightly watered after application to remove the chemical from the granules. Repeat treatments as necessary to control new infestations.

**FRUIT AND VEGETABLE CROPS (Not for Use in California)**

For control of Ants, Centipedes, Crickets, Earwigs, Grasshoppers, Millipedes, Sowbugs (Pillbugs), Springtails, and Ticks, refer to the Use Rate chart below for application information for various commodities. Apply recommended rate of product evenly over the soil surface. For best results water lightly after application to release the chemical from the granules. Applications may be made at any time before, during or after the growing season. If applying during the growing season, observe indicated pre-harvest intervals. (PHI).

**Use Rates for The Andersons Tee Time Fertilizer with Sevin**

Commodity	Use Rate
Asparagus	<p><b>Application to Spears:</b></p> <ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.2 lbs per 1,000 sq. ft. (6 lb ai/acre) per crop</li> </ul> <p><b>Application to Ferns (after harvest of spears):</b></p> <ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 1.4 lbs per 1,000 sq.ft. (4 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days when applied to ferns after harvest</li> <li>• Retreatment interval = 3 days when application are to spears</li> <li>• Do not apply within 1 day of harvest</li> <li>• The total maximum seasonal application rate for spears and ferns is 3.7 lbs per 1,000 sq. ft. (10 lb ai/acre) per crop</li> <li>• The total number of application is 3 prior to harvest with 2 additional applications to ferns only</li> </ul>
Beans, fresh and dried (snap, navy and kidney); pea, fresh and dried: cowpea, lentil	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.5 lbs per 1,000 sq. ft. (1.5 lb ai/acre) per application</li> <li>• Maximum season application rate of 2.2 lbs per 1,000 sq. ft. (6 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 3 days of harvest (succulents)</li> <li>• Do not apply within 21 days of harvest (dried)</li> <li>• Do not apply within 14 days of harvest (forage)</li> <li>• Do not apply within 21 days of harvest (hay)</li> </ul>

(BACK PANEL)

<p>Garden beets (roots), carrots, horseradish, radish, parsnip, rutabaga, salsify, turnips (roots)</p>	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.2 lbs per 1,000 sq. ft. (6 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 7 days of harvest</li> <li>• Total number of applications per crop can be 6 (some of which must be at lower than maximum single application rate)</li> </ul>
<p>Mustard greens</p>	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.5 lb per 1,000 sq.ft. (1.5 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.2 lb per 1,000 sq.ft. (6 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 13 to 14 days of harvest</li> <li>• Total number of applications/crop can be 4 (some of which must be at lower than maximum single application rate)</li> </ul>
<p>Brussels sprouts, broccoli, cabbage, cauliflower, Chinese cabbage, collards, kale, kohlrabi</p>	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.2 lbs per 1,000 sq. ft. (6 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 3 days of harvest of Brussels sprouts, broccoli, cabbage, kohlrabi, cauliflower</li> <li>• Do not apply within 14 days of Chinese cabbage, kale, collards</li> <li>• Total number of applications/crop can be 4 (some of which must be at lower than maximum single application rate)</li> </ul>
<p>Corn (field and pop)</p>	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.9 lbs per 1,000 sq. ft. (8 lb ai/acre) per crop</li> <li>• Do not apply within 14 days of harvest (forage, silage)</li> <li>• Do not apply within 48 days of harvest (grain, fodder)</li> </ul>
<p>Corn (sweet)</p>	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 5.9 lbs per 1,000 sq. ft. (16 lb ai/acre) per crop</li> <li>• Retreatment interval = 3 days</li> <li>• Do not apply within 2 days of harvest (ears)</li> <li>• Do not apply within 14 days of harvest (forage)</li> <li>• Do not apply within 48 days of harvest (fodder)</li> </ul>
<p>Cucumber, melon, squash</p>	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.3 lbs per 1,000 sq. ft. (1 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.2 lbs per 1,000 sq. ft. (6 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 3 days of harvest</li> </ul>

(BACK PANEL)

Lettuce, head and leaf; endive, celery, dandelion, parsley, spinach, Swiss chard, Hanover salad, garden beets (tops), turnips (tops)	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.2 lbs per 1,000 sq. ft. (6 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 14 to 15 days of harvest</li> <li>• Total number of applications/crop can be 5 (some of which must be at lower than maximum single application rate)</li> </ul>
Potato	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.2 lbs per 1,000 sq. ft. (6 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 7 days of harvest</li> <li>• Total number of applications/crop can be 6 (some of which must be at lower than maximum single application rate)</li> </ul>
Strawberry	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 3.7 lbs per 1,000 sq. ft. (10 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 7 days of harvest</li> </ul>
Tomato, pepper, eggplant	<ul style="list-style-type: none"> <li>• Maximum single application rate of 0.7 lbs per 1,000 sq. ft. (2 lb ai/acre) per application</li> <li>• Maximum seasonal application rate of 2.9 lbs per 1,000 sq. ft. (8 lb ai/acre) per crop</li> <li>• Retreatment interval = 7 days</li> <li>• Do not apply within 3 days of harvest</li> <li>• Total number of applications/crop can be 7 (some of which must be at lower than maximum single application rate)</li> </ul>

**SUGGESTED SPREADER SETTINGS**

*(AVAILABLE AT FINAL PRINTING)*

**IMPORTANT:** The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of terrain. Calculate the size of the area to be treated and do not over or under apply. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary, in order to assure correct rate of application. Close spreader when filling, stopping or turning, do not hand spread.

**Broadcast spreader {setting}s**

For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

*[diagram of path] [listing of settings]*

**Drop spreader {setting}s**

First apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.



9/11

(BACK PANEL)

[diagram of path] [listing of settings]

To determine square footage of application area, follow calculation chart. [lawn/turf diagram]

\_\_\_ length  
x \_\_\_ width  
= \_\_\_ Total Sq. Ft.  
- \_\_\_ Non-lawn areas  
= \_\_\_ Total Sq. Ft.

**STORAGE AND DISPOSAL**

**For Homeowner Use:**

Do not contaminate water, food or feed by storage and disposal.

**STORAGE:** Keep container closed and away from food, feedstuffs, and domestic water supplies.

**DISPOSAL:**

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If party filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**For Commercial Use:**

Do not contaminate water, food or feed by storage and disposal.

**STORAGE:** Keep container closed and away from food, feedstuffs, and domestic water supplies.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, burning. If burned, stay out of smoke.

**IMPORTANT: READ BEFORE USE:**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER and LIMITATION of LIABILITY:**

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS:**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVnot022504)

10711

(BACK PANEL)

Optional guarantee:

{brand} Satisfaction Guarantee:

If, for any reason, you are not completely satisfied with this product, our money will be refunded. Send proof of purchase to the address below with explanation of your problem.

**Manufactured By:**

The Andersons Lawn Fertilizer Division  
d/b/a Free Flow Fertilizer  
P.O. Box 119  
Maumee, OH 43537

**Optional marketing information:**

XX% Nitrogen – Gives lawns their green color. Fertilizers with controlled-release nitrogen formulas will feed and green your lawn over a longer period of time

X% Phosphate – This nutrient is important for root development. Good roots are vital for grass to take up water and nutrients

XX% Potassium – Also called potash, helps lawns better tolerate extreme weather conditions. Contributes to the overall vigor of grass

Sam Walton, founder of Wal-Mart, listened to his customers, understood their needs, and found ways to offer quality products to them at every day low prices.

The Sam's Choice brand continues his legacy by offering innovative products from around the world. We believe these products, available only at Wal-Mart and Wal-Mart Supercenters, offer exceptional quality and better value than the leading national brands at our always low Wal-Mart prices. We think you'll agree or we'll refund your money, no questions asked!

"Better products at the always low Wal-Mart prices..."


Always. We guarantee it. Sam Walton

Questions, Comments, Information? 1-888-246-0024

**Application Guidelines**

For best results, apply only one product during one period of time

ZONE WHERE YOU LIVE (See map)	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT
NORTHERN ZONE										
TRANSITIONAL ZONE										
SOUTHERN ZONE										



Types of grasses common to each zone are:  
 NORTHERN: Bluegrasses, Ryegrasses, Bentgrasses, Fine Fescue, Tall Fescue  
 TRANSITIONAL: Bluegrasses, Ryegrasses, Fine Fescue, Tall Fescue, Bermudagrass, Zoysiagrass  
 SOUTHERN: Bermudagrass, Zoysiagrass, Centipedegrass, Bahiagrass and St. Augustine

\*Note: This guide estimates the timing of lawn product applications and is a guideline only. In Transition Zones, be sure to apply the correct Weed & Feed for your grass type. Climatic conditions determine the of applying various product types. Consult a lawn and garden associate or your local county extension your product needs and the appropriate timing to apply these products to your lawn.

Date Printed	2/25/04
Date Approved	8/12/03
Supersedes	12/16/99
Label Changes	Notification – front claims, opt. guarantee
Filename	9198-106
Label Code	RVnot022504

11/11



**The Andersons, Inc.**

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

February 26, 2004

Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7505C)  
US Environmental Protection Agency  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, DC 20460-0001

RE: Notification of Minor Label Change  
SUBJECT: The Andersons Tee Time Fertilizer with Sevin  
EPA Reg. No. 9198-106

Per PR Notice 98-10 enclosed is an EPA Form 8570-1, an application notifying the Agency of a revision to the labeling for the subject registration. One copy of the revised labeling is also enclosed with the additions underlined and marked in blue.

We have made the following changes by notification:

- 1. Added optional front panel claims

For insect control on lawns  
Contains controlled release nitrogen for long-lasting feeding of your lawn

- 2. Added optional guarantee

If, for any reason, you are not completely satisfied with this product, your money will be refunded. Send proof of purchase to the address below with explanation of your problem.

Enclosed is a card indicating the acceptance or rejection of this notification. Please complete the card and return via mail. If there are any questions or comments regarding this notification, please feel free to contact me.

Sincerely,  
The Andersons Lawn Fertilizer Division, Inc.

*Dawn L. Walters*  
Dawn L. Walters  
Regulatory Administrator  
Phone 334-262-2766  
Fax 334-265-5703  
Email dawn\_walters@andersonsinc.com

enclosures