

U.S. ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (75-757)  
WASHINGTON, DC 20460

9198-108

AUG 2 1989

**NOTICE OF PESTICIDE:** ☒ REGISTRATION  
☐ REREGISTRATION  
(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

TERM OF REGISTRATION  
Conditional

NAME OF PESTICIDE PRODUCT

1.0% Aqueous Solution of 2,4-D  
1.0% Aqueous Solution of 2,4-D

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

1.0% Aqueous Solution of 2,4-D  
1.0% Aqueous Solution of 2,4-D  
1.0% Aqueous Solution of 2,4-D  
1.0% Aqueous Solution of 2,4-D

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit to all data required for registration, reregistration of your product under FIFRA section 3(c)(5) when the agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

- Add the phrase "EPA Registration No. 9198-108."
- Under Storage and Disposal, revise the pesticide disposal statement to include "Do not contaminate water when disposing of equipment washwaters."

3. Submit one (1) copy of your final printed labeling before you release the product for shipment.

**BEST AVAILABLE COPY**

☐ ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 5(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Richard P. Mountfort  
Product Manager (23)  
Fungicide-Herbicide Branch  
Registration Division (TS-767C)

Enclosure

Front Label

**The Andersons Tee-Time Lawn Feed With Crabgrass Control<sup>R</sup>**

(LOGO)

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established: Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysiagrass.

**Active Ingredients:**

N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-p-toluidine .....	.67%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine .....	.33%
Inert Ingredients .....	99.00%

TEAM<sup>R</sup> ---- the trademark for Elanco Products benefin plus trifluralin

**Keep Out of Reach of Children**

**Warning**

**See Back Panel For Additional Precautionary Statements**

Net Weight 23 lbs.

Treats Up To 5,000 Sq.Ft.

EPA REG. NO. 9198-

EPA Est. 9198-OH-1

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

AUG 2 1989

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

9198-108

**BEST AVAILABLE COPY**

Back Panel

Net Weight 23 lbs.

Treats up to 5,000 Sq.Ft.

**The Andersens Tee-Time Lawn Food With Crabgrass Control<sup>R</sup>**

**Guaranteed Analysis**

Total Nitrogen (N) .....	25%
1.2% Ammoniacal Nitrogen	
17.3% Urea Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
Available Phosphoric Acid ( $P_2O_5$ ) .....	3%
Soluble Potash ( $K_2O$ ) .....	8%
Sulfur .....	4.8%

Plant Nutrients Derived From: Ammoniated Phosphates, Urea, Muriate of Potash and Sulfur Coated Urea.

**Keep Out Of Reach Of Children**

**Precautionary Statements**

**Warning**

**Hazards To Humans And Domestic Animals**

**Warnings: DO NOT GET IN EYES. CAUSES EYE IRRITATION.** If in eyes, flush with plenty of water. Harmful if swallowed, inhaled or absorbed through the skin. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**Statement Of Practical Treatment**

**IF IN EYES:** Hold eyelids open and flush with a steady, gently stream of water for 15 minutes. Call a physician.

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching finger to back of throat. Never give anything by mouth to an unconscious or convulsing person.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF INHALED:** If breathing difficulty occurs, move individual to fresh air and get medical attention.

**BEST AVAILABLE COPY**

### Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, or marshes). Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

### Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product will control annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyard-grass (watergrass), and green and yellow foxtail. Tee-Time Plus Team<sup>R</sup> controls these annual weeds by killing their seeds as they germinate.

When this product is applied as directed, the following established turfgrasses are tolerant:

Kentucky bluegrass  
Perennial ryegrass  
Centipedegrass

Fescue  
Zoysiagrass  
Bermudagrass

St. Augustinegrass  
Bahia grass

This product will not harm most nearby established ornamental trees and shrubs.

### Application Directions

#### Rates, Frequency and Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.

**BEST AVAILABLE COPY**

- 6-7
1. Spring germinating summer annual grasses - crabgrass, foxtails and barnyardgrass.

Northern (cool season) turfgrasses

For preemergence control, apply 4.6 pounds per thousand square feet of this product (2.0 pounds active ingredient per acre) in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

2. Late spring, early summer germinating annual grasses - goosegrass.

Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications of this product should, therefore, be timed accordingly.

Northern (cool season) turfgrass

For preemergence control, apply 4.6 pounds per thousand square feet of this product (2.0 pounds active ingredient per acre) in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

3. Northern and Southern turfgrasses

In areas of the United States where preemergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications of this product spaced eight to ten weeks apart should be made to provide full season control of these annual grass species.

4. *Poa annua* (annual bluegrass)

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

Northern (cool season) turfgrasses

For preemergence control, apply 4.6 pounds per thousand square feet of this product (2.0 pounds active ingredient per acre) in the late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.

BEST AVAILABLE COPY

### Reseeding

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying this product. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using this product the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as Rogers Aero Seeder) should be employed. When using this product reseeding should be delayed 12 to 16 weeks after application.

### Application Equipment

Apply this product evenly over the turfgrass area without skips or misses. A more uniform application can be made with broadcast (rotary) type spreaders by spreading half the required amount over the area and then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly.

### Suggested Spreader Settings

(This section will be completed for final labeling)

### Use Precautions

Apply this product to the turf prior to the germination of weed grasses since this product will not control established weeds. This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of this product should not be used on putting greens. This product should not be used on dichondra. This product should not be applied to newly sprigged areas of bermudagrass, St. Augustine-grass or Centipede-grass until these turfs are well established.

### Storage and Disposal

**Storage:** Storage in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Do not contaminate water, food, or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.