



PM22 9198-111

Tacket 1049

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (M7505C) 401 M St., S.W. Washington, D.C. 20460	EPA Reg. Number: 9198-111	Date of Issuance: FEB 19 1993
		Term of Issuance: Conditional	
		Name of Pesticide Product: The Anderson's Tee Time with 1.0% Bayleton Fungicide	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)			
Name and Address of Registrant (include ZIP Code): The Andersons Lawn Fertilizer Division DBA/Free Flow Fertilizer P.O. Box 119 Maumee, OH 43537			
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.			
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none">1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.2. Make the following label changes listed below before you release the product for shipment: - Add the phrase, "EPA Reg. No. 9198-111".3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.			
Signature of Approving Official: 		Date:	

2019

page 2
EPA Reg. No. 9198-111

4. Submit within 9 months from the date of this Notice of Registration a dermal sensitization study conducted in accordance with Guideline Reference Number 81-6 with this product.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

PM22 9198-111

Jacket 3 of 9

ACCEPTED
with COMMENTS
In EPA Letter Dated:

FEB 10 1993

Under the Federal Insecticide
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9198-111

(FRONT PANEL)

THE ANDERSON'S TEE TIME WITH 1.0%

BAYLETON(R) FUNGICIDE

FOR CONTROL OF CERTAIN DISEASES ON TURF

ACTIVE INGREDIENT: 1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,
2,4-triazol-1-yl)-2-butanone 1.0%
INERT INGREDIENTS: 99.0%
TOTAL 100.0%

Bayleton(R) is a Registered Trademark of Bayer AG, Leverkusen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Net Wt. __ lbs.

Treats up to _____ sq. ft.

EPA Reg. No. 9198-

EPA Est. 9198-OH-1

(BACK PANEL)

THE ANDERSONS TEE TIME WITH 1.0%

BAYLETON(R) FUNGICIDE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Prolonged or frequent repeated skin contact may cause allergic skin reactions in some susceptible individuals. Keep out of reach of children.

Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed, administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately.

If on skin, wash skin immediately with soap and warm water. If irritation occurs, get medical attention.

If eyes are contaminated, wash with plenty of water for at least 15 minutes. Get medical attention.

If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEST AVAILABLE COPY

IMPORTANT: Read these entire Directions For Use before using The Andersons Tee Time With 1.0% Bayleton(R) Fungicide.

APPLICATION

BAYLETON(R) fungicide is absorbed rapidly and works systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Application should be applied at recommended intervals to maintain disease control.

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL:

DISEASE	Lbs. of Tee Time With 1.0% Bayleton(R) Fungicide Per 1,000 sq. ft.*		REMARKS
	Preventative	Curative	
Dollar Spot	1.5	-	PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of Bayleton fungicide may extend for as long as 60 days dependent environmental conditions.
	.75	-	PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of Bayleton fungicide may extend for as long as 30 days dependent upon environmental conditions.
	-	3	CURATIVE RATE: To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

Copper Spot
Powdery Mildew
Red Thread
(*Laetisaria*
fuciformis)

1.5

3

Rusts

Brown Patch
(*Rhizoctonia*
solani)
(Suppression)

PREVENTIVE RATE:

Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of Bayleton fungicide can be greater than 30 days depending on environmental conditions.

CURATIVE RATE:

To control existing infections, apply the curative rate. Re-applications should be not closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

Anthracnose

Poa Annua

3

-

Decline

PREVENTIVE RATE:

Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.

1.5

-

PREVENTIVE RATE:

Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days. ✓

-

3

CURATIVE RATE:

To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

Stripe Smut

Preventive Rates Only

3

Make a single application in spring as grass growth begins.

BEST AVAILABLE COPY

7049

Fusarium Blight (Fusarium culmorum) (Fusarium poae) Summer Patch (Phialophora graminicola) (Except California)	3	Apply first application in mid-June or 30 day prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.
Fusarium Blight (Fusarium culmorum) (Fusarium poae) Summer Patch (Phialophora graminicola) (California only)	3 to 6	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at the 6 lb. rate on 15 day intervals or at the 12 lb. rate on 30 day intervals.
Gray Snow Mold (Typhula incarnata) (Except California)	6	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
Pink Snow Mold Fusarium Patch (Fusarium nivale) Southern Blight (Sclerotium rolfsii) (Except California)	3 to 6	Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snow cover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.
Gray Snow Mold (Typhula incarnata) Pink Snow Mold (Fusarium nivale) (California only)	Single Application 7.5 to 12	Apply in late fall before first snow cover. Do not apply over snow cover. Use the high rate in areas with a history of severe snowmold damage.

Note*: Apply the specified amount of The Andersons Tee Time With 1.0% Bayleton(R) Fungicide with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft.

BEST AVAILABLE COPY

Make application when turf is dry and following the mowing operation. Do not mow until a minimum of .25 inch of water has been applied to treated area. Do not use clippings for animal feed.

SUGGESTED SPREADER SETTINGS

SPREADER	MODEL	SETTING
Cyclone	B1-84	4
Lely	W	5 1/2
Lesco		5 1/2
Scotts	R8	5 1/2
Vicon	402	33

NOTE: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area to be treated. Measure area to be treated. Do not over or under apply. Close spreader when filling, turning or stopping. DO NOT HAND SPREAD.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage and disposal. Store in cool, dry place. Prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container out of reach of children, preferably in a locked storage area.

DISPOSAL: Do not reuse empty container. Securely wrap in newspaper and discard in trash.

Pesticide Disposal (Except household): Wastes resulting from the use of this product may be disposed on site or at an approved waste disposal facility. ✓

Container Disposal (Except household): Do not use container in connection with food, feed, or drinking water. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

9 of 9

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by: The Andersons
Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537

✓