

9198-111

9/23/2002

4/7

SEP 23 2002

Ms. Dawn L. Walters
The Andersons, Inc.
P.O. Box 119
Maumee, Ohio 43537

Subject: The Andersons 1.0% Bayleton Fungicide
EPA Reg. No. 9198-111
Your Submission of June 12, 2002

Dear Ms. Walters:

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that the following labeling change is made prior to release of the product for shipment.

In the ingredients statement on the front panel of both labels, the common name of the active ingredient is misspelled. Correct to read "Triadimefon".

Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed.

Sincerely yours,



Cynthia Giles-Parker
Product Manager (22)
Fungicide Branch
Registration Division (7505C)

Enclosure

7505C:CGable:cg:9/20/02

2/7

(FRONT PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

THE ANDERSONS 1.0% BAYLETON® FUNGICIDE

FOR CONTROL OF CERTAIN DISEASES ON TURF

ACTIVE INGREDIENT:

Traiadimefon* 1.0%

OTHER INGREDIENTS 99.0%

TOTAL 100.0%

*CAS No. 43121-43-3

Bayleton® is a Registered Trademark of Bayer AG

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WT. _____ LBS.

TREATS UP TO _____ SQ. FT.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

SEP 23 2002

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

9198-111

EPA Reg. No. 9198-111

EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code on bag

(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

THE ANDERSONS 1.0% BAYLETON® FUNGICIDE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

FIRST AID

- | | |
|--------------|---|
| If swallowed | Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If on skin | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. |
| If inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice. |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use on only on home lawns.

4/7

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

APPLICATION

BAYLETON® fungicide is absorbed rapidly and works systematically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Application should be applied at recommended intervals to maintain disease control.

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL:

DISEASE	Lbs. of The Andersons 1.0% Bayleton® Fungicide Per 1,000 sq. ft.*		REMARKS
	PREVENTATIVE	CURATIVE	
Dollar Spot	1.5	-	PREVENTIVE RATE: Apply recommended rate on 30-day intervals. Protective activity of Bayleton fungicide may extend for as long as 60 days dependent upon environmental conditions
	0.75	-	PREVENTATIVE RATE: Apply recommended rate on 14-day intervals. Protective activity of Bayleton fungicide may extend for as long as 30 day dependent on environmental conditions.
	-	3	CURATIVE RATE: To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventative rate schedules for rates and timing.

(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Copper Spot Powdery Mildew Red Thread (<i>Laetisaria fuciformis</i>) Rusts Brown Patch (<i>Rhizoctonia solani</i>) (Suppression)	1.5	-	PREVENTATIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light-to-moderate disease development, use a longer interval. Protective activity of Bayleton fungicide can be greater than 30 days depending on environmental conditions.
	-	3	CURATIVE RATE: To control existing infections, apply the curative rate. Reapplications should be no closer than 14 days. When infections have been controlled, refer to the preventative rate schedule for rates and timing.
Anthracnose <i>Poa annua</i> decline	3	-	PREVENTATIVE RATE: Apply at 30-day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
	1.5	-	PREVENTATIVE RATE: Apply at 21-day intervals and repeat as necessary for seasonal conditions, residual control may be extended to 30 days.
	-	3.0	CURATIVE RATE: To control existing infections, apply the curative rate. Re-applications should be no closer than 14 days. When infections have been controlled, refer to the preventative rate schedule for rates and timing.
PREVENTIVE RATES ONLY			
Stripe Smut	3		Make a single application in spring as grass growth begins.
Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>) Summer Patch (<i>Phialophora graminicola</i>) (Except California)	3		Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at 30-day intervals.
Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>) Summer Patch (<i>Phialophora graminicola</i>) (California only)	3 to 6		Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at the 6 lb. rate on 15-day intervals.

(BACK PANEL)
HOMEOWNER USE ONLY*(The above is for internal use only and will not appear on the final label copy)*

Gray Snow Mold (Typhula incarnata) (Except California)	6	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover or when turf is dormant.
Pink Snow Mold Fusarium Patch (Fusarium nivale) Southern Blight (Sclerotium rolfsii) (Except California)	3 to 6	Apply before conditions favorable for infection occur. Reapplication should be made as needed at 60 to 90-day interval. Do not apply over snow cover or when turf is dormant. Use higher rate in areas with a history of severe disease damage.
Gray Snow Mold (Typhula incarnata) Pink Snow Mold (Fusarium nivale) (California Only)	Single Application 7.5 to 12	Apply in late fall before first snow cover. Do not apply over snow cover. Use the high rate in areas with a history of severe snow mold damage.

NOTE*: Apply the specified amount of The Andersons Tee Time With 1.0% Bayleton(R) Fungicide with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft.

Make application when turf is dry and following the mowing operation. Do not mow until a minimum of 0.25 inch of water has been applied to treated area. Do not use clippings for animal feed.

SUGGESTED SPREADER SETTINGS

(Note to EPA: spreader types and models will vary dependent on current market demand and availability. Actual spreaders and settings will be on final printed label)

SPREADER

MODEL

SETTING

Important: It is important to follow the directions as indicated on this quality lawn care product as overapplication may cause injury to your lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. Do not hand spread.

Broadcast Spreaders: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage. *{diagram of spread pattern}*

Drop Spreaders: First apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips. *{diagram of spread pattern}*

7/7

(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies.

DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (AP21993/RVamd61102)

Manufactured By:

The Andersons Lawn Fertilizer Division, Inc.
d/b/a Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537

For more information, contact your local retailer or call 1-800-4-A-Andersons. For more information, contact your local retailer or call 1-800-4-A-Andersons.

