

PM 22

9198-112

7/12/98

(approved stamp on p. 7)

1/7

9198-112H  
Page 1 of 7

(FRONT PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

## THE ANDERSONS TEE TIME WITH 0.5% BAYLETON® FUNGICIDE

FOR CONTROL OF CERTAIN DISEASES ON TURF

**ACTIVE INGREDIENT:**

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone ..... 0.5%

INERT INGREDIENTS ..... 99.5%

TOTAL 100.0%

Bayleton(R) is a Registered Trademark of Bayer AG, Leverkusen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Net Wt. \_\_\_ lbs.

Treats up to \_\_\_\_\_ sq. ft.

EPA Reg. No. 9198-112

EPA Est. 9198-OH-1

(BACK PANEL)

HOMEOWNER USE ONLY

*(The above is for internal use only and will not appear on the final label copy)*

**THE ANDERSONS TEE TIME WITH 0.5% BAYLETON(R) FUNGICIDE**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children.

Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

**STATEMENTS OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately.

**IF ON SKIN:** Wash skin immediately with soap and warm water. If irritation occurs, get medical attention.

**IF EYES ARE CONTAMINATED:** Wash with plenty of water for at least 15 minutes. Get medical attention.

**IF INHALED:** Remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

**NOTE TO PHYSICIAN:** No specific antidote is available. Treat symptomatically.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water by disposing of equipment washings. Apply this product only as specified on this label. Do not make applications when weather conditions favor drift from target area.

**Ground Water Advisory**

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

(BACK PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**For use only on home lawns.**

**Do not use on crops grown for food or forage.**

APPLICATION

BAYLETON® fungicide is absorbed rapidly and works systematically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Application should be applied at recommended intervals to maintain disease control.

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL:

DISEASE	Lbs. of Product Per 1,000 sq. ft.*		REMARKS
	<u>Preventive</u>	<u>Curative</u>	
Dollar Spot	3	-	<u>PREVENTIVE RATE:</u> Apply recommended rate on 30 day intervals. Protective activity of Bayleton fungicide may extend for as long as 60 days dependent environmental conditions.
	1.5	-	<u>PREVENTIVE RATE:</u> Apply recommended rate on 14 day intervals. Protective activity of Bayleton fungicide may extend for as long as 30 days dependent upon environmental conditions.
	-	6	<u>CURATIVE RATE:</u> To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

4/7

(BACK PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Copper Spot  
Powdery Mildew  
Red Thread  
(Laetisaria  
fuciformis)

3

6

Rusts

Brown Patch  
(Rhizoctonia  
solani)  
(Suppression)

PREVENTIVE RATE:

Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of Bayleton fungicide can be greater than 30 days depending on environmental conditions.

CURATIVE RATE:

To control existing infections, apply the curative rate. Re-applications should be not closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

Anthracnose  
Poa Annua  
Decline

Preventive

6

Curative

-

PREVENTIVE RATE:

Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.

PREVENTIVE RATE:

Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.

3

-

CURATIVE RATE:

To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

-

6

Stripe Smut

Preventive Rates Only

6

Make a single application in spring as grass growth begins.

(BACK PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Fusarium Blight ( <u>Fusarium</u> <u>culmorum</u> ) ( <u>Fusarium poae</u> ) Summer Patch ( <u>Phialophora</u> <u>graminicola</u> )  (Except California)	6	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident.  Applications should be made on 30 day intervals.
Fusarium Blight ( <u>Fusarium</u> <u>culmorum</u> ) ( <u>Fusarium poae</u> ) Summer Patch ( <u>Phialophora</u> <u>graminicola</u> )  (California only)	6 to 12	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident.  Applications should be made at the 6 lb. rate on 15 day intervals or at the 12 lb. rate on 30 day intervals.
Gray Snow Mold ( <u>Typhula</u> <u>incarnata</u> )  (Except California)	12	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
Pink Snow Mold Fusarium Patch ( <u>Fusarium</u> <u>nivale</u> ) Southern Blight ( <u>Sclerotium</u> <u>rolfsii</u> )  (Except California)	6 to 12	Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snow cover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.
Gray Snow Mold ( <u>Typhula</u> <u>incarnata</u> ) Pink Snow Mold ( <u>Fusarium</u> <u>nivale</u> ) (California Only)	Single Application  15 to 24	Apply in late fall before first snow cover. Do not apply over snow cover. Use the high rate in areas with a history of severe snowmold damage.

6/7

(BACK PANEL)

HOMEOWNER USE ONLY

*(The above is for internal use only and will not appear on the final label copy)*

Note\*: Apply the specified amount of The Andersons Tee Time With 0.5% Bayleton® Fungicide with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of .25 inch of water has been applied to treated area. Do not use clippings for animal feed.

SUGGESTED SPREADER SETTINGS

<u>SPREADER</u>	<u>MODEL</u>	<u>SETTING</u>
Cyclone	B1-84	4
Lely	W	5 1/2
Lesco		5 1/2
Scotts	R8	H 1/2
Vicon	402	33

NOTE: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area to be treated. Measure area to be treated. Do not over or under apply. Close spreader when filling, turning or stopping. DO NOT HAND SPREAD.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

(BACK PANEL)

HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)


MANUFACTURERS STATEMENT

This product conforms to the ingredient statement on the label. Manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage and handling of this product.

THIS PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED. IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND.

Manufactured by: The Andersons Lawn Fertilizer Division  
d/b/a Free Flow Fertilizer  
P.O. Box 119  
Maumee, OH 43537

Date Printed: 07/14/97  
Date Approved:  
Supersedes:  
Label Change: wps exclusionary/ground water advisory/mfg statement  
Filename: 9198112h.doc



PM 22

9198-112

1/12/98

1/7

9198-112C  
Page 1 of 7

(FRONT PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

## THE ANDERSONS TEE TIME WITH 0.5% BAYLETON(R) FUNGICIDE

FOR CONTROL OF CERTAIN DISEASES ON TURF

**ACTIVE INGREDIENT:**

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone ..... 0.5%

**INERT INGREDIENTS:** ..... 99.5%

TOTAL 100.0%

Bayleton(R) is a Registered Trademark of Bayer AG, Leverkusen.


KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Net Wt. \_\_\_ lbs.

Treats up to \_\_\_\_\_ sq. ft.



EPA Reg. No. 9198-112

EPA Est. 9198-OH-1



2/17

(BACK PANEL)

**COMMERCIAL USE ONLY**

*(The above is for internal use only and will not appear on the final label copy)*

**THE ANDERSONS TEE TIME WITH 0.5% BAYLETON(R) FUNGICIDE**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children.

Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

**STATEMENTS OF PRACTICAL TREATMENT**

IF SWALLOWED, administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately.

IF ON SKIN, wash skin immediately with soap and warm water. If irritation occurs, get medical attention.

IF EYES ARE CONTAMINATED, wash with plenty of water for at least 15 minutes. Get medical attention.

IF INHALED, remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washings. Apply this product only as specified on this label. Do not make applications when weather conditions favor drift from target area.

3/17

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Ground Water Advisory

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not use on crops grown for food or forage.

APPLICATION

BAYLETON(R) fungicide is absorbed rapidly and works systematically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Application should be applied at recommended intervals to maintain disease control.

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL:

DISEASE	Lbs. of Product Per 1,000 sq. ft.*		REMARKS
	<u>Preventative</u>	<u>Curative</u>	
Dollar Spot	3	-	<u>PREVENTIVE RATE:</u> Apply recommended rate on 30 day intervals. Protective activity of Bayleton fungicide may extend for as long as 60 days dependent environmental conditions.

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

1.5

PREVENTIVE RATE:

Apply recommended rate on 14 day intervals. Protective activity of Bayleton fungicide may extend for as long as 30 days dependent upon environmental conditions.

CURATIVE RATE:

To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

Copper Spot . . . Preventive

Powdery Mildew

Red Thread

(Laetisaria  
fuciformis)

3

Curative

6

Rusts

Brown Patch

(Rhizoctonia  
solani)

(Suppression)

PREVENTIVE RATE:

Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of Bayleton fungicide can be greater than 30 days depending on environmental conditions.

CURATIVE RATE:

To control existing infections, apply the curative rate. Re-applications should be not closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.

Anthrachnose

Poa Annua

Decline

6

PREVENTIVE RATE:

Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45

days.

5/7

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

		PREVENTIVE RATE:
3	-	Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
		CURATIVE RATE:
-	6	To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.
Stripe Smut	<u>Preventive Rates Only</u> 6	Make a single application in spring as grass growth begins.
Fusarium Blight ( <u>Fusarium culmorum</u> ) ( <u>Fusarium poae</u> )	6	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident.
Summer Patch ( <u>Phialophora graminicola</u> )  ( <u>Except California</u> )		Applications should be made on 30 day intervals.
Fusarium Blight ( <u>Fusarium culmorum</u> ) ( <u>Fusarium poae</u> )	6 to 12	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident.
Summer Patch ( <u>Phialophora graminicola</u> )  ( <u>California only</u> )		Applications should be made at the 6 lb. rate on 15 day intervals or at the 12 lb. rate on 30 day intervals.

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Gray Snow Mold ( <u>Typhula</u> <u>incarnata</u> )	12	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
( <u>Except</u> California)		
Pink Snow Mold Fusarium Patch ( <u>Fusarium</u> <u>nivale</u> )	6 to 12	Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snow cover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.
Southern Blight ( <u>Sclerotium</u> <u>rolfsii</u> )		
( <u>Except</u> California)		
Gray Snow Mold ( <u>Typhula</u> <u>incarnata</u> )	Single Application	Apply in late fall before first snow cover. Do not apply over snow cover. Use the high rate in areas with a history of severe snowmold damage.
Pink Snow Mold ( <u>Fusarium</u> <u>nivale</u> )	15 to 24	
(California Only)		

Note\*: Apply the specified amount of The Andersons Tee Time With 0.5% Bayleton(R) Fungicide with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of .25 inch of water has been applied to treated area. Do not use clippings for animal feed.

SUGGESTED SPREADER SETTINGS

<u>SPREADER</u>	<u>MODEL</u>	<u>SETTING</u>
Cyclone	B1-84	4
Lely	W	5 1/2
Lesco		5 1/2
Scotts	R8	H 1/2
Vicon	402	33

NOTE: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area to be treated. Measure area to be treated. Do not over or under apply. Close spreader when filling, turning or stopping. DO NOT HAND SPREAD.

(BACK PANEL)

COMMERCIAL USE ONLY

*(The above is for internal use only and will not appear on the final label copy)*

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

MANUFACTURERS STATEMENT

This product conforms to the ingredient statement on the label. Manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage and handling of this product.

THIS PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED. IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND.

Manufactured by: The Andersons  
Lawn Fertilizer Division  
P.O. Box 119  
Maumee, OH 43537

Date Printed: 07/11/97  
Date Approved:  
Supersedes:  
Label Change: wps exclusionary/mfg statement/ground water advisory  
Filename: 9198112c.doc