		9198	-112		женициот»	,	130/		
SEPA	United States  Environmental Protection Age Washington, DC 20460			ıcy	Form App	R	MB No. 2 egistra mendn ther	ion	OPP Identifier Number
	A	pplication	for Pe	esticid	e - Sect	ion I			· <u>}</u>
1. Company/Product Numb 9198-112	par				roduct Mane arker M		<b>ι</b> (∢√	·	posed Classification
4. Company/Product (Nam The Andersons 0.5%				PMy 38 21					
5. Name and Address of A The Andersons Law PO Box 119 Maumee, OH 4353	n Fert. Div. d/b/a Fr	-	rt.	(b)(i), my to: EPA Re		s similar	or identi	cal in cor	FIFRA Section 3(c)(3) mposition and labeling
	<del></del>	<del></del>		on - ii	I Ivaille				
Notification - Explain Statement of Practical Treats and spreader directions. All This notification is consistent confidentioal statement of for	ponse to Agency letter d in below.  In below.  In a page(s) if necessary.  In ange per PR Notice 98-10:  In ent' header to "First Aid". Also  changes are in bold and highlique  with the provisions of PR Notice  mula of this product. I underthe  with the terms of PR Notice 98-	(For section I prand name chang to added all Ander the ghed in yellow.  e 98-10 and EPA ad that It is a viola	ge, revised from estab regualtions tion of 18 U 52.46, this p	tion II.) Warranty silishment nu	*Me Too* A Other - Expli- tatement, replay umbers to from 152.46 and no 1001 to wilfully y be in violation	ain belov ace "Inert" t panel. O	with "Other" n homeowne	MA in ilngrediel r version, a een made t	lso added optional marketing to the labeling or the
1. Material This Product W	Fil Re Packaged In:		Secur	JII - III					
Child-Resistant Packaging Yes No Pertification must Submitted  3. Location of Net Contents	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per container . Size(s) Retail	N N If "Yes" Package		No, per container	5. Locati	Type of C	Metal Plastic Glass Paper Other (S	
6. Manner in Which Label i		Lithograp Paper gid Stenciled	h ed		Other				
	<del></del>	Staticited		n - IV		. <u></u>			
1. Contact Point (Complet	e items directly below for	identification (	of individu	uel to be	contacted, i	f necess	ery, to pro	cess this	epplication.)

I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. 3. Title

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete.

6. Date Application Received

Telephone No. (Include Area Code)

334-262-2766

(Stamped)

2. Signature

Regulatory Administrator Dawn & Walters

5. Date

4. Typed Name Dawn L. Walters

Name

Dawn L. Walters

5/15/00

**Regulatory Administrator** 

9198-112H Page 1 of 6

#### (FRONT PANEL) HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

# THE ANDERSONS 0.5% BAYLETON® FUNGICIDE

# Optional Marketing Claims:

- FOR CONTROL OF CERTAIN DISEASES ON TURF
- · Prevents most common turf diseases that damage or destroy bluegrass, ryegrass, fescue and bentgrass lawns
- Controls most existing fungus infections when used at the curative rate
- {Product Name} is absorbed rapidly and systemically by the roots of the plants for control of disease
- · For the best lawn in town
- Ready-to-use granules for systemic control of lawn diseases
- · Controls: dollar spot, brown patch, powdery mildew, red thread and many others
- · Controls certain turf diseases including Dollar Spot, Strip Smut, and Brown Patch
- Contains Bayleton for systemic control

ACTIVE INGREDIENT:	
1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone	0.5%
OTHER INGREDIENTS	<u>99.5%</u>
	TOTAL 100.0%
Bayleton(R) is a Registered Trademark of Bayer AG, Leverkusen.	
KEEP OUT OF REACH OF CHILDREN CAUTION SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS	MOTERATION
	NOTIFICATION
Net Wt lbs. Treats up to sq. ft.	MAY 3 0 2000
Possible treatment coverage: Treats up to 3,000 sq. ft.	

EPA Reg. No. 9198-112
EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code on bag

#### (BACK PANEL) HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

# THE ANDERSONS 0.5% BAYLETON(R) FUNGICIDE

#### PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

# **CAUTION**

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children.

Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

## **FIRST AID**

IF SWALLOWED: Administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately.

IF ON SKIN: Wash skin immediately with soap and warm water. If irritation occurs, get medical attention.

IF EYES ARE CONTAMINATED: Wash with plenty of water for at least 15 minutes. Get medical attention.

IF INHALED: Remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

#### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water by disposing of equipment washings. Apply this product only as specified on this label. Do not make applications when weather conditions favor drift from target area.

Ground Water Advisory

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

# (BACK PANEL) HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For use only on home lawns . Do not use on crops grown for food or forage.

## **APPLICATION**

BAYLETON® fungicide is absorbed rapidly and works systematically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Application should be applied at recommended intervals to maintain disease control.

# RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL:

Disease	Lbs. of Product Per 1,000 sq.ft.		Remarks		
	Preventive	Curative			
Dollar Spot	3	-	Preventative Rate: Apply recommended rate on 30 day intervals. Protective activity of Bayleton fungicide may extend for as long as 60 days of dependent environmental conditions.		
	1.5	-	Preventative Rate: Apply recommended rate on 14 day intervals. Protective activity of Bayleton fungicide may extend for as long as 30 days dependent upon environmental conditions.		
		6	Curative Rate: To control existing infections, apply the curative rate. Reapplication should be no closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.		
Copper Spot Powder Mildew Red Thread (Laetisara fuciformis) Rusts Brown Patch (Rhizoctonia solani) (suppression)	3	6	Preventive Rate: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of Bayleton fungicide can be greater than 30 days depending on environmental conditions.  Curative Rate: To control existing infections, apply the curative rate. Reapplications should not be closer than 14 days. When infections have been controlled, refer to the preventive rate schedule for rates and timing.		

9198-112H Page 4 of 6

## (BACK PANEL) HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Anthracnose	6		Preventive Rate: Apply at 30 day intervals and
Poa Annua Decline		}	repeat as necessary for seasonal control.
t out thinds boom to			Dependent upon environmental conditions,
			residual control may be extended to 45 days.
	3	<del> </del>	Preventive Rate: Apply at 21 day intervals and
	J		repeat as necessary for seasonal control.
Í		1	Dependent upon environmental conditions,
<i>'</i>		!	residual control may be extended to 30 days.
		6	Curative Rate: To control existing infections,
1	-	0	
}		ļ	apply the curative rate. Reapplication should be no closer than 14 days. When infections
			have been controlled, refer to the preventive
	Drovent	ius Patas Only	rate schedule for rates and timing.
Stain Smout	Prevent	ive Rates Only	Make a significant in the signif
Strip Smut			Make a single application in spring as grass growth begins.
Fusarium Blight		6	Apply first application in mid-June or 30 days
(Fusarium culmorum)			prior to time this blight normally becomes
(Fusarium poae)			evident.
Summer patch			
(Phialophora			Applications should be made on 30 day
graminicola)			intervals.
(Except California)		· · · · · · · · · · · · · · · · · · ·	
Fusarium Blight		6 to 12	Apply first application in mid-June of 30 days
(Fusarium culmorum)			prior to time this blight normally becomes
(Fusarium poae)			evident.
Summer patch			
(Phialophora			Applications should be made at the 6 lb. rate
graminicola)			on 15 day intervals or at the 12 lb. rate on 30
			day intervals.
(California only)			
Gray Snow Mold	12		Apply in fall 30 days prior to turf dormancy. If
(Typhula incarnata)			turf breaks dormancy during winter months a
			second application should be made. Do not
(Except California)			apply over snow cover, or when turf is
			dormant.
Pink Snow Mold	•	6 to 12	Apply before conditions favorable for infection
Fusarium Patch			occur. Reapplication should be made as
(Fusarium nivale)			needed at a 60 to 90 day interval. Do no apply
Southern Blight			over snow cover, or when turf is dormant. Use
(Sclerotium rolfsii)			higher rate in areas with a history of severe
İ			disease damage.
(Except California)			
Gray Snow Mold	Single	Application	Apply in late fall before first snow cover. Do
(Typhula incarnata)	_	• •	not apply over snow cover. Use the high rate
Pink Snow Mold	1	5 to 24	in areas with a history of severe snowmold
(Fusarium nivale)			damage.
(California only)			

Note\*: Apply the specified amount of The Andersons Tee Time With 0.5% Bayleton® Fungicide with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of .25 inch of water has been applied to treated area. Do not use clippings for animal feed.

9198-112H Page 5 of 6

#### (BACK PANEL) HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

## SUGGESTED SPREADER SETTINGS

SPREADER	MODEL	SETTING
Cyclone Lely Lesco	B1-84 W	4 5 1/2 5 1/2
Scotts	R8	H 1/2
Vicon	402	33

NOTE: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area to be treated. Measure area to be treated. Do not over or under apply. Close spreader when filling, turning or stopping. DO NOT HAND SPREAD.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

## WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted. (AP032891/RVnot051500)

## Optional spreader directions:

It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Calibrate your spreader on 100 square feet and adjust as necessary. Refer to the table and picture below to determine the size of your lawn.

Close spreader hopper and adjust the spreader setting using the suggested spreader settings chart below. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn. Refer to "Path A" when using a broadcast spreader and "Path B" when using a drop spreader. Close the spreader hopper whenever spreader stops or turns.

OR

Coverage Hints Directions:

#### (BACK PANEL) HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

1. Close spreader hopper

2. Adjust the spreader setting using the spreader suggestions chart

3. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn

4. Apply product to lawn. Refer to Path A when using a broadcast spreader and Path B when using a drop spreader. Close the spreader hopper whenever spreader stops or turns.

5. If using a broadcast spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

This bag should cover up to 6,000 sq. ft. (100 ft x 60 ft) at the 1.5 lbs. per 1,000 sq. ft. rate. Refer to the table and picture to determine the size of your lawn.

{diagrams of lawn, spreader paths and calculation chart}

Optional (subregistrant) marketing information:

Frank's Quality Guarantee: Frank's takes pride in offering only the highest quality. Every Frank's product is formulated by our experts to help you enjoy a beautiful lawn. You'll feel confident that you made the 'quality' choice when you choose Frank's.

Manufactured by: The Andersons Lawn Fertilizer Division

d/b/a Free Flow Fertilizer

P.O. Box 119

Maumee, OH 43537

Date Printed:	5/15/00 (notification)
Date Approved:	01/12/98
Supersedes:	3/28/91
Label change:	Notification: name change/ other ingredient/first aid heading/warranty statement/optional marketing & spreader directions
Filename:	9198112H
Label code:	AP032891/RVnot051500