· 9198-112 5-17-2004

≎EPA	Environmenta Washi	Inited States I Protecti ngton, DC 20	460	,	V	Registra Amenda Other	ation	OPP Identifier Number
Application for Pesticide - Section I								
1. Company/Product Number	r			PA Product Ma	nager		3. Pr	oposed Classification
9198-112			Gil	es-Parker			V	None Restricted
4. Company/Product (Name) The Andersons 0.5% Baylete				PM# 22				
5. Name and Address of Ap The Andersons Lawn F d/b/a Free Flow Fertilize PO Box 119 Maumee OH 43537 Check if this	6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name							
			Section	-]]				
Amendment - Explain Resubmission in resp	onse to Agency letter	dated	[Final printo Agency let "Me Too" Other - Ex	tter dat Applic	ation.		NOTIFICATION MAY 1 7 2004
Notification of Minor Label changes per PR Notice 98-10: addition of California Prop 65 language; see cover letter for details This notification is consistent with the provisions of PR Notices 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.								
			Section	<u>- III</u>				
1. Material This Product Will	Be Packaged In:							
Child-Resistant Packaging	Unit Packaging					2. Type of Container Metal		
Yes	Yes		Yes					
No	No		No			Plastic Glass		
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wg	Wgt container Paper Other (Specify)			ipecify)	
3. Location of Net Contents	nformation	4. Size(s) Ret	tail Container		5. Lo	cation of Lab	el Directio	ាន
Label C	ontainer				l	On Labe	1	
6. Manner in Which Label is	Affixed to Product	Lithog Paper Stenc	reph glued led	Othe	or			
			Section	- IV		<u> </u>		
1. Contact Point (Complete	items directly helow for	or identificatio			if nec	essery, to pro	ocass this	englication I
Name Dawn L. Walters			1 '				No. (include Area Code)	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					Received			
2. Signature			3. Title				● # *	
Dawn Klineiers			Regulatory Administrator					*
4. Typed Name Dawn L. Walters			5. Date	5/3/04			-	,, 3 8

9198-112

(FRONT PANEL)

THE ANDERSONS 0.5% BAYLETON® FUNGICIDE

Optional Marketing Claims:

- FOR CONTROL OF CERTAIN DISEASES ON TURF
- Prevents most common turf diseases that damage or destroy bluegrass, ryegrass, fescue and bentgrass lawns
- Controls most existing fungus infections when used at the curative rate
- {Product Name} is absorbed rapidly and systemically by the roots of the plants for control of disease
- For the best lawn in town
- Ready-to-use granules for systemic control of lawn diseases
- Controls: dollar spot, brown patch, powdery mildew, red thread and many others
- Controls certain turf diseases including Dollar Spot, Stripe Smut, and Brown Patch
- Contains Bayleton for systemic control

ACTIVE INGREDIENT: TriadimefonF OTHER INGREDIENTS			
TCAS No. 43121-43-3	TOTAL 100.0%		
Bayleton® is a Registered Trademark of Bayer C	Crop Science		
KEEP OUT OF REACH OF CHILDREN CAUTION SEE BACK PANEL FOR ADDITIONAL PRECA	UTIONS		
Net Wt lbs. Treats up to sq. ft.			
Possible treatment coverage: Treats up to 3,000 sq. ft.			
EPA Reg. No. 9198-112 EPA Est. 9198-OH-1 <u>M,</u> 9198-OH-2 <u>B,</u> 9198-AL-0 Underlined letter is first letter used in run cod			; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;
`-	NOTIFICATION MAY 1 7 2004	*****	6

THE ANDERSONS 0.5% BAYLETON® FUNGICIDE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **CAUTION**

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

FIRST AID

Call a poison control center or doctor for treatment advice. Have person sip a If swallowed

> glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-

20 minutes. Call a poison control center or doctor for treatment advice.

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove If in eyes

contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison

control center or doctor for treatment advice.

if inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then

give artificial respiration, preferably mouth-to mouth if possible. Call a poison

control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

ENVIRONMENTAL HAZARDS

For homeowner Products:

Do not apply directly to water. Do not contaminate water by disposing of equipment washings. Apply this product only as specified on this label. Do not make applications when weather conditions favor drift from target area.

For commercial products:

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washings. Apply this product only as specified on this label. Do not make applications when weather conditions favor drift from target area.

Ground Water Advisory

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not use on crops grown for food or forage.

APPLICATION

BAYLETON® fungicide is absorbed rapidly and works systematically from within the plant. Rainfall or sprinkler imigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Application should be applied at recommended intervals to maintain disease control.

APPLICATION RATES FOR TURF GRASS DISEASE CONTROL:

		Product	Remarks		
Disease		00 sq.ft.			
	Preventative	Curative			
Dollar Spot	3	-	Preventative Rate: Apply recommended rate on 30 day intervals. Protective activity of Bayleton fungicide may extend for as long as 60 days of dependent environmental conditions.		
	1.5	.*	Preventative Rate: Apply recommended rate on 14 day intervals. Protective activity of Bayleton fungicide may extend for as long as 30 days dependent upon environmental conditions.		
	-	6	Curative Rate: To control existing infections, apply the curative rate. Reapplication should be no closer than 14 days. When infections have been controlled, refer to the Preventative rate schedule for rates and timing.		
Copper Spot Powder Mildew Red Thread (Laetisara fuciformis) Rusts Brown Patch (Rhizoctonia solani) (suppression)	3	6	Preventative Rate: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of Bayleton fungicide can be greater than 30 days depending on environmental conditions. Curative Rate: To control existing infections, apply the curative rate. Reapplications should not be closer than 14 days. When infections have been controlled, refer to the Preventative rate schedule for rates and timing.		
Anthracnose Poa Annua Decline	6	-	Preventative Rate: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.		
	3	•	Preventative Rate: Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmenta conditions, residual control may be extended to 30 days.		
	-	6	Curative Rate: To control existing infections, apply the curative rate. Reapplication should be no closer than 14° days. When infections have been controlled, refer to the Preventative rate schedule for rates and timing.		

Disease	Lbs. of Product Per 1,000 sq.ft. Preventative Rates Only	Remarks		
Stripe Smut	6	Make a single application in spring as grass growth begins.		
Fusarium Blight (Fusarium culmorum) (Fusarium poae) Summer patch (Phialophora graminicola)	6	Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.		
(Except California)				
Fusarium Blight (Fusarium culmorum) (Fusarium poae) Summer patch	6 to 12	Apply first application in mid-June of 30 days prior to time this blight normally becomes evident.		
(Phialophora graminicola) (California only)		Applications should be made at the 6 lb. rate on 15 day intervals or at the 12 lb. rate on 30 day intervals.		
Gray Snow Mold (Typhula incarnata) (Except California)	12	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.		
Pink Snow Mold Fusarium Patch (Fusarium nivale) Southern Blight (Sclerotium rolfsii) (Except California)	6 to 12	Apply before conditions favorable for infection occur. Reapplication should be made as needed at a 60 to 90 day interval. Do no apply over snow cover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.		
Gray Snow Mold (Typhula incarnata) Pink Snow Mold (Fusarium nivale)	Single Application 15 to 24	Apply in late fall before first snow cover. Do not apply over snow cover. Use the high rate in areas with a history of severe snowmold damage.		

Note*: Apply the specified amount of The Andersons 0.5% Bayleton® Fungicide with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of 0.25 inch of water has been applied to treated area. Do not use clippings for animal feed.

SUGGESTED SPREADER SETTINGS (available on final printed label)

Directions for homeowner products:

It is important to follow the directions as indicated so that injury will not occur to your lawn.

Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to check spreader at the beginning of the application; adjust spreader if necessary. Close spreader when filling, turning or stopping.

DO NOT HAND SPREAD.

Broadcast Spreaders: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of wheels will ensure complete coverage. (directional diagram)

Drop Spreaders: First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, shutting off the spreader before turning as you reach the edge of the border strips. (directional diagram)

To determine how much you need, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, gardens, etc.) One bag covers X,XXX sq. ft.

Optional:		
	Length	(Property Diagram)
×	Width	
-	Total Sq. Ft.	
	Non-lawn areas	
	Total sq. ft. coverage you need	

Directions for commercial products:

Suggested Spreader Settings

These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.

> XX lb. treats XX,XXX sq. ft. at the 1.5 lb./1,000 sq.ft. rate XX lb. treats XX,XXX sq. ft. at the 3.0 lb./1,000 sq.ft. rate XX lb. treats XX,XXX sq. ft. at the 6.0 lb./1,000 sq.ft. rate XX lb. treats XX,XXX sq. ft, at the 12.0 lb./1,000 sq.ft. rate

SPREADER	GROUND OR PTO SPEAD	WIDTH OF COVERAGE	SPREADER SETTINGS lbs./1,000 sq. ft.				
			1.5	3.0	6.0	12.0	15.0
The Andersons Rotaries AccuPro/SR-2000 (cone X)	X mph	XX ft/	х	х	х	х	х
Scotts Rotaries** R-8, R-8A (cone X)	X mph	XX ft	х	х	х	х	х
Lely Models WTR, WFT, HR 1250 (PTO at 450 rpm)	X mph	XX ft	х	Х	х	×	х

Note to EPA: actual spreader models may vary depending on market demand and availability

STORAGE AND DISPOSAL

For homeowner products:

STORAGE: Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies.

DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available. If party filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

For commercial products

Do not contaminate water, food or feed by storage and disposal. PESTICIDE STORAGE: Keep ** container closed and away from food, feedstuffs, and domestic water supplies. PESTICIDE. DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, ... or rinsate, is a violation of Federal Law and may contaminate groundwater. If tkeໍ້ຂ ພໍ້astes cannot be disposed of by use according to label instructions, contact your State Pesticide or **** Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA****

Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVamd052203)

Attention: This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.

Optional (subregistrant) marketing information:

Frank's Quality Guarantee: Frank's takes pride in offering only the highest quality. Every Frank's product is formulated by our experts to help you enjoy a beautiful lawn. You'll feel confident that you made the 'quality' choice when you choose Frank's.

Manufactured by: The Andersons Lawn Fertilizer Division d/b/a Free Flow Fertilizer P.O. Box 119 Maumee, OH 43537

Date Printer			
[13]#	the contract of the contract o		
inipelite F3	1.60.95	pagagagan ang pagamanan an	(· k (
John William	1 (ct 1) shon = 0.4 From 65		در در او آواز کار کر سالگرستان
្តីមិន្តស្រាក់ 	Market Co.		G 1 E
		5 1 9 1	٠
	+	. ,	6 1
		• • • •	•
		2 1	• • •
		 • → * * * 	* * *
		® a o a a e	•
			8 4 2 3
			• :
			ş. ,





The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

May 3, 2004

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7505C)
US Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, DC 20460-0001

RE:

Notification of Minor Label Change

SUBJECT:

The Andersons 0.5% Bayleton Fungicide

EPA Reg. No. 9198-112

Per PR Notice 98-10 enclosed is an EPA Form 8570-1, an application notifying the Agency of a revision to the labeling for the subject registration. One copy of the revised labeling is also enclosed with the additions underlined and marked in blue.

We have made the following change by notification:

1. Added the following required statement for California:

Attention: This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.

Enclosed is a card indicating the acceptance or rejection of this notification. Please complete the card and return via mail. If there are any questions or comments regarding this notification, please feel free to contact me.

Sincerely.

The Andersons Lawn Fertilizer Division, Inc.

Dawn & Walters

Dawn L. Walters Regulatory Administrator Phone 334-262-2766 Fax 334-265-5703

Email dawn walters@andersonsinc.com

enclosures

