

9198-117 4-18-2008

1/36



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

APR 18 2008

Ms. Dawn Walters
The Andersons Lawn Fertilizer Division, Inc.
d/b/a Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537

Subject: Label Notifications for Pesticide Registration Notice 2007-4

Dear Ms. Walters,

The Agency is in receipt of your Applications for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated March 3, 2008 and April 8, 2008 for the following products:

The Andersons Dimension Herbicide IV	EPA Registration Number 9198-117
The Andersons Fertilizer w/0.38% Barricade Herbicide	EPA Registration Number 9198-163
The Andersons 0.2% Imidacloprid Insecticide + Fertilizer	EPA Registration Number 9198-236

The Registration Division (RD) has conducted its review of these requests for their applicability under PRN 2007-4 and finds that the label changes requested fall within the scope of PRN 2007-4. The labels submitted with the application have been stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please call me directly at 703-305-6249 or Steve Schaible of my staff at 703-308-9362.

Sincerely,



Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

3/26



United States
Environmental Protection Agency
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 9198-117	2. EPA Product Manager Kraft	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Dimension Herbicide IV	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fertilizer Division, Inc. d/b/a Free Flow Fertilizer PO Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Label change per PR Notice 2007-4.

This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
				<input checked="" type="checkbox"/> Glass	<input checked="" type="checkbox"/> Paper
				Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 15-2,000 lbs.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label	
6. Manner in Which Label is Affixed to Product		<input checked="" type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input checked="" type="checkbox"/> Other Pocket	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Dawn L. Walters	Title Regulatory Administrator	Telephone No. (Include Area Code) 419-897-6790
<p align="center">Certification</p> <p>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</p>		6. Date: Application Received (Stamped)
2. Signature	3. Title Regulatory Administrator	
4. Typed Name Dawn L. Walters	5. Date 4/3/08	



The Andersons, Inc.
P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

March 3, 2008

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
US Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

RE: Notification of Minor Label Change
SUBJECT: The Andersons Dimension Herbicide IV
EPA Reg. No. 9198-117

Per PR Notice 2007-4 enclosed is EPA Form 8570-1, an application notifying the Agency of a revision to the labeling for the subject registration. There are separate homeowner and commercial versions of the labeling. Copies of the both versions of the revised labeling are enclosed with the additions underlined and marked in blue.

As consistent with the guidance in PR Notice 2007-4, "Labeling Revisions Required by the Final Rule 'Pesticide Management and Disposal; Standards for Pesticide Containers and Containment'", we have added the following statements to our Storage and Disposal section:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.


If there are any questions or comments regarding this notification, please contact me.

Sincerely,
The Andersons Lawn Fertilizer Division, Inc.

Dawn L. Walters

Dawn L. Walters
Regulatory Administrator
Phone: 419-897-6790
Fax: 419-891-2745
Email: dawn_walters@andersonsinc.com

Enclosures



(FRONT PANEL)
COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

THE ANDERSONS DIMENSION HERBICIDE IV

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

Optional marketing claims:

- Utilizes Poly-S® for slow release nitrogen feeding
- Includes DG Pro® for enhanced herbicide distribution
- Contains potassium sulfate for reduced burn potential

ACTIVE INGREDIENTS:

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S- dimethyl ester.....	0.25%
---	-------

INERT INGREDIENTS	<u>99.75%</u>
--------------------------------	---------------

TOTAL.....	100.00%
------------	---------

EPA Reg. No. 9198-117
EPA Est. No. 9198-OH-1


KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Visit our website: AndersonsGolfProducts.com
SGN XXX
Covers up to XX,XXX Square Feet

NET WEIGHT _____ LBS.



(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

NET WEIGHT _____ LBS.

The Andersons Dimension Herbicide IV

GUARANTEED ANALYSIS

Total Nitrogen (N).....	16%
Available Phosphate (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	4%
Primary Plant Food derived from: Urea, Muriate of Potash, Ammoniated Phosphates.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN
CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

- If in eyes Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
- If swallowed Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- If on skin or clothing Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
- If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Cool-Season Grasses

- Bentgrass, creeping*
- Agrostis palustris
- Bluegrass, Kentucky
- Poa pratensis
- Fescue, fine**
- Festuca rubra
- Fescue, tall
- Festuca arundinacea
- Ryegrass, perennial
- Lolium perenne

Warm-Season Grasses

- Bahiagrass
- Paspalum notatum
- Bermudagrass
- Cynodon dactylon
- Buffalograss***
- Buchloe dactyloides
- Carpetgrass
- Axonopus affinis
- Centipedegrass
- Eremochloa ophiuroides
- Kikuyugrass
- Pennisetum clandestinum
- St. Augustinegrass
- Stenotaphrum secundatum
- Zoysiagrass
- Zoysia japonica

* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties.

**Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

RESEEDING, OVERSEEDING, OR SPRIGGING

Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

GOLF COURSE PUTTING GREENS

Putting greens should be well-drained and managed to promote the development of deep rooting systems. The use of this product on greens without good root development may result in undesirable turfgrass injury.

For golf courses which practice overseeding putting greens consisting of seasonal turfgrasses, application of this product to overseeded greens should be made within four weeks or less before the desired transition period from the overseeded turfgrass(es) back to the established "base" turfgrass.

Avoid application of this product to golf course putting greens that are under stress.

Undesirable thinning, yellow and rapid decline of established *Poa annua* may occur under stress conditions. DO NOT use this product on greens that have desirable *Poa annua* cover where summer heat and/or other stresses to *Poa annua* can occur and loss of some or all of that *Poa annua* cover is unacceptable. Use this product where the goal is to control and/or eliminate some or all of the *Poa annua* and where this loss can be tolerated.

Use of this product on 'Tifgreen' (328) hybrid bermudagrass greens may result in injury. Prior to use, test on a small section of turf.

DO NOT apply this product less than 60 days before covering greens for winter protection.

OTHER USE PRECAUTIONS

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

Keep people and pets off treated areas until dust has settled.

(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

APPLICATION DIRECTIONS

Table 1: Recommended Application Use Rates*
(lbs of this product/1000 sq. ft.)

PROGRAM	1	2	3	4
USE (turfgrass cut)	Preemergence (high)	Preemergence (low) Postemergence (high)	Preemergence (low) Postemergence (low)	Golf Course Putting Green Mowing Height less than 1/4 inch**
REGION				
North	1.2	1.7	2.3	2.3
Transition	1.7	2.3	3.4	2.3
South	2.3	3.4	4.6	2.3 + 2.3
Coastal South	3.4	4.6**	4.6**	3.4 + 2.3 + 2.3

*DO NOT apply more than 4.6 lbs. per application, and no more than 13.8 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

- North: All other areas.
- Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.
- South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.
- Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

**May require split or sequential applications for full control.

CONTROL OF CRABGRASS

Preemergence and Early Postemergence Control

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

DO NOT apply more than 4.6 lbs. per 1000 sq. ft. per application, and no more than 13.8 lbs. per 1000 sq. ft. per year.

Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring.

The fall application should be followed by an appropriate spring application to provide season-long control.

Tips For Improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications. Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

barnyardgrass	lespedeza (common)
bluegrass (annual, roughstalk)	oxalis (buttercup, creeping & yellow woodsorrel)
goosegrass	purslane (common)
crowfootgrass	speedwell (corn)
foxtail (yellow & green)*	spurge (prostrate & spotted)
kikuyugrass	
ryegrass (annual & perennial)	
smutgrass	

*Also controlled at the Program 2 rates.

Fall Applications. Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

bluegrass (annual)-Poa annua	geranium (Carolina)
bittercress	parsley-piert
chickweed	pineappleweed
henbit	shepherdspurge

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

Suggested Spreader Settings
(Available at final printing)

(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION - LANDSCAPE ORNAMENTALS

This product provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces where ornamental plants are grown for aesthetic purposes in ornamental gardens, and residential properties.

Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES

Apply this product with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following table for product recommendations when treating a small area.

DO NOT apply more than 4.6 pounds per 1,000 square feet per application and no more than 13.8 pounds (4.6 pounds in New York) per 1,000 square feet per year. Do not apply in Nassau and Suffolk Counties of New York.

**Pounds of Product
per 1,000 square feet**
4.6

**Ounces of Product
per 100 Square Feet**
7.36

APPLICATION TIMING AND RECOMMENDATIONS

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS

- Only apply this product to established ornamentals.
- Do not apply this product directly to bare root of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.

(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- Do not apply this product on grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing condition. Follow directions given above to determine plant tolerance under your growing conditions prior to large scale use.

COMMON NAME	BOTANICAL NAME	TOLERANT CULTIVARS
Abelia, Dwarf	Abelia X Grandiflora	Nana
Ajuga	Ajuga reptans Ajunga genevensis	Bronze Bronze Beauty
Almond, Flowering	Prunus gladulosa	
Apple*	Malus pumila	
Arbovitae	Thuja occidentalis	Nigra Pyramidalis Smaragh Techny Woodwardii
Arborvitae, Golden	Thuja orientalis	
Aster, Chinese	Callistephus chinensis	Dwarf Queen
Ash, Green	Fraxinus pennsylvanica	
Ash, Mountain	Sorbus acucparia	
Ash, Purple	Fraxinus americana	
Azalea	Rhododendron spp.	Brillant Buccaneer Carror Chimes (Belgian) Exbury Fashion Hardijzer Beauty Hershey Red Higasa Hinocrimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow Southern Charm
Azalea, Flame	Rhododendron calendulaceum	
Azalea, Kirishima		
Bamboo, Heavenly		
Barberry	Berberis thunbergii	Aurea Dwarf Pigmy

(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Green Kobold Pygmy Red Rose Glow
Barberry, Purple		Atropurpurea
Bearberry (common)	Arctostaphylos uva-ursti	Massachusetts
Birch, River	Betula nigra	
Blueberry*	Vaccinium spp.	Bluecorp Blue Jay Jersey North Blue Northland
Bottlebrush	Callistemon citrinum	
Boxwood, Japanese		Japonica
Boxwood, Weller	Buxus sempervirens	
Broom	Cytisus spp.	Moonlight
Broom	Genista pilosa	Vancouver Gold
Bugle Carpet		
Carex, Variegated	Carex	
Camellia	Camellia japonica Camellia sasangua	Debutante Mathotiana Supreme Chansonette
Cedar, Red	Juniperus virginiana	
Cockscomb, Plumosa	Celosia cristata	Red Kewpie
Coleus	Coleus blumei	
Copper leaf	Acalypha wilkesiana	
Cotoneaster	Cotonester apiculatus	
Cycads	Cycads revoluta	
Coyotebrush	Baccharis pelularis	
Cypress, Bald	Taxodium distichum	
Cypress, Italian	Cupressus sempervirens	Glauca
Cypress, Leyland	Cupressocyparis leylandii	
Cypress, Japanese False	Chamaecyparis obtusa	Gracilis
Daffodil	Narcissus spp.	King Alfred
Daylilly	Hemerocallis spp.	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilson's Yellow
Dogwood	Cornus florida	
Dogwood, American	Cornus sericea	
Douglas Fir	Pseudotsuga menziesii	
Dusty Miller	Senecio cineraria	
Elm, Drake	Ulmus parvifolia	
Euonymus	Euonymus fortunei	Argenteo-variegata Auereo-marginata Colorata Emerald Gaiety Emerald 'N Gold Gold Edge Gold Princess

(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Silver King Tricolor Vegetus
Fan Palm, European	Chamaerops humilis	
Fan Palm, Mexican	Washingtonia robusta	
Fescue	Festuca glauca	
Fetterbush	Leucothoe fontiana	Rainbow
Ficus	Ficus retusa	Nitidia
Forsythia	Forsythia X intermedia	Arnold Dwarf Bronxensis Dwarf Lynwood Gold Meadowlark Weeping
	Spring Glory	
Fir Fraser	Abies fraseri	
Gardenia	Gardenia jasminoides	Mystery Radicans
Geranium	Pelargonium X hortorum	
Gum	Eucalyptus citriodora	
Hawthorn	Crataegus spp.	Cockspur White Crimson Cloud Enchantress Jack Evans Washington White
Heather, Twisted	Erica cinerea	Mediterranean Pink
Hibiscus	Hibiscus spp.	Brilliant Hula Girl Blue Bird
Hemlock, Canada	Tsuga canadensis	
Holly	Ilex spp. Ilex X meserveae	Blue Boy Blue Girl Burfodii China Girl Ciompacta Forsteri Hellerie Japanese Northern Beauty Needlepoint Nellie R. Stevens Savanah
	Ilex X attenuata	
Holly, Chinese	Ilex cornuta	
Holly, Japanese	Ilex crenata	
Holly, Yaupon	Ilex vomitoria	
Honeysuckle	Lonicera japonica	Clavey's Dwarf Halliana Tatarian Canadian White Zebelli Red Hosta
Hosta	Hosta spp. Hosta lancifolia	Algo Marginata
Ice Plant	Carpobrotus edulis	
Impatiens	Impatiens wallerana	
Iris	Iris spp.	Wedgewood

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

		Dwarf blue
Ivy, English	Hedera helix	Bulgaria Thorndale
Jasmine, Asian	Trachelopermum asiaticum	
Juniper	Juniperus spp.	Arcadia Armstrong Bar Harbor
	Juniperus horizontalis	Blue Chip Blue Pacific Blue Rug
	Juniperus horizontalis	Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Endora Compacta Fuitlandi Green Gold Tip Hetzi Hughes Manhattan Blue. Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blude Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan
	Juniperus chinensis	Tamariscifolia
	Juniperus sabina	Weberi Youngstown Yukon Belle
King Palm	Archontophoenix cunninghamiana	
Laurel, Australian	Pittosporum tobira	
Laurel, Mountain	Kalmia latifolia	
Leucothoe	Leucothoe fontanesiana	
Lily, Africian	Agapanthus africanus	Albus Peter Pan
Lily, African Blue		
Lily of the Valley	Pieris japonica	Mt. Fire
Lilyturf	Liriope muscari	Evergreen Giant Lilac Beauty Majestic Monroe White

(BACK PANEL)

COMMERCIAL USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Variegata
Ligustrum, Japanese	Ligustrum japonicum	
Liriope, Green	Liriope spicata	
Maple, Japanese	Acer japonicum	
Maple, Norway	Acer platanoides	
Maple, Red	Acer rubrum	
Maple, Siviler	Acer saccharinum	
Maple, Sugar	Acer saccharum	
Marigold	Tagetes patula	Honeycomb Wheeler's Dwarf Variegata
Mock Orange*	Philadelphus spp	Golden Snowflake Double White
Mopndo Grass	Ophiopogon japonicus	
Moss Rose	Portulaca grandiflora	Sunnyside
Myrtle, Crape	Lagerstoemia indica	Faurei Langer Muskogee Standard Pink
Myrtle, Wax	Myrica cerifer	
Nadina	Nadina domestica	Compacta Nana
Oleander	Nerium oleander	Hardy Red Petite Pink Sister Agnes
Oak, Laurel	Quercus laurifolia	
Oak, Pin	Quercus palustris	
Oak, Red	Quercus rubra	
Oak, Southern L	Quercus virginiana	
Oak, Willow	Quercus phellos	
Oteospermum	Osteospermum fruitcosum	Wirligig
Palm, Bangalow		
Pampas Grass	Cortadaria selloana	
Paper Flower	Bougainvillea glabra	Barbara Karst
Pachysandra	Pachysandra terminalis	
Peach*	Prunus persica	
Periwinkle, Dwarf	Vinca minor	
Petunia	Petunia X hybrida	Picoti
Photinia, Red Tip	Photinia X fraseri	
Pieris	Pieris japonica	
Pine, Australian	Pinus nigra	
Pine, Japanese Black	Pinus thumbergiana	
Pine, Loblolly	Pinus teada	
Pine, Longleaf	Pinus palustris	
Pine, Mugho	Pinus mugho	
Pine, Scotch	Pinus sylvestris	
Pine, Slash	Pinus eliottii	
Pine, Swiss Mt.	Pinus mugo	
Pine, Virginia	Pinus virginiana	
Pine, White	Pinus strobus	
Pineapple, Guava	Feijoa fellowaiana	

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

Pittosporum, Japan		
Potentilla	Potentilla fruticosa	Abbotswood
Privet	Ligustrum japonicum	Golden Vicary Regal Texanum Yellow Tipped Wax
Privet, Glossy	Ligustrum lucidum	
Pyracantha	Pyracantha koidzumii	
Queen Pam	Arecastrum rammanzoffianum	
Quince, Japanese*		
Rhododendron	Rhododendron spp	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina	Rhododendron carolinianum	
Rhododendron, Catawba	Rhododendron catawbiense	
Rodie Max (Rosebay)	Rhododendron maximum	
Rose	Rosa banksiaae	Luta
Rosemary*	Rosmarinum officinalis	
Rosemary, Bog	Andromeda polifolia	Nana
Salvia	Salvia farinacea	Rhea
Sedum	Sedum spurium	Dragon Blood Red Red Carpet Yellow
Spiraea	Astilbe X arendsii	Fanall
Spiraea	Spiraea spp.	Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houttee White
Spiraea, Garland	Spiraea spp.	
Spruce, Black Hills	Picea glauca	
Spruce, Colorado Blue	Picea pungens	
Spruce, Norway	Picea abies	
Spruce, White	Picea glauca	
Sourwood	Oxydendrum arboreum	
Sweetflag, Grassyy-Le		
Sweetgum	Liquidambar styraciflua	
Sycamore	Platanus occidentalis	
Tree Fern (Tiki Fern)	Asparagus virgatus	
Trumpet Flower, Evening	Gelsemium sempervirens	
Tulip	Tulip, spp	Apeldoorn
Verbena, Shrub	Lantana sellowiana	
Vervain	Verbena spp.	St. Paul
Viburnum	Viburnum spp.	American Cranberry Bush Arrowood

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

		Common Snowball European Cranberry Bush Linden Mohican Wright
Windmill Palm	<i>Trachycarpus fortunei</i>	
Xylosma	<i>Xylosma congestum</i>	
Yaupon	<i>Ilex vomitoria</i>	Dwarf
Yew	<i>Taxus cuspidata</i> <i>Taxus X media</i>	Denisiformis

***Ornamental species only: DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS**

WEEDS CONTROLLED

When used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established grasses except for crabgrass in early stages of development, and broadleaf weeds, therefore, the area to be treated should be free of weeds prior to application.

GRASSES

- barley (*Hordeum* spp.)
- barnyardgrass (*Echinochloa crus-galli*)
- bluegrass, annual (*Poa annua*)
- brome (*Bromus* spp.)
- crabgrass, large (*Digitaria sanguinalis*)
- crabgrass, smooth (*Digitaria ischaemum*)
- crabgrass, Southern (*Digitaria ciliaris*)
- crowfootgrass (*Dactyloctenium aegyptium*)
- dallisgrass (seedling) (*Paspalum dilatatum*)
- goosegrass (*Eleusine indica*)
- foxtail, green (*Setaria verdi*)
- foxtail, yellow (*Setaria faberi*)
- kikuyugrass (*Pennisetum clandestinum*)
- oats, wild (*Avena fatua*)
- ryegrass (annual & pernnial) (*Lolium* spp.)
- smutgrass (*Sporobolus poireti*)

BROADLEAF WEEDS

- bittercress (*Cardamine* spp.)
- carpetweed (*Mollugo verticillata*)
- chickweed (*Stellaria* spp.)
- geranium, Carolina (*Geranium carolinianum*)
- henbit (*Lamium* spp.)
- knotweed, prostrate (*Polygonum saviculare*)
- lespedeza, common (*Lespedeza striata*)
- marestail (*Hippuris vulgaris*)
- medic, black (*Medicago lupulina*)
- mustard (*Brassica* spp.)
- oxalis, buttercup (*Oxalis pes-caprae*)
- pineappleweed (*Matricaria matricarioides*)
- pigweed, redroot (*Amaranthus retroflexus*)
- parsley-piert (*Alchemilla avrensis*)
- purslane, common (*Portulaca oleracea*)
- rocker, London (*Sisymbrium irio*)
- shepherdspurse (*Capsella bursa-pastoris*)
- speedwell, corn (*Veronica arvensis*)
- spurge, garden (*Euphorbia hirta*)
- spurge, prostrate (*Euphorbia humistrata*)
- spurge, spotted (*Euphorbia maculata*)
- woodsorrel, creeping (*Oxalis corniculata*)
- woodsorrel, yellow (*Oxalis stricta*)

2/1/36

(BACK PANEL)

COMMERCIAL USE ONLY

(The above is for internal use only and will not appear on the final label copy)

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVnot040808)

Manufactured by:

The Andersons Lawn Fertilizer Division, Inc.
d/b/a Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537
Visit our website: AndersonsGolfProducts.com

Date Printed	4/8/08
Date Approved	5/8/03
Supersedes	5/26/98
Label Changes	Notification -container
Filename	9198-117c
Label Code	RVnot040808

36

(FRONT PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

The Andersons Dimension Herbicide IV

- A Herbicide Plus Fertilizer Combination Product that Provides Control of Crabgrass and Other Weeds in Established Lawns and Ornamental Turfgrasses.
- Premium Granular Fertilizer that Feeds Your Lawn - Providing The Nutrients Needed for A Beautiful, Green and Vigorous Lawn.
- Contains Dimension® Herbicide - A Herbicide for Crabgrass Control!
- Can Be Applied Up to 4 Weeks Later Than Other Crabgrass Preventers and Controls Crabgrass All Season Long.

ACTIVE INGREDIENTS:

*Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)
-6-(trifluoromethyl)-S,S- dimethyl ester..... 0.25%

INERT INGREDIENTS..... 99.75%
TOTAL..... 100.00%

® Dimension is a registered trademark of Dow AgroSciences LLC


KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT _____ LBS.
Covers {over} X,XXX square feet

EPA Reg. No. 9198-117
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code on bag



(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

NET WEIGHT _____ LBS.

The Andersons Dimension Herbicide IV

GUARANTEED ANALYSIS

Total Nitrogen (N).....	16%
Available Phosphate (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	4%
Primary Plant Food derived from: Urea, Muriate of Potash, Ammoniated Phosphates.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN

CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.


FIRST AID

- If in eyes Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
- If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
- If swallowed Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- If on skin or clothing Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.



(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 4.6 pounds per 1000 square feet (equal to 0.5 pound active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

Use only according to label instructions.

GENERAL INFORMATION

This product is a combination product that provides both crabgrass control and the required nutrients for optimum turfgrass growth all in a single product.

This product contains Dimension, a crabgrass preventer that can be applied up to 4 weeks later than other crabgrass preventers and provide up to 4 months control of crabgrass and other grass and broadleaf weeds prior to their establishment in lawns and ornamental turfgrasses. This product should be applied to lawns and ornamental turfgrasses that are well established (See "Use Precautions" section below for a list of the tolerant turfgrasses). This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

This product should only be applied to lawns and ornamental turfgrasses that are well established. Use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury.

The Andersons Dimension Herbicide IV can be applied to the following turfgrasses when applied as directed:	
<u>Cool-Season Turfgrasses</u> <ul style="list-style-type: none">• Creeping Bentgrass• Fine Fescues• Kentucky Bluegrass• Perennial Ryegrass• Tall Fescues	<u>Warm-Season Turfgrasses</u> <ul style="list-style-type: none">• Bahiagrass• Bermudagrass• Buffalograss• Carpetgrass• Centipedegrass• St. Augustinegrass• Zoysiagrass

This product should be applied in the spring or fall prior to the weed seed germination in established lawns and ornamental turfgrasses.

28/36

(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

This product can be applied up to 4 weeks later than other crabgrass preventers and still provide season-long weed control of crabgrass. Dimension herbicide provides preemergence control of crabgrass and other weeds when applied prior to their germination, and it also provides postemergence control of crabgrass. The postemergence control provided by this product is limited to crabgrass only and will provide control of crabgrass when applied up to 4 weeks after crabgrass germination. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass.

For control of other grass and broadleaf weeds, this product must be applied as a preemergent treatment. Only crabgrass can be controlled with delayed applications after germination.

Tips for Improved Weed Control: For best results, apply this product within a few days after mowing and delay further mowings after the application of this product until the lawn or ornamental turfgrass has been watered or received significant rainfall to dissolve the fertilizer granules. Improved weed control may be observed when treated lawn or ornamental turfgrass is watered or receives rainfall within a few days after application of this product.

Reseeding, Overseeding, or Sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until 16 weeks after application. Reseeding, overseeding, or sprigging upto 16 weeks after application may prevent the establishment of the newly reseeded, overseeded or sprigged turfgrasses.

•DO NOT apply more than 4.6 pounds of this product per 1,000 square feet of turfgrass area during the year (1,000 square feet area = 100 feet x 10 feet).

•DO NOT apply to flowers, vegetables, or fruit trees.

•DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees.

•Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

The Andersons Dimension Herbicide IV will provide control of crabgrass and other grasses and broadleaf weeds listed below when applied prior to weed seed germination in established lawns and ornamental turfgrasses.

Weeds Controlled and/or Suppressed by The Andersons Dimension Herbicide IV

Grass Weeds:

Annual Bluegrass
Annual Ryegrass
Barnyardgrass
Crabgrass
Crowfootgrass
Goosegrass

Green Foxtail
Kikuyugrass (seedling)
Perennial Ryegrass
Smutgrass (seedling)
Yellow Foxtail

Broadleaf Weeds:

Bittercress
Buttercup Oxalis
Carolina Geranium
Chickweed
Common Lespedeza
Common Purslane
Corn Speedwell

Creeping Oxalis
Henbit
Parsley-piert
Pineappleweed
Spotted Spurge
Yellow Woodsorrel

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

Application Timings:

This product may be applied either in the spring or in the fall for the control of weeds.

When this product is applied in the fall, a sequential or second application should be made the following spring to provide season-long control of weeds.


Length of Control:

This product will provide up to 4 months of weed control if applied according to label directions.

The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States that this product is being applied.

Please check the map below to determine which "Region" of the U.S. that applies to your particular use area.

Use "Regions" within the United States:



Recommended Use Rates:

Recommended Use Rates for The Andersons Dimension Herbicide IV (Pounds of this product to apply per 1,000 square feet of turfgrass)		
<u>Region of U.S.</u>	<u>Length of Weed Control</u>	
	<u>2-3 Months Control</u>	<u>3-4 Months Control</u>
North	1.15	1.6
South	1.8	2.3

(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

SUGGESTED SPREADER SETTINGS

(Available at Final Printing)

Broadcast Spreaders: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

Drop Spreaders: First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

Important: It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. Do not hand spread.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs and domestic water supplies.

DISPOSAL:

If empty: ~~Do not reuse this container. Place in trash or offer for recycling if available. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.~~

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL INFORMATION - LANDSCAPE ORNAMENTALS

This product provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces where ornamental plants are grown for aesthetic purposes in ornamental gardens, and residential properties.

Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES

Apply this product with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following table for product recommendations when treating a small area.

DO NOT apply more than 4.6 pounds per 1,000 square feet per year. Do not apply in Nassau and Suffolk Counties of New York.

**Pounds of Product
per 1,000 square feet
4.6**

**Ounces of Product
per 100 Square Feet
7.36**

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

APPLICATION TIMING AND RECOMMENDATIONS

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS

- Only apply this product to established ornamentals.
- Do not apply this product directly to bare root of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- Do not apply this product on grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing condition. Follow directions given above to determine plant tolerance under your growing conditions prior to large scale use.

COMMON NAME	BOTANICAL NAME	TOLERANT CULTIVARS
Abelia, Dwarf	<i>Abelia X Grandiflora</i>	Nana
Ajuga	<i>Ajuga reptans</i> <i>Ajunga genevensis</i>	Bronze Bronze Beauty
Almond, Flowering	<i>Prunus glandulosa</i>	
Apple*	<i>Malus pumila</i>	
Arbovitae	<i>Thuja occidentalis</i>	Nigra Pyramidalis Smaragh Techny Woodwardii
Arborvitae, Golden	<i>Thuja orientalis</i>	
Aster, Chinese	<i>Callistephus chinensis</i>	Dwarf Queen
Ash, Green	<i>Fraxinus pennsylvanica</i>	
Ash, Mountain	<i>Sorbus acucparia</i>	

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

Ash, Purple	<i>Fraxinus americana</i>	
Azalea	<i>Rhododendron spp.</i>	Brilliant Buccaneer Carror Chimes (Belgian) Exbury Fashion Hardijzer Beauty Hershey Red Higasa Hinocrimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow Southern Charm
Azalea, Flame	<i>Rhododendron calendulaceum</i>	
Azalea, Kirishima		
Bamboo, Heavenly		
Barberry	<i>Berberis thunbergii</i>	Aurea Dwarf Pigmy Green Kobold Pygmy Red Rose Glow
Barberry, Purple		Atropurpurea
Bearberry (common)	<i>Arctostaphylos uva-ursti</i>	Massasschusetts
Birch, River	<i>Betula nigra</i>	
Blueberry*	<i>Vaccinium spp.</i>	Bluecorp Blue Jay Jersey North Blue Northland
Bottlebrush	<i>Callistemon citrinum</i>	
Boxwood, Japanese		Japonica
Boxwood, Weller	<i>Buxus sempervirens</i>	
Broom	<i>Cytisus spp.</i>	Moonlight
Broom	<i>Genista pilosa</i>	Vancouver Gold
Bugle Carpet		
Carex, Variegated	<i>Carex</i>	
Camellia	<i>Camellia japonica</i> <i>Camellia sasangua</i>	Debutante Mathotiana Supreme Chansonette
Cedar, Red	<i>Juniperus virginiana</i>	

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

Cockscomb, Plumosa	<i>Celosia cristata</i>	Red Kewpie
Coleus	<i>Coleus blumei</i>	
Copper leaf	<i>Acalypha wilkesiana</i>	
Cotoneaster	<i>Cotoneaster apiculatus</i>	
Cycads	<i>Cycads revoluta</i>	
Coyotebrush	<i>Baccharis pelularis</i>	
Cypress, Bald	<i>Taxodium distichum</i>	
Cypress, Italian	<i>Cupressus sempervirens</i>	Glaucia
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	
Cypress, Japanese False	<i>Chamaecyparis obtusa</i>	Gracilis
Daffodil	<i>Narcissus spp.</i>	King Alfred
Daylily	<i>Heemerocallis spp.</i>	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilson's Yellow
Dogwood	<i>Cornus florida</i>	
Dogwood, American	<i>Cornus sericea</i>	
Douglas Fir	<i>Pseudotsuga menziesii</i>	
Dusty Miller	<i>Senecio cineraria</i>	
Elm, Drake	<i>Ulmus parvifolia</i>	
Euonymus	<i>Euonymus fortunei</i>	Argenteo-variegata Aureo-marginata Colorata Emerald Gaiety Emerald 'N Gold Gold Edge Gold Princess Silver King Tricolor Vegetus
Fan Palm, European	<i>Chamaerops humilis</i>	
Fan Palm, Mexican	<i>Washingtonia robusta</i>	
Fescue	<i>Festuca glauca</i>	
Fetterbush	<i>Leucothoe fontiana</i>	Rainbox
Ficus	<i>Ficus retusa</i>	Nitidia
Forsythia	<i>Forsythia X intermedia</i> <i>Spring Glory</i>	Arnold Dwarf Bronxensis Dwarf Lynwood Gold Meadowlark Weeping
Fir Fraser	<i>Abies fraseri</i>	
Gardenia	<i>Gardenia jasminoides</i>	Mystery Radicans
Geranium	<i>Pelargonium X hortorum</i>	
Gum	<i>Eucalyptus citriodora</i>	
Hawthorn	<i>Crataegus spp.</i>	Cockspur White

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Crimson Cloud Enchantress Jack Evans Washington White
Heather, Twisted	<i>Erica cinerea</i>	Mediterranean Pink
Hibiscus	<i>Hibiscus spp.</i>	Brilliant Hula Girl Blue Bird
Hemlock, Canada	<i>Tsuga canadensis</i>	
Holly	<i>Ilex spp.</i> <i>Ilex X meserveae</i> <i>Ilex X attenuata</i>	Blue Boy Blue Girl Burfodii China Girl Ciompacta Forsteri Hellerie Japanese Northern Beauty Needlepoint Nellie R. Stevens Savanah
Holly, Chinese	<i>Ilex cornuta</i>	
Holly, Japanese	<i>Ilex crenata</i>	
Holly, Yaupon	<i>Ilex vomitoria</i>	
Honeysuckle	<i>Lonicera japonica</i>	Clavey's Dwarf Halliana Tatarian Canadian White Zebelli Red Hosta
Hosta	<i>Hosta spp.</i> <i>Hosta lancifolia</i>	Algo Marginata
Ice Plant	<i>Carpobrotus edulis</i>	
Impatiens	<i>Impatiens wallerana</i>	
Iris	<i>Iris spp.</i>	Wedgewood Dwarf blue
Ivy, English	<i>Hedera helix</i>	Bulgaria Thorndale
Jasmine, Asian	<i>Trachelopermum asiaticum</i>	
Juniper	<i>Juniperus spp.</i>	Arcadia Armstrong Bar Harbor
	<i>Juniperus horizontalis</i>	Blue Chip Blue Pacific Blue Rug
	<i>Juniperus horizontalis</i>	Blue Star Broadmoor Buffalo Calgary Carpet

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Emerald Sea Emerald Spreader Endora Compacta Fuitlandi Green Gold Tip Hetzi Hughes Manhattan Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blude Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan
	<i>Juniperus chinensis</i>	Tamariscifolia
	<i>Juniperus sabina</i>	Weberi Youngstown Yukon Belle
King Palm	<i>Archontophoenix cunninghamiana</i>	
Laurel, Australian	<i>Pittosporum tobira</i>	
Laurel, Mountain	<i>Kalmia latifolia</i>	
Leucothoe	<i>Leucothoe fontanesiana</i>	
Lily, Africian	<i>Agapanthus africanus</i>	Albus Peter Pan
Lily, African Blue		
Lily of the Valley	<i>Pieris japonica</i>	Mt. Fire
Lilyturf	<i>Liriope muscari</i>	Evergreen Giant Lilac Beauty Majestic Monroe White Variegata
Ligustrum, Japanese	<i>Ligustrum japonicum</i>	
Liriope, Green	<i>Liriope spicata</i>	
Maple, Japanese	<i>Acer japonicum</i>	
Maple, Norway	<i>Acer platanoides</i>	
Maple, Red	<i>Acer rubrum</i>	
Maple, Sivler	<i>Acer saccharinum</i>	
Maple, Sugar	<i>Acer saccharum</i>	
Marigold	<i>Tagetes patula</i>	Honeycomb

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Wheeler's Dwarf Variegata
Mock Orange*	<i>Philadelphus spp</i>	Golden Snowflake Double White
Mopndo Grass	<i>Ophiopogon japonicus</i>	
Moss Rose	<i>Portulaca grandiflora</i>	Sunnyside
Myrtle, Crape	<i>Lagerstoemia indica</i>	Faurei Langer Muskogee Standard Pink
Myrtle, Wax	<i>Myrica cerifer</i>	
Nadina	<i>Nadina domestica</i>	Compacta Nana
Oleander	<i>Nerium oleander</i>	Hardy Red Petite Pink Sister Agnes
Oak, Laurel	<i>Quercus laurifolia</i>	
Oak, Pin	<i>Quercus palustris</i>	
Oak, Red	<i>Quercus rubra</i>	
Oak, Southern L	<i>Quercus virginiana</i>	
Oak, Willow	<i>Quercus phellos</i>	
Oteospermum	<i>Osteospermum fruitcosum</i>	Wiriligig
Palm, Bangalow		
Pampas Grass	<i>Cortadaria selloana</i>	
Paper Flower	<i>Bougainvillea glabra</i>	Barbara Karst
Pachysandra	<i>Pachysandra terminalis</i>	
Peach*	<i>Prunus persica</i>	
Periwinkle, Dwarf	<i>Vinca minor</i>	
Petunia	<i>Petunia X hybrida</i>	Picoti
Photinia, Red Tip	<i>Photinia X fraseri</i>	
Pieris	<i>Pieris japonica</i>	
Pine, Australian	<i>Pinus nigra</i>	
Pine, Japanese Black	<i>Pinus thumbergiana</i>	
Pine, Loblolly	<i>Pinus teada</i>	
Pine, Longleaf	<i>Pinus palustris</i>	
Pine, Mugho	<i>Pinus mugho</i>	
Pine, Scotch	<i>Pinus sylvestris</i>	
Pine, Slash	<i>Pinus eliottii</i>	
Pine, Swiss Mt.	<i>Pinus mugo</i>	
Pine, Virginia	<i>Pinus virginiana</i>	
Pine, White	<i>Pinus strobus</i>	
Pineapple, Guava	<i>Feijoa fellowaiana</i>	
Pittosporum, Japan		
Potentilla	<i>Potentill fruticosa</i>	Abbotswood
Privet	<i>Ligustrum japonicum</i>	Golden Vicary Regal

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Texanum Yellow Tipped Wax
Privet, Glossy	<i>Ligustrum lucidum</i>	
Pyracantha	<i>Pyracantha koidzumii</i>	
Queen Pam	<i>Arecastrum rammanzoffianum</i>	
Quince, Japanese*		
Rhododendron	<i>Rhododendron spp</i>	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina	<i>Rhododendron carolinianum</i>	
Rhododendron, Catawba	<i>Rhododendron catawbiense</i>	
Rodie Max (Rosebay)	<i>Rhododendron maximum</i>	
Rose	<i>Rosa banksiaae</i>	Luta
Rosemary*	<i>Rosmarinum officinalis</i>	
Rosemary, Bog	<i>Andromeda polifolia</i>	Nana
Salvia	<i>Salvia farinacea</i>	Rhea
Sedum	<i>Sedum spurium</i>	Dragon Blood Red Red Carpet Yellow
Spiraea	<i>Astilbe X arendsii</i>	Fanall
Spiraea	<i>Spiraea spp.</i>	Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houttee White
Spiraea, Garland	<i>Spiraea spp.</i>	
Spruce, Black Hills	<i>Picea glauca</i>	
Spruce, Colorado Hlue	<i>Picea pungens</i>	
Spruce, Norway	<i>Picea abies</i>	
Spruce, White	<i>Picea glauca</i>	
Sourwood	<i>Oxydendrum arboreum</i>	
Sweetflag, Grassyy-Le		
Sweetgum	<i>Liquidambar styraciflua</i>	
Sycamore	<i>Platanus occidentalis</i>	
Tree Fern (Tiki Fern)	<i>Asparagus virgatus</i>	
Trumpet Flower, Evening	<i>Gelsemium sempervirens</i>	
Tulip	<i>Tulip, spp</i>	Apeldoorn
Verbana, Shrub	<i>Lantana sellowiana</i>	
Vervain	<i>Verbena spp.</i>	St. Paul
Viburnum	<i>Viburnum spp.</i>	American Cranberry Bush

(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

		Arrowwood Common Snowball European Cranberry Bush Linden Mohican Wright
Windmill Palm	<i>Trachycarpus fortunei</i>	
Xylosma	<i>Xylosma congestum</i>	
Yaupon	<i>Ilex vomitoria</i>	Dwarf
Yew	<i>Taxus cuspidata</i> <i>Taxus X media</i>	Denisiformis

*Ornamental species only: DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS

WEEDS CONTROLLED

When used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established grasses except for crabgrass in early stages of development, and broadleaf weeds, therefore, the area to be treated should be free of weeds prior to application.

GRASSES

- barley (*Hordeum* spp.)
- barnyardgrass (*Echinochloa crus-galli*)
- bluegrass, annual (*Poa annua*)
- brome (*Bromus* spp.)
- crabgrass, large (*Digitaria sanguinalis*)
- crabgrass, smooth (*Digitaria ischaemum*)
- crabgrass, Southern (*Digitaria ciliaris*)
- crowfootgrass (*Dactyloctenium aegyptium*)
- dallisgrass (seedling) (*Paspalum dilatatum*)
- goosegrass (*Eleusine indica*)
- foxtail, green (*Setaria verti*)
- foxtail, yellow (*Setaria faberi*)
- kikuyugrass (*Pennisetum clandestinum*)
- oats, wild (*Avena fatua*)
- ryegrass (annual & pernnial) (*Lolium* spp.)
- smutgrass (*Sporobolus poireti*)

BROADLEAF WEEDS

- bittercress (*Cardamine* spp.)
- carpetweed (*Mollugo verticillata*)
- chickweed (*Stellaria* spp.)
- geranium, Carolina (*Geranium carolinianum*)
- henbit (*Lamium* spp.)
- knotweed, prostrate (*Polygonum saviculare*)
- lespedeza, common (*Lespedeza striata*)
- marestail (*Hippuris vulgaris*)
- medic, black (*Medicago lupulina*)
- mustard (*Brassica* spp.)
- oxalis, buttercup (*Oxalis pes-caprae*)
- pineappleweed (*Matricaria matricarioides*)
- pigweed, redroot (*Amaranthus retroflexus*)
- parsley-piert (*Alchemilla avrensis*)
- purslane, common (*Portulaca oleracea*)
- rocker, London (*Sisymbrium irio*)
- shepherdspurse (*Capsella bursa-pastoris*)
- speedwell, corn (*Veronica arvensis*)
- spurge, garden (*Euphorbia hirta*)
- spurge, prostrate (*Euphorbia humistrata*)
- spurge, spotted (*Euphorbia maculata*)
- woodsorrel, creeping (*Oxalis corniculata*)
- woodsorrel, yellow (*Oxalis stricta*)

(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy).

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes **NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.** To the extent consistent with applicable law, it is Manufacturer's intent to **LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES** to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer **DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS.** Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVnot053007)

Manufactured by:

The Andersons, Lawn Fertilizer Division, Inc. d/b/a Free Flow Fertilizer
P.O. Box 119
Maumee, OH 43537

Date Printed	5/30/07
Date Approved	
Supersedes	5/26/98
Label Changes	Notification - warranty
Filename	9198-117h
Label Code	RVnot053007