8-117

5181200



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Ms. Debbie Ziehr Product Registration The Andersons Lawn Fert. Div., Inc PO Box 119 Maumee, OH 43537

MAY 8 2009

SUBJECT: Application for Pesticide Notification (PRN 98-10) Request Alternate Brand Name "The Andersons Turf Products Fertilizer with 0.25 Dimension Turf Herbicide10-6-4"

> EPA Reg. No. 9198-117 Application Dated April 14 2009

Dear Ms Ziehr:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 04/14/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

2~

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

| Picase read instructions on | reverse before comple | ating form. | | Form Appro | <u>ved. OMB No. 2</u> | <u>070-0</u> 06 | aofy |
|--|--|--|---|---|---|---|-------------------------------|
| \$-EPA | ر Environmenta | Jnited States | | | Registra Amendn Other | tion | OPP Identifier Number |
| | | Applicati | on for Pestici | de - Sectio | on I | | |
| 1. Company/Product Numb 9198-117 | ər | | 2. EPA Miller | Product Manag | er | 3. Pr | oposed Classification |
| 4. Company/Product (Name The Andersons Dimension I | | | РМ# 23 | | | | |
| 5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow Fert. P.O. Box 119 Maumee, OH 43537 | | | 6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No | | | | |
| Check if thi | s is a new address | | | | | | |
| have and | | | Section - | | | | |
| Amendment - Explai Resubmission in resp Notification - Explain Explanation: Use additio Notification of Alternate Brand Na This notification is consistent with made to the labeling or the confic further understand that if this not violation of FIFRA and I may be s | ponse to Agency letter below. nal page(s) if necessar ame per PR Notice 98-10: In the guidance of PR Notice Jential statement of formula fication is not consistent wi | y. (For sectio The Andersons 98-10 and the ro of this product. th the guidance of | n I and Section II.) Turf Products Fertilizer v equirements of EPA's re I understand that it is a of PR Notice 98-10 and | Agency letter "Me Too" App Other - Explain with 0.25% Dimens gulations at 40 CF violation of 18 U.S he requirements c | blication. n below, ion Turf Herbicide 1 R 156.10 and 40 CF .C. Sec. 1001 to will | 0-6-4 R 152.46 a fully make a | iny false statement to EPA. I |
| | | n and penanies | Section - I | | | | |
| 1. Material This Product Wi | ll Be Packaged In: | | | | | | |
| Child-Resistant Packaging Yes No * Certification must be submitted | Unit Packaging Yes V No If "Yes" Unit Packaging wgt. | No. per container | Water Soluble P Yes No If "Yes" Package wgt | ackaging No. per container | 2. Type of (| Container Metal Plastic Glass Paper Other (S | |
| 3. Location of Net Contents | Information | 4. Size(s) Re | tail Container | 5 | Location of Lab | el Directio | |
| | Container | | 10 - 2,000 lbs. | | On Label | | · |
| 6. Manner in Which Label is | Affixed to Product | Litho Paper Stend | graph glued iled | 🖌 Other | Pocket | | |
| | | | Section - I | V. | | | |
| 1. Contact Point <i>(Complete</i> | items directly below f | for identification | on of individual to b | e contacted, if i | necessary, to pro | cess this | app:lication.) |
| Name Debbie Ziehr | | Title Regulatory Administrator | | | Telephone No. (Include Area Code) 419-891-6671 a Co | | |
| • | ments I have made on by knowlinglly false or law. | | all attachments the | | or imprisonment | · · · · | 6. Date Application |
| 2. Signature Delebie | Ziehr | | 3. Title Regulatory Adminis | strator | | | |
| 4. Tγped Name U Debbie Ziehr | | | 5. Date (((())) | | | | |





The Andersons, Inc. P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

April 13, 2009

Document Processing Desk (NOTIF) Office of Pesticide Programs - (7504P) U. S. Environment Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

Subject: NOTIFICATION - Additional Brand Name

RE: The Andersons Dimension Herbicide IV EPA Reg. No. 9198-117

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons Turf Products Fertilizer with 0.25% Dimension Turf Herbicide 10-6-4

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

rehr ebbee ~

Debbie Ziehr Regulatory Administrator Phone #419-891-6671 Fax #419-891-2745 Email: Debbie_ziehr@andersonsinc.com

Enclosure



• • • • • • • • • • •



Fertilizer with 0.25% Dimension® Turf Herbicide 10-6-4

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

GUARANTEED ANALYSIS

10.00%

| 2.34% Ammoniacal Nitrogen 7.66% Urea Nitrogen | |
|--|-----------------------------------|
| Available Phosphate (P_2O_5). Soluble Potash (K_2O). | 6.00% 4.00% |
| Plant nutrients derived from ammonium phosphate, urea, p coated sulfur coated urea and potassium chloride. | olymer |
| Chlorine (Cl) Max *3.00% Slowły available nitrogen from polymer coated sulfu urea. | 3.40% ur coated |
| | F292 |
| ACTIVE INGREDIENT: Dithicpyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4- (2-methylpropyl)-6-(trifluoromethyl)-5,S-dimethyl ester INERT INGREDIENTS: Total | 0.25% <u>99.75%</u> 100.00% |

Total Nitrogen (N).....

INERT INGREDIENTS: Total

EPA Reg. No. 9198-117 EPA Est, No. 9198-OH-1M, 9198-OH-28, 9198-AL-001A (Underlined letter is first letter used in run code on end of bag.)

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing clust. Wash thoroughly with scap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

It in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If swallowed; Call a poison control center or doctor for treatment advice. Have person sig a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person

If on skin or clothing: Take off contaminated clothing. Binse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inheled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is taxic to fish and highly taxic to other aquatic organisms including system and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertirial areas below the mean high water mark. Do not apply when weather conditions lavor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR LISE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 4.6 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

GENERAL INFORMATION

This product is a herbicide that provides control of craborass and other annual grasses and broadleaf weeds in established lawns and ornamental turls, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turigrass Safety: This product may be used on seeded, sodded, or sprigged lawns and ornamental turis that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seed-ing, sodding, or sprigging before it can receive its first application of this product. Use of this product on Jawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

| Cool-Sesson Grasses Benigrass, creeping" (Agrostis palustris) Bluegrass, Kontucky (Poa pratensis) Fescue, fine" (Festuca rubra) Fescue, dial (Festuca rubra) Regrass, perennial (Lolium perenne) | Warm-Soacon Grasses Bahagrass (Pasnahum notatum) Bermudagrass (Cynodon dactylon) Buttalograss" (Buchloe dactyloides) Carpeigrass (Axonopus affinis) Centipedograss (Pennisetun clandeshum) Kikuyugrass (Pennisetun clandeshum) |
|---|--|
| | Kikuyugrass (Pennisetum clandestinum) St. Augustinegrass (Stenotaphrum secunda Zoysiagrass (Zoysia japonica) |
| *Use of this product on certain varieties | of creeping bentgrass, such as 'Cohansey' |

'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (Agrostis tenuis) varieties.

"Use of this product on certain varieties of fine feacue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary'. 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf' and "Waldina" Not all varieties of fine feacue have been tested

""DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turforass is fully green and has established new roots

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

Reserving, Overseeding, or Sprigging; Reserving, overseeding, or springing of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its lifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply rs, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

Control of Crabgrass Preemergence and Early Postemergence Control: This product provides "preemargence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of craborass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it i often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for craburass control in the spring, summer, or fall. DO NOT apply more than 4.6 lbs. per 1,000 sq. ft, per application, and no more than 13.8 Lbs. per 1,000 sq. ft, per year.

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somehat depending on weather, weed pressure, turigrass competitiveness, and the user's location within a region.

Use Program 1 for pregnergence control at sites where the turgrass is cut high (e.g., homeowner lawne). The program provides 3-5 months of preemergence crabgrass control. This program affocula not be used for early postemergence crabgrass control.

Use Program 2 for programs gence control at sites where (a) turigrass is cut low (e.g., golf fairways) and (b) turigrass maintenance or weed control has been conducted during the previous year. This grogram may allow used for any postemergence crabgrass control. This program may allow used for any postemergence control at sites where turigrass is cut high (e.g., horizowner laws)

Use Program 3 for reamergence control at sites where (a) lufgrass is cut low (e.g., golf fairways), and (b) rist maintenents or weed control has not been conducted during the previous year. This program movies 4-6 months of preemergence crabgrass control. This program may the beacting early postemergence control at sites where turfgrass is cut low (e.g., go antways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

| | TABLE 1: Recommended (Lbs. of this prod | | • | |
|------------------------|--|-------------------------|------------------------|--|
| PROGRAM | 1 | 2 | 3 | |
| USE (turfgrase cut) | Preamergence (high) | Presmergence (low) | Presmargance (low) | |
| REGION | | Postemergence (high) | Postamergance (low) | |
| North | 12 | 1.7 | 2.3 | |
| Transition | 1.7 | 2.3 | 3.4 | |
| South | 2.3 | 3.4 | 4.6** | |
| Coastal South | 3.4 | 4,6** | 4.6** | |

Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below

North: All other areas

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL,

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control: For best results, apply this product within a lew days after mowing and gelay mowing again for a few days after the application. When treated laws or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of solit (half-rate) applications spaced 6-8 weeks apart may provide improved weed control

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barryarguns, blaggass (annual, roughstalk), crawfootgrass, fortail (yellow & green)', goosegrass, klusygrass yegrass (annual & perennial), amutgrass, lespedeza (common), osalis (buttercup, creeping & yellow woodsorrel), purslane (common), speedwell (corn), spurge (prostrate & spotted).

Fall Applications: Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence; bluegrass (annual) - Poa annua, bittercress, chickweed, pineappleweed, henbit, geranium (Carolina), parsley-piert, shepherdspurse

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning property and applying uniform distribution of granules.

| These sugger Piec 50 lbs. covers 29,400 sq. 50 lbs. covers 21,700 sq. 1 50 lbs. covers 14,700 sq. 1 50 lbs. covers 19,800 sq. 1 | sted spreader se calibrate (t. (0.67 acre) (0.50 acre) al t. (0.33 acre) | the 100 lbs. pro at the 152 lbs. p | l intended to slore applyi: oduct/acre l duct/acre M roduct/acre | n replace cal ng product. LOW RATE IEDIUM RATE HIGH RATE | 1.7 lbs/1,00 E - 2.3 lbs/1, • 3.4 lbs/1,00 | 000 sq. tt. 10 sq. tt. |
|--|--|---------------------------------------|--|--|--|-------------------------------------|
| t L | 2 | , | | SPREDER SETTINGS | | |
| | | WID, 1 OP COVEPAGE | LOW ,VATE , 0,12 B ,ai acre | McDRUM+ _ATE _ r ^S lb, eilacte | hait fTE 9.*9.16. aitacre | HEAVY RATE 0.50 B. ai'acre |
| Andersons Accufro 2000~ SIR-2000 (cone 4) | 5 mpn 🦷 🤇 | 10 teet | J 1/2 | - x - | ~L 54 | 814 |
| Andersons SS-2 Drop | 3 mph | averlap wheels | 2 1/4 | 2 3/4 | 119 | 4 1/4 |
| Scotts R-8, R-8A (cons 4) | 5 moh | 10 feet | I | 11.2 | K 1J2 | L 1-2 |
| Lesco 020093 (pattern 0.25) | 3 hiph | g leat | D 1/2 | £ | F 1/2 | н |
| Lesco 705622 (pattern 0.25) | | jae: | <u>_</u> 11 | 12 | 15 | 18 |
| Lety Wodels WTR. WFR HR, 1250 (PTC at 450 mph) Pattern III-B | 4.5 mph - | 36 leet | -4 1/2 | 4 3/4 | 5 1/2 | 5 1/4 |
| Vicon (03 Series) | 4.5 mch | 30 leet | 15 | 22 | 27 | 32 |

STORAGE AND DISPOSAL Do not contaminate water, food, or feed by storage and disposal

STORAGE: Keep container closed and away from food, leedstuffs, and domestic water supplies.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

CONTAINER DISPOSAL: Completely empty bag into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landful or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURFOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SEECIAL CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER BIGHTS THAT VARY FROM STATE/JUBISDICTION TO STATE/JUBISDICTION

Product of USA

Manufactured by: The Andersons Lawn Fertilizer Division, Inc. P.O. Box 119 Maumee, OH 43537 www.andersonslawnproducts.com

The Andersons loop is a repistered trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences LLC

SGN 215 Net Wt. 50 Lbs. (22.68 kg)



"Also controlled at the "Program 2" rates.

IN, OH, & coastal areas of CT, NY, & RI, South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX, Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX, "May require split or sequential applications for full control

latum}

*DO NOT apply more than 4.6 lbs, per application, and no more than 13.8 lbs, per year.