

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Ms. Debbie Ziehr Product Registration The Andersons Lawn Fert. Div., Inc PO Box 119 Maumee, OH 43537

MAY 2 1 2009

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request Alternate Brand Names "The Andersons Turf Products

Fertilizer with 0.25% Dimension Turf Herbicide 14-0-5"

EPA Reg. No. 9198-117

Application Dated April 20, 2009

Dear Ms. Ziehr:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 04/20/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

Places and instructions		dina da		C 4		40 N	70 0000	Ammuourd armiras 8 20 0
SEPA	Environmenta:	Inited States			Re Ar	gistrati nendmo her	on	O. Approval expires 2-28-9 OPP Identifier Number
		Applicatio	n for Pestici	de - Section	on I			
1. Company/Product Number 9198-117			2. EPA Miller	Product Manag	er		3. Pro	posed Classification
4. Company/Product (Name) The Andersons Dimension H			PM# 23]	, notine
5. Name and Address of App The Andersons Lawn Fert d/b/a Free Flow Fert. P.O. Box 119 Maumee, OH 43537		ode)	(b)(i), n to: EPA F	ny product is	similar	or identica	al in cor	FIFRA Section 3(c)(3) mposition and labeling FICATION 2 1 2009
			Section -	——— <u> </u>			IVI MI	<u>4 1 7009</u>
Amendment - Explain Resubmission in resp Notification - Explain Explanation: Use addition Notification of Alternate Brand Nat This notification is consistent with made to the labeling or the confide further understand that if this notification of FIFRA and I may be set	below. nel page(s) if necessar me per PR Notice 98-10: the guidance of PR Notice ential statement of formula ication is not consistent wi	Ty. (For section The Andersons To 98-10 and the red of this product. I th the guidance of	urf Products Fertilizer of quirements of EPA's re understand that it is a PR Notice 98-10 and	gulations at 40 Cl violation of 18 U.S the requirements	r dated oplication in below asion Turf F	Herbicide 14-	0-5 152.46 ai	ny false statement to EPA. I
	· 		Section - I	<u> </u>				
1. Material This Product Will Child-Resistant Packaging Yes No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per container	Water Soluble P Yes ✓ No If "Yes" Package wgt	Ackaging No. per container	2.	<i>J</i>	onteiner Metal Plastic Glass Paper Other (S	pecify)
3. Location of Net Contents Lebel ✓ C 6. Manner in Which Label is	ontainer	4. Size(s) Reta	10 - 2,000 lbs.	√ Other		on of Label On Label	Directio	ns
			Section - I'	V				
Contact Point //Complete Name Debbie Ziehr	items directly below t	T	n of individual to b Title Regulatory Admir		necessa	T		No. (Include Area Code)
I certify that the stater I acknowledge that an both under applicable	y knowlinglly false or	Certifica this form and	tion all attachments the	ereto are true,		and comp	ilete.	6. Date Application Regeived' ((Stamped)
2. Signature	Ziehr	:	3. Title Regulatory Adminis	strator			,	9937 7 9 9

5. Date

4/20/09

4. Typed Name

Debbie Ziehr



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

April 20, 2009

Document Processing Desk (NOTIF)
Office of Pesticide Programs - (7504P)
U. S. Environment Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Subject:

NOTIFICATION - Additional Brand Name

RE:

The Andersons Dimension Herbicide IV

EPA Reg. No. 9198-117

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons Turf Products Fertilizer with 0.25% Dimension Turf Herbicide 14-0-5

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

Debbie Ziehr

Regulatory Administrator Phone #419-891-6671

Fax #419-891-2745

Email: Debbie_ziehr@andersonsinc.com

Enclosure



Andersons Turf Products

Fertilizer with 0.25% Dimension® Turf Herbicide 14-0-5

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

GUARANTEED ANALYSIS Total Nitrogen (N) 14.00%

14.00% Urea Nitrogen* Soluble Potash (K ₂ O)	5.00%
Plant nutrients derived from urea, potassium chloride and potated sulfur coated urea.	olymer
Chlorine (CI) Max	4.24% ur coate

ACTIVE INGREDIENT:	
Dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-	
Dittiopyl. 3,3-pylicitiedicardoution acid, 2-(minorotteatyl)-4-	0.000
(2-methylpropyl)-6-(trifluoromethyl)-5,5-dimethyl ester	0.25%
(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester	99,75%
Total	100.00%

EPA Reg. No. 9198-117 EPA Est. No. 9198-OH-1<u>M.</u> 9198-OH-2<u>B.</u> 9198-AL-001<u>A</u> (Underlined letter is first letter used in run code on end of bag.)

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation, Harmful if inhaled, Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughtly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact white handling litis material may cause allergic reaction in some individuals.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, it present, after the first 5 minutes, then continue rinsing. Call a poison control center or clocker for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not include oranizing unless loid to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth it possible. Call a poison control center or obcor for freatherst activice.

Have the product container or label with you when cating a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including systems and shimin. Use with care when splying to furt areas actacent to any body of water. Dritt and runoff from treaded furt may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertisal areas below the mean high water mark. Do not apply when weather conditions tava "artiful time treated areas. Do not contaminate water when disposing of equipment washington."

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on truth being using this product, Use only according to label instructions. Not for use on truth being useful and use as od, or for commercial seed production, or for massively product.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more highly typoungs ppr 1,000 square feet (equal to 0.5 pounds active ingredient) per acrol braden; and some highly programs in Nassau and Suffolk Counties.

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turts, including golf course fairways, roughs, and too boxes. This product will not harm nearby established conamentals when used according to label directions.

LISE PRECAUTIONS

Turigrass Safety: This product may be used on seeded, sockled, or sprigged lawns and omamental turis that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two movings tollowing its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and omamental turis that are not well-established, or on those that have been weakened by weather, peat, disease, chemical-, or mechanical-related stress, may increase the chances for furf injury.

This product should only be applied to lawns and ornamental turts that are composed of the following turigrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turigrasses are tolerant to this product:

Cool-Season Grasses
Bentgrass, creeping "Agrostis palustris)
Bluegrass, Kentucky (Poa pratensis)
Fescue, tine" (Festura nubra)
Fescue, tall (Festura arundra)
Ryegrass, peremial (Jollium peremie)

F292

Warm-Season Grassas Bahagrass (Paspatum notatum) Bermudagrass (Cyrnodro dactylon) Buttalograss (Wachibe dactyloties) Carpalgrass (Axonopus attinophiluroides) Kituyugrass (Fernisetum Cardestrium) SI. Augustinegrass (Stenotaptrium secundatum) Zoyagrafass (Zoyaia japoman)

"Use of this product on certain varieties of creeping bentgrass, such as "Cohansey", "Carrieri," Seaside, and "Washington" may result in undesirable turtgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (Agrostis tenuis) varieties.

"Use of this product on cortain varieties of line fescue may result in undestable furt injury. The following fine fuscue varieties have been found to be sensitive to this product: 'Allanta', 'Banner,' Beauty, 'Bigart', CF-2', Enjoy', 'HF-93', 'Highlight', 'Nalo', 'Jamssfown', 'Koket', 'Majerita', 'Mary', 'Pennlawn', 'Tamara', 'Tatjaras', 'Wakfort' and 'Wakfora', 'Not all varieties of line fescue have been fested.

""OO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turtgrass is fully green and has established new roots.

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

Reseabling, Overseading, or Sprigging: Reseabling, overseading, or sprigging of treated a reas within a months after application of this product may inhibit the establishment of desirable turigrasses. When reseabing or overseading, proper cultural practices such as soil cultivation, irrigation and tertitization about the follower. For heat results, use mechanical or power seading equipment (six seeders) designed to give good seed to soil contact.

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass it applied before crabgrass has displayed its fitth leaf or first titler.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydroaertication, and verticuting, should be done before application of this product. DO NOT apply altion product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turts only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

Control of Crabgrass

Premiergence and Early Postemergence Control: This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species when applied prior to the emergence of crabgrass from the ground in established lawns and omamental turts.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very arrail, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass pitch generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) if can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 4.6 lbs. per 1,000 sq. ft. per application, and no more than 13.8 Lbs. per 1,000 sq. ft. per year,

Spring Applications: For single applications made in the spring or early surrows, this product should be applied at the appropriate rate corresponding to one of the three control programs lated in Table 1 helow, depending on the user's location, the turtgrass moving height, and whether the use is considered to be preemergence or early postermergence at the time of the application. The duration or resolutal weed control provided by this product is directly related to the total rate applied, but will vary somewhall depending on wealther, weed pressure, turtgrass competitiveness, and the user's location within a renion.

Use Program 1 for preemergence control at sites where the turigrass is cut high (e.g., homeowner lawns). This program provides 3.5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turtigrass is out low (e.g., golf atmways) and (b) turtigrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence catignass control. This program may also be used for early postemergence control at sites where turtigrass is cut high (e.g., homeowner farms).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf lairways), and (b) fur maintlenance or weed control has not been conducted during the previous year. This program provides 4.6 months of preemergence carbgrass control. This program may also be used for early postemergence control at sites where furfgrass is cut low (e.g., oolt fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split lentilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to craburass emergence.

PROGRAM USE (turigrass cut)	11	2	Proamergence (low)	
	Preemergence (high)	Preemergence (low)		
REGION		Postemergence (high)	Postemergence (low)	
North	12	1.7	2.3	
Transition	1.7	2.3	34	
South	23	3.4	4.6**	
Coastal South	34	4.6"	4.5	

"DO NOT apply more than 4.6 lbs. per application, and no more than 13.8 lbs. per year. Postemergence control is limited, see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.
Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

"May require split or sequential applications for full control.

Fall Applications: This product can also be applied in the late summer or early fall (fale August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-from control.

Tips for Improved Control: For best results, apply this product within a lew days after nowing and delay mowing again for a lew days after the application. When treated lawn or crnamental furl areas are watered or receive significant rainfall within a lew days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barryardyrass, bluegrass (annual, roughstalk), crowfootgrass, lostall (yellow & green)", goosgass, kikuyugrass, ryograss (annual & perinnial), smrligrass, lepedera (common), crais (full crup, creeping & yellow woodsorrus), pursiane (common), speedwell (com), spurge (prostrate & spoting).

"Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for tate summer or early fall use, for crabgrass control incugin the early part of the next spring, this product will also control (at the Program 3' rates) or suppress (at the "Program 2' rates) the following weeds when applied prior to their emergence; bluegrass (annual): Poa annua, bitforcress, chickweed, princaptieweed, herbit, greatnum (Cacrinia), parsley prier; hepherdepure.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with dop or rotary type spreaders designed to apply granular her bicities. Avoid her see of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental furfures. A more uniform application can be made by sureading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, steps, or overlaps during application. Check exaginment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

SUGGESTED SPREADER SETTINGS

These suggested spreader settings are not intended to replace calchation.

Plans calchat spreader settings are not intended to replace calchation.

50 lbs. covers 21,700 sq. 11, (0.53 acre) at the 100 lbs. productares the EDUM RATE - 2.3 lbs.1, 0.00 sq. 11.

50 lbs. covers 14,700 sq. 11, (0.53 acre) at the 152 lbs. productares the HAT RATE - 3.6 lbs.1,000 sq. 11.

50 lbs. covers 10,500 sq. 11, (1.55 acre) at the 152 lbs. productares the HAT RATE - 4.6 bs.1,000 sq. 11.

			SPREADER SETTINGS			
SPREADER	GRC IND SPEED.	WIDTH OF COVERAGE	MEL-A R. E 0.2 b. avacre	FATE O 0.38 lb	TAYE 150 lb.	
Andersons AccuPro 2000, SR-2000 (cone 4)	3 mph	11 feet	К	L	M 1/4	
Scotts R-B, R-BA (cone 4)	3 mph	11 feet	1 1/2	Ķ	L 1/4	
Lesco 020093 (pattern 0 25)	3 mph	~ - 10 fred -~	- ~.€	- F.1/2	G 3/4	
Lesco 705699 (patiem 0.25)	3 mph	10 feet	, Ç	15	17 V2	
Lety Models WTR, WFR, HR, 1250 (PTO at 450 mph) Pattern RI-B	ار ار 4.5 mph	32 feet	4 1/2	s Ve	6 1/2	
Vicon (03 Series)	4.5 mph	36 test	21	24	28	

STORAGE AND DISPOSAL

Do not contaminate water, lood, or leed by storage and disposal

STORAGE: Keep container closed and away from food, feedstuffs, and domestic water supplies.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Fodoral Law. It these wastes cannot be disposed of by use according to laber instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office

CONTAINER DISPOSAL: Completely empty bag into application equipment. Nonrefilable container. Do not reuse or refill this container. Ofter for recycling, if available. In oit available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of recycle.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the table when used in stitle accordance with the directions, Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERICHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LUBBLITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to retund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATIORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY. TO LABEL DIRECTIONS.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS, YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATEJURISDICTION TO STATEJURISDICTION

Product of USA

Manufactured by: The Andersons Lawn Fertilizer Division, Inc. P.O. Box 119 Maumee, OH 43537

The Andersons logo is a registered trademark of The Andersons, Inc.; Damension is a registered trademark of Dow AgroSciences LLC.

SGN 215 Net Wt. 50 Lbs. (22.68 kg)



