

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

JUN 4 2009

Mrs. Debbie Ziehr Regulatory Administrator The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow Fert. P.O. Box 119 Maumee, OH 43537

RE: Notification of Alternate Brand Names: The Andersons Turf Products Fertilizer with 0.25% Dimension Turf Herbicide 21-0-10

Registration Number: 9198-117 Submission Dated: March 31, 2009

Dear Mrs. Ziehr:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated, March 31, 2009, for the product, The Andersons Dimension Herbicided IV. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely yours,

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

Please read instructions on reverse before completing	ng form.		Form Ap	proved.	OMB No.	2070-006	O. Approval expires 2-28-9		
United States Environmental Protection Washington, DC 2046		on Agency			Registration Amendment ✓ Other		OPP Identifier Number		
A	pplication	n for Pesti	cide - Sec	tion I					
1. Company/Product Number 9198-117		2. EPA Product Manager Miller				3. Proposed Classification None Restricted			
4. Company/Product (Name) The Andersons Dimension Herbicide IV		PM# 23					Nosaliciae		
5. Name and Address of Applicant (Include ZIP Code The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow Fert. P.O. Box 119 Maumee, OH 43537 Check if this is a new address		(b)(i) to: EPA		is simil	ar or ident	tical in co	FIFRA Section 3(c)(3) mposition and labeling FICATION 0 4 2009		
		Section	- 11						
Amendment - Explain below. Resubmission in response to Agency letter do Notification - Explain below.	ated	Final printed labels in repsonse to Agency letter dated "Me Too" Application. Other - Explain below.							
Notification of Alternate Brand Name per PR Notice 98-10: The This notification is consistent with the guidance of PR Notice 98 made to the labeling or the confidential statement of formula of further understand that if this notification is not consistent with the violation of FIFRA and I may be subject to enforcement action as	3-10 and the re this product. I the guidance of	quirements of EPA's understand that it is FPR Notice 98-10 ar	regulations at 40 a violation of 18 nd the requiremen) CFR 156 U.S.C. Se	6.10 and 40 C cc. 1001 to wil	FR 152.46 a lifully make a	iny false statement to EPA. I		
		Section -	[]]						
1. Material This Product Will Be Packaged In:									
Child-Resistant Packaging Yes ✓ No * Certification must be submitted Unit Packaging Yes ✓ No If "Yes" Unit Packaging wgt. No. per container		Water Soluble Packaging Yes ✓ No If "Yes" Package wgt No. per Package wgt No. per				of Container Metal Plastic Glass Paper Other (Specify)			
	. Size(s) Ret	ail Container	<u> </u>	5. Loca	ation of Lat	oel Directio	ons		
Label V Container		10 - 2,000 lbs.		[-	On Labe	el			
6. Manner in Which Label is Affixed to Product V									
<u>, , , , , , , , , , , , , , , , , , , </u>		Section -	IV				(
1. Contact Point (Complete items directly below for	identificatio	n of individual to	be contacted,	, if nece:	ssary, to pr	ocess this			
Name Debbie Ziehr		Title Regulatory Administrator				Telephone No. (Include Area Code)			
f certify that the statements I have made on the lacknowledge that any knowlingfly false or mit both under applicable law.		all attachments			ate and co		6. Date Application Received (Stamped)		
2. Signature Lebbie Ziekr		3. Title Regulatory Administrator							
4. Typed Name Debbie Ziehr		5. Date 3/31/09							



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

March 31, 2009

Document Processing Desk (NOTIF)
Office of Pesticide Programs - (7504P)
U. S. Environment Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Subject:

NOTIFICATION - Additional Brand Name

RE:

The Andersons Dimension Herbicide IV

EPA Reg. No. 9198-117

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons Turf Products Fertilizer with 0.25% Dimension Turf Herbicide 21-0-10

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

Debbie Ziehr

Regulatory Administrator

Phone #419-891-6671

Fax #419-891-2745

Email: Debbie_ziehr@andersonsinc.com

Enclosure



Fertilizer with 0.25% Dimension® Turf Herbicide 21-0-10

Contains NutriSphere-N®

This product is a herbicide that provided the product on laws and ornamental turls that are not well-established, or on those annual grasses and broadleaf weeds the stablished away that have been weakened by weather, pest, disease, chemical, or mechanical-related ornamental turfs.

JUN 0 4 2009

GUARANTEED ANALYSIS

21.00% Urea Nitrogen Soluble Potash (K ₂ O)	10.00%
Plant nutrients derived from urea and potassium chloride. Chlorine (CI) Max	8.46%
	F292
ACTIVE INGREDIENT: Dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4- (2-methylpropyl)-6-(trifluoromethyl)-5,5-dimethyl ester INERT INGREDIENTS: Total	0.25% 99.75%

EFA Est. No. 9198-OH-1M. 9198-OH-2B. 9198-AL-001A (Underlined letter is first letter used in run code on end of bag.)

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing, Avoid breathing dust. Wash thoroughly with scop and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If In eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes, Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including cysters and shrimp. Use with care when applying to furl areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Bead the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research nurposes.

New York Use Restrictions: In New York State this product may paly be used by professional applicators and at no more than 4.6 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties

GENERAL INCORMATION

This product is a herbicide that provides control of craborass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee hoves. This product will not horm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turforess Safety: This product may be used an seeded, sodded, or springed lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seedig, sodding, or sprigging before it can receive its first application of this product. Use stress, may increase the chances for furf injury

This product should only be applied to lawns and ornamental turfs that are composed of the following turigrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turforasses are tolerant to this product:

Cool-Season Grasses Bentgrass, creeping" (Agrostis palustris) Riueursss, Kentucky (Poa pratensis) Ryegrass perennial (I olium perenne)

Warm-Season Grasses Bahiagrass (Paspalum notatum) Carting ass (Fashaum rotatum)
Buffalograss (Cynodon dactylon)
Buffalograss*** (Buchloe dactyloides)
Carpetgrass (Axonopus affinis)
Centipedegrass (Eremochloa ophiuroides) Kikuyugrass (Pennisetum clandestinum) St. Augustinegrass (Stenotaphrum sec Zoysiagrass (Zoysia japonica)

"Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', "Carmen" "Seaside" and 'Washington' may result in undesirable turforass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Benturass (Agrostis tenuis) varieties

"Use of this product on certain varieties of fine feacue may result in undesirable turi Injury. The following fine feacue varieties have been found to be sensitive to this product: Adanta, Beauty, Bligart, CF-2, 'Enjoy, 'HF-93, 'Highight,' Ivab', Jamestown, Koket, 'Majerta, 'Mary, 'Pennlawn, 'Tamara', 'Tajana', 'Waldorf and 'Waldina'. Not all varieties of fine fescue have been tested

***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results. use mechanical or power seeding equipment (slit seeders) designed to give good seed

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on craborass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydroaerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turks paly. DO NOT apply

DO NOT use clippings from treated turf for multihing around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled

APPLICATION DIRECTIONS

Control of Crabgrass Preemergence and Early Postemergence Control: This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 4.6 lbs, per 1,000 sq. ft, per application, and no more than 13.8 Lbs. per 1,000 sq. ft, per year

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control guarants listed in Table 1 below depending on the user's location, the turburass mowing height, and whether the use is considered to be preemergence or early postamergence at the time of the application. The digration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turtorass competitiveness, and the user's

Use Program 1 for preemergence control at sites where the turforass is cut high (e.g. ~ homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence graborass control. --

Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemgrance grandrass control This program may also be used for early postemergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemargence applications should be made where known periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

PROGRAM	1 1	2	3
USE (turigrass cut)	Preamergence (high)	Preemergence (low)	Preemergence (low)
REGION		Postemergance (high)	Postsmergence (low)
North	12	1.7	2.3
Transition	1.7	2.3	3.4
South	2.3	3.4	4.6**
Coastal South	3.4	4.6**	4.6**

*DO NOT apply more than 4.6 lbs, per application, and no more than 13.8 lbs. per year. Postemergence control is limited; see "Craburass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other area

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL. IN, OH, & coastal areas of CT, NY, & RI.
South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX,

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX,

"May require split or sequential applications for full control

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next apring. The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barnyardgrass, bluegrass (annual, roughstalk), crowfootgrass, foxtail (yellow & green)*, goosegrass, kikuyugrass, ryegrass (annual & perennial), smutgrass, lespedeza (common), oxalis (buttercup creeping & yellow woodsorrel), pursiane (common), speedwell (corn), spurge (prostrate

*Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for late summer or early fall use, for craborase control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: bluegrass (annual) - Poa annua, bittercress, chickweed, pineappleweed, henbit, geranium (Carolina), paraley-piert, shepherdspurse,

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular horizontals. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, cathorate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking skins, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules,

CHOCCETED CODE A DED CETTIMOS

SUGGESTED SPREADER STITMGS
These suggested scenario et alimps are not internded to replace calibration.
Places calibrate your spreader before applying product.
50 lbs. covers 23,400 sq. ft. (0.57 scenar) at the 74 be productabers (DM PATE - 1.7 lbs/1,000 sq. ft.
50 lbs. covers 21,700 sq. ft. (0.50 scray) at the 100 lbs. productabers (MEDUM PATE - 2.3 lbs/1,000 sq. ft.
50 lbs. covers 11,700 sq. ft. (0.33 scray) at the 100 lbs. productabers HECHM PATE - 3.4 lbs/1,000 sq. ft.
50 lbs. covers 11,000 cq. ft. (0.33 scray) at the 200 lbs. productabers HEAY PATE - 4.5 lbs/1,000 sq. ft.

	2 22		1	~ -	¬, SP	TINGS	
^	C TREADER	GROWND SPLED	WIDTH COVERAGE	COW TATE (-19 lb. avacre	MEUTUM TE 0-1 lb, akacre	HIGH RATE 0.38 lb. sl/acre	HEAVY RATE 0,50 lb. ai/acre
	Andersons AccuPro 2000, SR-2000 (core 4)	3 mph	11 leet	J 1/2	K 1/4	, t 1/4	M 1/2
١	Andersons SS-2 Orop	3 mph	overlap wheels	2.1/2	2 3/4	3 3/4	5
1	Scotta R-8, R-8A (cops_4)	_ 3,mph	11 feet	1	13/4	K 1/2	L 1/4
1	Lesco 020090 (par 7-m 0.25)	3 nC3	11Clast	บาล	ŧ	FHA	G 172
Ì	Lesce 705698 (pattern U.25)	3 mph	10 feet	10 1/2	12	14 1/2	17
	Lely Models WYR. WFR HR, 1250 — — (PTO at 450 mph) - datern #-A	-, 45 mph	C 4C 381	4 1/2	5	5 3/4	6 1/2
Ì	Vicon (03 Series)	4.5 mph	3E leet	20	23	. 28	34

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs, and domestic

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty had into application equipment Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke

IMPORTANT: READ REFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability

WADDAMTY DISCHAIMED and LIMITATION of LIABILITY.

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTARILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY-LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS, YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATEJURISDICTION TO STATEJURISDICTION

Manufactured by: The Andersons Lawn Fertilizer Division, Inc. PO Box 119 Maumee, OH 43537 www.andersonslawnnmducts.com

The Andersons logo is a registered trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences LLC; NutriSphere is a registered trademark of Specialty Fertilizer Products, LLC.

> **SGN 215** Net Wt. 50 Lbs. (22.68 kg)



