



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

JAN 8 2010

Ms. Debbie Ziehr Product Registration The Andersons Lawn Fert. Div., Inc PO Box 119 Maumee, OH 43537

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request Alternate Brand Name "The Andersons Turf Products Fertilizer with

0.25% Dimension Turf Herbicide 12-4-8"

EPA Reg. No. 9198-117

Application Dated November 10, 2009

Dear Ms. Ziehr:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 11/10/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

Please read instructions on I	reverse before com	ing form.		Form A	pproved	.∷OMB No.	2070-006	O. Approval expires 2-2		
United States Environmental Protection Washington, DC 20460				Regis Agency Amer		Registr Amend Other		OPP Identifier Number		
		Application	for Pesi	ticide - Se	ction	J		. 		
1. Company/Product Number 9198-117			2. EPA Product Manager Miller			3. Pr	3. Proposed Classification None Restricted			
Company/Product (Name) The Andersons Dithiopyr Herbicide IV			PM# 23							
The Andersons Lawn Fert d/b/a Free Flow Fert. P.O. Box 119 Maumee, OH 43537		de)	(b)(to:	i), my produc	t is sim	ilar or iden	itical in co	FIFRA Section 3(c)(3 mposition and labeling		
Check if this	is a new address		Pr	oduct Name						
			Section	- 11						
Amendment - Explain Resubmission in resp Notification - Explain Explanation: Use addition Notification of Alternate Brand Na This notification is consistent with made to the labeling or the confide further understand that if this notification of FIFRA and I may be so	below. hal page(s) if necessary me per PR Notice 98-10: the guidance of PR Notice 98-10 in the guidance of promula of the page of the pa	. (For section The Andersons Tu 98-10 and the requ of this product. I u the guidance of F	orf Products Ferti uirements of EP/ anderstand that it PR Notice 98-10	"Me Too" Other - Est II.) ilizer with 0.25% E A's regulations at is a violation of 1 and the requirement.	* Applica explain be Dimension 40 CFR 18 8 U.S.C. 8	ation. Siow. Turf Herbicide 56.10 and 40 (Sec. 1001 to w	JAN 12-4-8 CFR 152.46 a illfully make a	ny false statement to EPA. I		
	·		Section	- 111	· · · · · · · · · · · · · · · · · · ·			· · · · · · · · · · · · · · · · · · ·		
1. Material This Product Will	Be Packaged In:		A Suffer of a Maria				<u> </u>			
Child-Resistant Packaging Yes V No Certification must be submitted	Unit Packaging Yes ✓ No If "Yes" Unit Packaging wgt.	No. per container	Water Solut Yes No If "Yes" Package wg	No. pe		2. Type of	Metal Plastic Glass Paper Other (S	pecify)		
3. Location of Net Contents	Information container	4. Size(s) Reta	il Container 15 - 50 lbs.		5. Lo	cation of La		ns		
6. Manner in Which Label is	Affixed to Product	Lithogra Paper g Stencile	aph lued ed	✓ Oth	ner Po	ocket				
			Section	- IV						
1. Contact Point Complete	items directly below fo	r identification	of individual	to be contacte	d, if nec	essary, to p	rocess this	application.)		
Name Debbie Ziehr			Title Regulatory Administrator			Telephone No. (Include Area Code) 419-891-6671				
	ments I have made on t y knowlinglly false or n		il attachment					6. Date Application Pectived (Stamped)		

3. Title

5. Date

Regulatory Administrator

11/10/09

2. Signature

4. Typed Name

Debbie Ziehr



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

November 10, 2009

Document Processing Desk (NOTIF)
Office of Pesticide Programs - (7504P)
U. S. Environment Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Subject:

NOTIFICATION - Additional Brand Name

RE:

The Andersons Dithiopyr Herbicide IV

EPA Reg. No. 9198-117

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons Turf Products Fertilizer with 0.25% Dimension Turf Herbicide 12-4-8

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

Debbie Ziehr

Regulatory Administrator Phone #419-891-6671

Fax #419-891-2745

Email: Debbie_ziehr@andersonsinc.com

Enclosure



Andersons Turf Products

Fertilizer with 0.25% Dimension® Turf Herbicide 12-4-8

This product is a herbicide that provides the broadleaf weeds in established lawns and the transfer of the broadleaf weeds in established lawns and the broadleaf weeds in established lawns and the broadleaf weeds in established lawns are the broadleaf weeds are the broa

JAN - 8 2010

GUARANTEED ANALYSIS

Tr' 'Mitrogen (N)	12.00%
a Phosphate (P2O5)	4.00%
Dotash (K ₂ O).	8.00% 14.60%
14.60% Combined Sulfur (S)	

Plant nutrients derived from ammonium phosphate, ammonium suttate and potassium

FLORIDA APPLICATIONS: We recommend that you follow the Florida Green industries BMP's at http://www.flaes.org/pdt/BMP_Book_tinal.pdf

FLORIDA APPLICATIONS: We recommend that you follow the Florida Golf Course BMP's at http://www.tlaes.org/pdf/uffbmp07.pdf

ACTIVE INGREDIENT: Dithiopyr: 3,5 syndimedicarbothioic acid, 2-(difluoromethyl)-4- (2-methylpropyl)-8-(trisfluoromethyl)-S,S-dimethyl ester MERT INGREDIENTS:	0.25% 99.75%
Total	100.00%

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation, Harmful fi inhaled. Avoid contact with eyes or clothing, Avoid ing clust. Wash thoroughly with soap and water after handling. Remove aled clothing and wash before euce. Prolonged or I requently repeated skin ii ib handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person

If on skin or clothing: Take of contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice

If Inhaled: Move person to tresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth it possible. Call a poison control center or chotor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment

ENVIRONMENTAL HAZARDS

This product is toxic to lish and bighly toxic to other aquatic organisms including cysters and shrimp. Use with care when applying to turf areas adjacent to any body of and runoff from treated but may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions tayor drill from treated areas. Do not contaminate water when disposing of equipment

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to tabel instructions. Not for use on furf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 4.6 pounds por 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleal weeds in established lawns and ornamental turts, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions

USE PRECAUTIONS

Turigrass Safety: This product may be used on seeded, sodded, or sprigged lawns and ornamental turis that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings tollowing its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turts that are not well-established, or on those that have been weakened by weather, pest, disease, chemical, or mechanical related stress, may increase the chances for furl injury.

This product should only be applied to lawns and ornamental turks that are composed of the following turigrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following shed turiorasses are tolerant to this product

Cool-Season Grasses Bentgrass, creeding (Agrostis palustris) Bluegrass, Kentucky (Poa pratensis)

Fescue tine" (Festuca rubra) Fescue, tali (Festuca arundinacea) Ryegrass, perennial (Lolium perenne) Werm-Season Grasses

werm-season crasses
Bahagrass (Paspairm notatum)
Bermudagrass (Cymodon dacyfoides)
Butlalograss ("Buchloe dacyfoides)
Carpelgrass (Avonçous atlinis)
Centpedograss (Fernnisetum clandestrum)
Kikuvugrass (Pennisetum clandestrum)
SI Aurustinienrass (Stonortschalm serverta St. Augustinegrass (Stenotaphrum secundatum) Zoysiagrass (Zoysia japonica)

"Use of this product on certain varieties of creening hentgrass, such as "Cohansev" 'Carmon', 'Seaside', and 'Washington' may result in undestrable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (Agrostis tenuis) varieties.

"Use of this product on certain varieties of fine fescue may result in undestrable turl injury. The following fine feacue varieties have been found to be sensitive to this product: 'Allanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', Jamestown', 'Kokel', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Wakdorf' and 'Wakdina', Not all varieties of tine lescue have been tested.

""DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turigrass is fully green and has established new roots

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of freated areas within 4 months after application of this product may inhibit the establish-ment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, singation and tertitization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabonass if applied before craborass has displayed its fifth leaf or first title

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turks only. DO NOT apply to flowers vegetables shrubs or trees

DO NOT use clippings from treated turf for mulching around vegetables or truit trees Keep people and pols off treated areas until dust has settled.

APPLICATION DIRECTIONS

Control of Crabgrass Preemergence and Early Postemergence Control: This product provides rgence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns

It can also provide "early posternergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is on difficult to see the very small, early stages of crabgrass in well-established to and ornamental turis. Postemorgence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the graborass plants in the lawn or turl. So the practical benefit of this product's additional, early posternergence activity is that compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 4.6 lbs, per 1,000 sq. (I, per application, and no more than 13.8 Lbs. per 1,000 sq. fl. per year.

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turtgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turtgrass competitiveness, and the user's

Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turbrass is cut low (e.g., golf, fairways) and (b) furigrass maintenance or weed control has been concluded during the previous year. This program provides 4-6 months of preomergence crabgrass control. This program may also be used for early postemergence control at sites where turigrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turigrass is cut low (e.g., golf fairways), and (b) furf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control This program may also be used for early posternergence control at sites where turigrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emerge

TABLE 1: Recommended Application Use Rates* (Lbs. of this product/1,000 sq. It.)					
PROGRAM	1	2	3		
USE (turigrass cut) REGION	Preemergence (high)	Preemergence (low) Postemergence (high)	Preemergence (low) Postemergence (low)		
North	12	1.7	2.3		
Transition	1.7	23	34		
South	23	34	4.6"		
Coastal South	3,4	4.6**	4.6"		

*DO NOT apply more than 4.6 lbs, per application, and no more than 13.6 lbs, per year rgence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below

North: All other areas

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.
South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

"May require split or sequential applications for full control

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control: For best results, apply this product within a lew days after mowing and delay mowing again for a few days after the application. When treated lawn or organization and areas are watered or receive significant rainfall within a lew days after lication of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for craborass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barnyardgrass, bluegrass (annual, roughstalk), crowlootgrass, loxiail (yellow & green)", goosegrass, kikuyugrass, ryegrass (annual & perennial), smutgrass, lespedeza (common), oxalis (buttercup) creeping & yellow woodsorret), pursiane (common), speedwell (com), spurge (prostrate & spotted).

'Also controlled at the "Program 2" rates

Fall Applications: Used as directed for late summer or early fail use, for crabonass control through the early part of the next spring, this product will also control (at the Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: bluegrass (annual) - Poa annua, bittercress, chickweed, pineappleweed, henbit, geranium (Carolina), parsley-piert, shepherdspurse.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granutar herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or organization area. A more uniform application can e made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking skins or overtaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

SUGGESTED SPREADER SETTINGS

These suggested greater settings are not intended to replace calibration.

These suggested greater settings are not intended to replace calibration.

Solibs, covers 29,400 sq. ft, 607 and get that 7 like productions to PARE - 17 lbs./i p00 sq. ft.

Solibs, covers 29,400 sq. ft, 605 down) at the 100 lbs, productions to PARE - 17 lbs./i p00 sq. ft.

Solibs, covers 17,700 sq. ft, 625 down) at the 100 lbs, productions to PARE - 1.4 lbs./i p00 sq. ft.

Solibs, covers 10,800 sq. ft, 625 acre) at the 200 lbs, productions the RAY RATE - 4.6 lbs./i,000 sq. ft.

		1		SPREAMER STYTING			
SPREADER	GROUM) \$1 .E0	WIDTH OF COVERAGE	LOW RATT 0.25 Ib all'acte	MEDIUS. RATE 025 (b. sl'acre	HDH. 'ATE C 18 lb. erseve	RAT 0.50 lb. allson	
Andersons AcquPro 2000, SR-2000 (con+6)	3 mph	10 feet	JI#	E	L 1/4	14 1/4	
Souts R-8, R-8A (cone 6)	3 mph	10 feet	H 3/4	13/4	J 3/4	K 3/4	
Andersons SS-2 Drop	3 mph	overlap wheels	?	217	3 144	4	
Lesco 020093 (pattern 0.25)	3 mph	91eet	ē	D	E ·	E V2	
Lesco 705695 (pattern 0.25)	3 mph	, bled l	A	10	12	13	
Lely Models WTR, WFR, HR, 1250 (PTO at 450 mph) Pattern III-B	45 mph	1 3010-1 1 3010-1	4	. 1/4	5	53/4	
Vicon (03 Serias)	4.5 mph	32 feet	12	15	18	21	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Keep container closed and away from tood, teedstuffs, and domestic vater supplies

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for

CONTAINER DISPOSAL: Completely empty hag into application equipment.

Nonrelitlable container. Do not reuse or refit this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability fore using this product. If terms are not acceptable, return the unoponed product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably lift for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyor's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LUBBLITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL BIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS: YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATEJURISDICTION TO STATEJURISDICTION.

Manufactured by: The Andersons Lawn Fertilizer Division Inc. Maumee, OH 43537 www.andersonstawnproducts.com

The Andersons logo is a registered trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences LLC

> **SGN 150** Net Wt. 50 Lbs. (22.68 kg)



* : * * /七