

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

MAR 2 1 2008

Ms. Dawn Walters Product Registration The Andersons Lawn Fert. Div., Inc PO Box 119 Maumee, OH 43537

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request Alternate Brand Name "The Andersons Turf Products Fertilizer with

0.072% Dimension Turf Herbicide"

EPA Reg. No. 9198-118

Application Dated February 28, 2008

### Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 02/25/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

			2084
Please read instructions on reverse before completing form.  United States Environmental Protecti Washington, DC 20	on Agency	Registration Amendmen  Other	
Application	on for Pesticide - Sec	ction I	
1. Company/Product Number 9198-118	2. EPA Product Ma Kraft	nager	3. Proposed Classification  None Restricted
4. Company/Product (Name) The Andersons Dithiopyr Herbicide I	<b>PM#</b> 23		Nastricted
5. Name and Address of Applicant (Include ZIP Code)  The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow IPO Box 119  Maumee, OH 43537  Check if this is a new address	Fert. (b)(i), my product	t is similar or identical i	with FIFRA Section 3(c)(3) n composition and labeling
	Section - II		
Amendment - Explain below.  Resubmission in response to Agency letter dated  Notification - Explain below.  Explanation: Use additional page(s) if necessary. (For section Notification of Alternate Brand Name per PR Notice 98-10: The Andersons This notification is consistent with the guidance of PR Notice 98-10 and the made to the labeling or the confidential statement of formula of this product. Further understand that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this notification is not consistent with the guidance of the second that if this product.	Agency le "Me Too" Other - Ex On I and Section II.) Turf Products Fertilizer with 0.072% If requirements of EPA's regulations at 4 I understand that it is a violation of 18	10 CFR 156.10 and 40 CFR 15 3 U.S.C. Sec. 1001 to willfully n	nake any false statement to EPA. I
violation of FIFRA and I may be subject to enforcement action and penalties		5/13 0/ 40 OF IT 130. TO allo 40 V	or trioz.40, this product may be in
Material This Product Will Be Packaged In:			
Child-Resistant Packaging  Yes  No  No  * Certification must be submitted  Unit Packaging  Yes  No. per Container	Water Soluble Packaging  Yes  No  If "Yes"  Package wgt  Contain	Gla Pal	stal estic
3. Location of Net Contents Information 4. Size(s) Re	stail Container	5. Location of Label Di	rections
	graph Oth	ner	cocc
	Section - IV		
1. Contact Point (Complete items directly below for identification	on of individual to be contacted	d, if necessary, to process	s this application.)
Name Dawn L. Walters	Title Regulatory Administrator		phone No. (Include Area Code) 8897-6790° ေ
Certification is certify that the statements I have made on this form and acknowledge that any knowlingly false or misleading state both under applicable law.	d all attachments thereto are tr	τας σε ue, accurate and οςτόρΙσί fine or imprisonment or	E. Eate Application Received  Cocc (Stamped)
2. Signature	3. Title		( ( (
Dawn Xwalter	Regulatory Administrator		- ( ( )
4. Typed Name Dawn L. Walters	5. Date 2/25/0	g	



## The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

February 26, 2008

Document Processing Desk (NOTIF) Office of Pesticide Programs - (7504P) U. S. Environment Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

Subject:

**NOTIFICATION - Additional Brand Name** 

RE:

The Andersons Dithopyr Herbicidel

EPA Reg. No. 9198-118

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons Turf Products Fertilizer with 0.072% Dimension® Turf Herbicide

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

Dawn Lualturs

Dawn L. Walters

Regulatory Administrator

Phone 419-897-6790

Fax

419-891-2745

Email dawn\_walters@andersonsinc.com

Enclosure



18.00%

## Fertilizer with 0.072% Dimension® Turf Herbicide

### 18-0-0

If inhaled:

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

#### **GUARANTEED ANALYSIS**

70.00 70 Grow Hill oggil	
Plant nutrients derived from urea.	
ACTIVE INGREDIENT:	F292
Dithiopyr: 3,5-pyridinedicarbothiolc acid, 2-(difluoromethyl)-4-	
(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester	0.072%
INERT INGREDIENTS:	99.928%

Total Nitrogen (N).....

EPA Reg. No. 9198-118 EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A (Underlined letter is first letter used in run code on end of bag.)

Total .....

### KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eve irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prokonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals

#### FIRST AID

If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20
	minutes. Remove contacts, if present, after the first 5 minutes, then
	continue rinsing. Call a poison control center or doctor for treatment
	advice

Move person to tresh gir. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give any-

thing by mouth to an unconscious person. Take off contaminated clothing. Rinse skin immediately with plenty of if on skin

water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including gysters and shrimp. Use with care when applying to turt areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions lavor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 16.0 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau

#### GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turts, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ntals when used according to label directions

#### USE PRECAUTIONS

Turfgrass Safety: This product may be used on seeded, sodded, or sprigged lawns and ornamental turis that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turigrasses are tolerant to this product:

Cool-Season Grasses Caol-Sexson Grasses
Bentgrass, creeping\* (Agrostis palustris)
Bluegrass, Kentucky (Poa pratensis)
Fescue, fine\*\* (Festuca rubra)
Fescue, all (Festuca arundinaces)
Ryegrass, perennial (Lolium perenne)

warm-Season Grasses
Bahagrass (Paspalum notatum)
Bermudagrass (Cyrodon dactylon)
Buttalograss "" (Buchlee dactyloides)
Carpetgrass (Axonopus affinis)
Certipedegrass (Fernochba ophiuroides)
Kikuyugrass (Pennisetum clandestinum)
Si. Aupustinegrass (Staubraham esen St. Augustinegrass (Steriotaphrum secunda Zoyslagrass (Zoysia japonica)

'Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to onial Bentgrass (Agrostis tenuis) varieties

\*\*Use of this product on certain varieties of line fescue may result in undesirable turf Injuy. The following fine feacus varieties of his beaction in the seast of index and the seast of the product: Atlanta, Banner, Beauty, Bilgant, CF-2, Enjoy, HF-93, Highlight, Ivalo', Jameslown, Koket, Majenta, Mary, Pennlawn, Tamara', Tajiana', Waldorf and 'Walding' Not all varieties of fine feacue have been tested.

\*\*\*DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turl management practices before making

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseading, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

Other Use Precoutions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its lifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turks only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled.

### APPLICATION DIRECTIONS

Control of Crabgrass Preemergence and Early Postemergence Control: This product provides "preemergence" control of crabgrass (including the targe, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established fawns and ornamental turis. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 16 lbs. per 1,000 sq. ft. per cation, and no more than 48 lbs. per 1,000 sq. ft. per year.

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three product should be applied at the legislating and application and application of the state of the control programs fisted in Table 1 below, depending on the user's location, the turigrass mowing height, and whether the use is considered to be pre-merged or early postemergence at the time of the application. The duration of residued weed control ovided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turigrass competitiveness, and the user's

Use Program 1 for preemergence control at sites where the turfgrass is cut high ( $\widetilde{\omega}g_{*},\; \overline{\omega}$ homeowner lawns). This program provides 3-5 months of preemergence crabgration control. This program should not be used for early postemergence crabgrass control. O

Use Program 2 for preemergence control at sites where (a) turigrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early posternergence control at sites where turfgrass is cut high (e.g., homeowner lawns),

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turl maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turigrass is cut low (e.g., goll fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1: Recommended Application Use Rales* {Lbs. of this product/1,000 sq. ft.}						
PROGRAM	1	2	3 Preemergence (low)			
USE (turfgrass cut)	Preemergence (high)	Preemergence (low)				
REGION		Postemergence (high)	Pastemergence (low)			
North	4	6	8			
Transition	6	8	12			
South	8	12	16			
Coastal South	12	16**	16"			

\*DO NOT apply more than 16 lbs, per application, and no more than 48 lbs, per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI. South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

"May require split or sequential applications for full control.

Fall Applications: This product can also be applied in the tate summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control

Tins for improved Control: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) lications spaced 6-8 weeks apart may provide improved weed control

#### CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barnyardgrass, bluegrass (annual, roughstalk), crowfootgrass, foxtail (yellow & green)\*, goosegrass, kikuyugrass, ryegrass (annual & perennial), smutgrass, tespedeza (common), oxalis (buttercup, creeping & yellow woodsorrel), pursiane (common), speedwell (corn), spurge (prostrate

\*Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for late summer or early fall use, for craburass rail Applications: Used as utelected for fate southine for early failuse, for Jourglass control through the early part of the next spring, this product will also control (at the "Program 3" rates) for suppress of the "Program 2" rates) the following weeds when applied prior to their emergence; bluegrass (annual) - Pod annua, bittercress, chickweed. henbit, geranium (Carolina), paraley-piert, pineappleweed, shepherdspurse.

#### APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotaly-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or over-laps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

#### SUGGESTED SPREADER SETTINGS

These suggested speader settings are no linemede to replace calibration.

Please calibrate your speader before applying product.

50 lbs. treats 12,500 sq. n. (0,29 acre) at the 174 lbs. productbacre - LIGHT RATE (4,0 lbs./1,000 sq. lt.)

50 lbs. treats 3,000 sq. n. (0,19 acre) at the 261 lbs. productbacre - LIGHT RATE (4,0 lbs./1,000 sq. lt.) 50 fbs. treats 6,200 sq. ft. (0.14 acre) at the 348 fbs. product acre - MEDIUM RATE (8.0 fbs./1,000 sq. ft.)

SPREADER, C	GROUND SF-CED	WHOTH OF COVERAGE	PGHT RATE 0,13 ids, UAcre	LOW RATE 0.19 lbs. al/Acre	MEDIUM RATE 0.25 lbs. ai/Acre
Andersons AccuPr=2000, 57-500 (cone 2)	3-noh	Si fee	วี เมื	V 3/4	N 3/4
Scotts R-8, R-BA (come 2)	3 mph	11 feet	L 1/2	V. 3/4	0
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern III-A	4.5 mph	26 faet	5 1/4	6 1/4	7 1/4
Vicon (03 Ciries)	4,5 mph.	28 lest	28	35	42

STOCAGE AND DISPOSAL

Do not Contaminate water, Ictid, otifeed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs, and domestic water supplies.

PESTICIDE DISPOSAL. Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary tandfill or by lincineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

#### IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

#### WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the tabel when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/

Product of USA

Manufactured by: The Andersons Lown Fertilizer Division, Inc. P.O. Box 119 Maumee, OH 43537

The Andersons logo is a registered trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences LLC.

> **SGN 215** Net Wt. 50 Lbs. (22.68 kg)

> > APT18DM50 B8 (APTBLKPMBRNB)

