

PM23

9198-120

1.8.19

9198-120C
Page 1 of 10

(FRONT PANEL)

COMMERCIAL USE ONLY

THE ANDERSONS DIMENSION™ HERBICIDE II

This product is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

ACTIVE INGREDIENTS:

*Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester..... 0.103%

INERT INGREDIENTS..... 99.897%

TOTAL..... 100.000%

™ Dimension is a Trademark of the Monsanto Company

KEEP OUT OF REACH OF CHILDREN**CAUTION****SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

NET WEIGHT _____ LBS.

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUN 9 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9198-120

EPA Reg. No. 9198-120

EPA Est No. 9198-OH-1

BEST AVAILABLE COPY

0719

(BACK PANEL)
COMMERCIAL USE ONLY

Net Weight _____ Lbs.

The Andersons Dimension™ Herbicide II

GUARANTEED ANALYSIS

| | |
|--|-----|
| Total Nitrogen(N)..... | 16% |
| Available Phosphoric Acid(P_2O_5)..... | 4% |
| Soluble Potash(K_2O)..... | 4% |

Primary Plant Food derived from: Urea, Muriate of Potash, Ammoniated Phosphates.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

CAUTION

CAUSES EYE IRRITATION. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

IF IN EYES: Immediately flush with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove individual to fresh air. Seek medical attention if breathing difficulty develops.

IF ON SKIN: Wash with plenty of soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent

BEST AVAILABLE COPY

(BACK PANEL)

COMMERCIAL USE ONLY

to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Read the entire label before using this product.

Use only according to label instructions.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

GENERAL INFORMATION

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

TURFGRASS SAFETY

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

BEST AVAILABLE COPY

4 of 19

(BACK PANEL)

COMMERCIAL USE ONLY

Cool-Season Grasses

Bentgrass, creeping*
Agrostis palustris
Bluegrass, Kentucky
Poa pratensis
Fescue, fine**
Festuca rubra
Fescue, tall
Festuca arundinacea
Ryegrass, perennial
Lolium perenne

Warm-Season Grasses

Bahiagrass
Paspalum notatum
Bermudagrass
Cynodon dactylon
Buffalograss***
Buchloe dactyloides
Carpetgrass
Axonopus affinis
Centipedegrass
Eremochloa ophiuroides
Kikuyugrass
Pennisetum clandestinum
St. Augustinegrass
Stenotaphrum secundatum
Zoysiagrass
Zoysia japonica

*DO NOT apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties. Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

**Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

RESEEDING, OVERSEEDING, OR SPRIGGING

Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

BEST AVAILABLE COPY

(BACK PANEL)
COMMERCIAL USE ONLY

GOLF COURSE PUTTING GREENS

DO NOT use this product on golf course putting greens.

OTHER USE PRECAUTIONS

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

) For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

DO NOT apply this product to pastures.

Keep people and pets off treated areas until dust has settled.

) APPLICATION DIRECTIONS

CONTROL OF CRABGRASS

Preemergence and Early Postemergence Control

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly

BEST AVAILABLE COPY

6 of 19

(BACK PANEL)

COMMERCIAL USE ONLY

preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

DO NOT apply more than 11 lbs. per 1000 sq. ft. per application, and no more than 33 lbs. per 1000 sq. ft. per year.

Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

BEST AVAILABLE COPY

(BACK PANEL)

COMMERCIAL USE ONLY

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Table 1: Recommended Single Application Use Rates*
(lbs/1000 sq. ft.)

| PROGRAM | 1 | 2 | 3 |
|------------------------|------------------------|--|---|
| USE (turfgrass cut) | Preemergence (high) | Preemergence (low) Postemergence (high) | Preemergence (low) Postemergence (low) |
| REGION | | | |
| North | 2.8 | 4 | 5.5 |
| Transition | 4 | 5.5 | 8.4 |
| South | 5.5 | 8.4 | 11 |
| Coastal South | 8.4 | 11 | 11** |

*DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

**May require split or sequential applications for full control.

Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring.

The fall application should be followed by an appropriate spring application to provide season-long control.

BEST AVAILABLE COPY

(BACK PANEL)

COMMERCIAL USE ONLY

Tips For Improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications. Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

| | |
|-------------------------------|--|
| barnyardgrass | lespedeza (common) |
| bluegrass (annual) | oxalis (buttercup, creeping & yellow woodsorrel) |
| crowfootgrass | |
| foxtail (yellow & green)* | purslane (common) |
| goosegrass | speedwell (corn) |
| kikuyugrass | spurge (prostrate & spotted) |
| ryegrass (annual & perennial) | |
| smutgrass | |

*Also controlled at the Program 2 rates.

Fall Applications. Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

| | |
|--------------------|---------------------|
| bluegrass (annual) | henbit |
| bittercress | geranium (Carolina) |
| chickweed | parsley-piert |
| pineappleweed | |

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader

BEST AVAILABLE COPY

(BACK PANEL)

COMMERCIAL USE ONLY

settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

Manufacturer's Warranty

This product conforms to the ingredient statement on the label. Manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed under normal use conditions. **FOLLOW DIRECTIONS CAREFULLY.** Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. When product is not used according to label instruction, buyer assumes all risks of use, storage and handling of this product.

THIS PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED. IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND.

Manufactured by: The Andersons
Lawn Fertilizer Division
P.O. Box 119
Maumee, OH 43537

BEST AVAILABLE COPY

10819

(BACK PANEL)

COMMERCIAL USE ONLY

Date Printed: 5/24/94
Date Approved:
Supersedes:
Label Change:

11819

(Front Panel)

Homeowner Use Only

The Andersons Dimension™ Herbicide II

This product is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

ACTIVE INGREDIENTS:

*Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester.....0.103%

INERT INGREDIENTS.....99.897%
100.000%

™ Dimension is a Trademark of the Monsanto Company

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel For Additional Precautionary Statements

Net Weight _____ Lbs.

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUN 9 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9198-120

EPA Reg. No. 9198-120

EPA Est. No. 9198-OH-1

12 of 19

(Back Panel)

HOMEOWNER USE ONLY

Net Weight _____ Lbs.

The Andersons Dimension™ Herbicide II

Guaranteed Analysis

Total Nitrogen(N) 16%
Available Phosphoric Acid(P_2O_5)
4%
Soluble Potash(K_2O) 4%
Primary Plant Food derived from: Urea, Muriate of Potash,
Ammoniated Phosphates.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals
Keep out of reach of children.

CAUTION

CAUSES EYE IRRITATION. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists.

IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

IF ON SKIN, wash with plenty of soap and water.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

BEST AVAILABLE COPY

13 of 19

(Back Panel)

HOMEOWNER USE ONLY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Read the entire label before using this product.

Use only according to label instructions.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

GENERAL INFORMATION

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turfgrass Safety

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses

Bentgrass, creeping
Bluegrass, Kentucky
Fescue, fine
Fescue, tall
Ryegrass, perennial

Warm-Season Grasses

Bahiagrass
Bermudagrass
Buffalograss
Carpetgrass
Centipedegrass
Kikuyugrass
St. Augustinegrass

BEST AVAILABLE COPY

Zoysiagrass
(Back Panel)

HOMEOWNER USE ONLY

Reseeding, Overseeding, or Sprigging

Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

Other Use Precautions

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf.

For best results, cultural practices that disturb the soil, such as core aerification or verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

DO NOT apply this product to pastures.

Keep people and pets off treated areas until dust has settled.

BEST AVAILABLE COPY

(Back Panel)

HOMEOWNER USE ONLY

APPLICATION DIRECTIONS

Control of Crabgrass

Preemergence and Early Postemergence Control.

- This product provides "preemergence" control of crabgrass (both large and smooth species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.
- It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to identify the very small, early stages of crabgrass in well-established lawns and ornamental turfs.
- So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly pre-emergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Timings.

- This product may be applied as a single application or as a sequential application for crabgrass control. DO NOT apply more than 33.6 lbs. of this product/1000 sq. ft. per year.
- Applications may be made in the spring and in the fall.

Length of Control.

- The length of time of residual weed control provided by this product is directly related to the total rate applied.
- Depending on the desires of the user, this product may be applied at the use rates recommended in Table 1 below for the user's particular region to get 2-3 months of weed control, or it may be applied at rates listed in Table 2 below to get 3-5 months of control from date of application.
- The actual length of control will vary somewhat depending on weather, weed pressure, and the user's location within the North or South region.

BEST AVAILABLE COPY

11/8/19

(Back Panel)

HOMEOWNER USE ONLY

Table 1:

Recommended Use Rates for 2-3 Months of Crabgrass Control*
(lbs. of this product/1000 sq. ft.)

| Region | Preemergence | Postemergence |
|--------|--------------|---------------|
| North | 2.2 | 4.0 |
| South | 4.0 | 8.3 |

Table 2:

Recommended Use Rates for 3-5 Months of Crabgrass Control*
(lbs. of this product/1000 sq. ft.)

| Region | Preemergence | Postemergence |
|--------|--------------|---------------|
| North | 2.8 | 5.6 |
| South | 5.6 | 11.2 |

*South: includes AL, AR, AZ, CA, FL, GA, HI, LA, MS, NC, NM, NV, OK, SC, TN, TX, and VA. North: includes all other states.

Postemergence control is limited; see the "Crabgrass Control" and "Precautions" sections above.

Spring Applications.

- When applied in the spring, a single application of this product at the appropriate "preemergence" rate listed in Table 1 above will provide control of crabgrass for 2-3 months after application.
- Use the preemergence rate listed in Table 2 when the application is made very early in the spring (mid-February to mid-April) and longer control is required.
- Alternatively, a second ("sequential") application at the preemergence rate listed in Table 1 can be made 6-8 weeks after an early first application at the same rate to provide extended control where sequential fertilizer applications are recommended.
- When crabgrass has (or is likely to have) already emerged from the ground in your area and early postemergence control

BEST AVAILABLE COPY

of crabgrass is desired, use the appropriate "postemergence" rate listed in Table 1.

(Back Panel)

HOMEOWNER USE ONLY

- Use the postemergence rate listed in Table 2 when crabgrass has already emerged, but it is still relatively early in the spring.

Fall Applications.

- This product can also be applied in the late summer or early fall (late August through October) at the preemergence use rates listed in Table 2 to provide control of crabgrass through the early part of the next spring.
- The fall application should be followed by a mid-to-late spring application at the appropriate postemergence rate to provide season-long control.

Tips for Improved Control.

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Control of Other Grass and Broadleaf Weeds

Spring Applications. When applied as directed for crabgrass control in the spring, this product will also control (at the "postemergence" rates) or suppress (at the "preemergence" rates) the emergence of the following weeds:

| | |
|-------------------------------|--|
| barnyardgrass | lespedeza (common) |
| bluegrass (annual) | oxalis (buttercup, creeping & yellow woodsorrel) |
| crowfootgrass | |
| foxtail (yellow & green)* | purslane (common) |
| goosegrass | speedwell (corn) |
| kikuyugrass | spurge (prostrate & spotted) |
| ryegrass (annual & perennial) | |
| smutgrass | |

*Also controlled at the preemergence rates.

BEST AVAILABLE COPY

18 of 19

(Back Panel)

HOMEOWNER USE ONLY

Fall Applications. When applied as directed for early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "postemergence" rates) or suppress (at the "preemergence" rates) the emergence of these weeds:

bluegrass (annual)
bittercress
chickweed
pineappleweed

henbit
geranium (Carolina)
parsley-piert

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

BEST AVAILABLE COPY

19 of 19

(Back Panel)

HOMEOWNER USE ONLY

Manufacturer's Warranty

This product conforms to the ingredient statement on the label. Manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. When product is not used according to label instruction, buyer assumes all risks of use, storage and handling of this product.

THIS PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED. IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND.

Manufactured by:
THE ANDERSONS
LAWN FERTILIZER DIVISION
P.O. BOX 119
MAUMEE, OH 43537

Date Printed: 5/24/94
Date Approved:
Supersedes:
Label Change:

BEST AVAILABLE COPY