

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

9198-120

3/13/2007

8 A

5 3

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Dawn L. Walters
Product Registration
The Andersons Lawn Fert. Div., Inc
Fert. PO Box 120
Maumee, OH 43537

MAR 13 2007

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Alternate Brand Name "Intermountain Farmers Association Crabgrass &
Spurge Preventer with Nitrogen"
EPA Reg. No. 9198-120
Application Dated January 11, 2007

Dear: Ms. Walters,

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 01/11/07 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

802



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 9198-120	2. EPA Product Manager Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Dimension Herbicide II	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow Fert. PO Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Alternate Brand Names per PR Notice 98-10 : Intermountain Farmers Association Crabgrass & Spurge Preventer with Nitrogen

This notification is consistent with the guidance of PR Notice 98-10 and the requirements of EPA's regulations at 40 CFR 156.10 and 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 98-10 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Dawn L. Walters	Title Regulatory Administrator	Telephone No. (Include Area Code) 334-262-2766	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature Dawn L. Walters	3. Title Regulatory Administrator		
4. Typed Name Dawn L. Walters	5. Date 1/11/87		



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

January 11, 2007

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U. S. Environment Protection Agency
1200 Pennsylvania Ave., NW
Washington, D.C. 20460-001

Subject: NOTIFICATION - Additional Brand Names

**RE: The Andersons Dimension Herbicide II
EPA Reg. No. 9198-120**

Per PR Notice 98-10, enclosed are two EPA Forms 8570-1, applications notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand names are:

**Intermountain Farmers Association Crabgrass & Spurge Preventer with Slow Release Nitrogen
Intermountain Farmers Association Crabgrass & Spurge Preventer with Nitrogen**

One copy of each final printed label is enclosed.

Enclosed are self-addressed cards indicating the acceptance or rejection of these notifications. Please complete the cards and return via mail.

If there are any questions, please contact me.

Sincerely,
The Andersons Lawn Fertilizer Division, Inc.

Dawn L. Walters

Dawn L. Walters
Regulatory Administrator
Phone 419-351-3717
Email dawn_walters@andersonsinc.com

enclosure

3/4



Crabgrass & Spurge Preventer with Nitrogen

23-3-3

- This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

GUARANTEED ANALYSIS

Total Nitrogen (N).....	23.00%
1.18% Ammoniacal Nitrogen	
21.82% Urea Nitrogen	
1-methylphosphoryl-6-(trifluoromethyl)-S,S-dimethyl ester	3.00%
1-methylphosphoryl-6-(trifluoromethyl)-S,S-dimethyl ester	3.00%

Plant nutrients derived from ammonium phosphate, urea and potassium chloride.
Chlorine (Cl) Max

2.54%

F292

ACTIVE INGREDIENT:

Diflupyr: 3,5-pyridinedicarboxylic acid, 2-(difluoromethyl)-4-(2-methylphosphoryl)-6-(trifluoromethyl)-S,S-dimethyl ester

0.103%

INERT INGREDIENTS:

99.897%

Total 100.000%

EPA Reg. No. 9198-120

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

(Underlined letter is first letter used in run code on end of bag.)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Keep the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8991 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for commercial sod production, or for research purposes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 11.0 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year, and is prohibited from use in Nassau and Suffolk Counties.

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turfgrass Safety: This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses

Bentgrass, creeping (Agrostis palustris)
Bentgrass, Kentucky (Poa pratensis)
Fescue, fine (Festuca tenuis)
Fescue, tall (Festuca arundinacea)
Ryegrass, perennial (Lolium perenne)

Warm-Season Grasses

Bahagrass (Paspalum notatum)
Bermudagrass (Cynodon dactylon)
Buckgrass (Echinochloa crusgalli)
Carpetgrass (Axonopus affine)
Centipedegrass (Eriochloa fasciculata)
Cynodactylon (Cynodon dactylon)
St. Augustinegrass (Stenotaphrum secundatum)
Zoysiagrass (Zoysia japonica)

*Use of this product on certain varieties of creeping bentgrass, such as "Cohasset", "Carmen", "Seaside", and "Washington" may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (Agrostis tenuis) varieties.

**Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: "Atlanta", "Banner", "Beauty", "Bigcat", "CF-2", "Enjoy", "HF-8", "Highlight", "Ivala", "Jamestown", "Kokoi", "Majesta", "Mary", "Pentlawn", "Tanjana", "Tanjana", "Waklont", and "Watkins". Not all varieties of fine fescue have been tested.

***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully grown and has established new roots.

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

Re-seeding, Overseeding, or Sprigging: Re-seeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When re-seeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (sift seeders) designed to give good seed to soil contact.

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aeration, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

Control of Crabgrass

Preemergence and Early Postemergence Control: This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 11 lbs. per 1,000 sq. ft. per application, and no more than 33 lbs. per 1,000 sq. ft. per year.

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split between two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1: Recommended Application Use Rates* (Lbs. of this product/1,000 sq. ft.)				
PROGRAM	1	2	3	
USE (turfgrass cut)	Preemergence (high)	Preemergence (low)	Preemergence (low)	
REGION		Postemergence (high)	Postemergence (low)	
North	2.6	4.0	5.5	
Transition	4.0	5.5	8.4	
South	5.5	8.4	11.0	
Coastal South	8.4	11.0	13.6	

*DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

**May require split or sequential applications for full control.

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barnyardgrass, bluegrass (annual, roughstalk), crowfootgrass, foxtail (yellow & green), goosegrass, karyograss, ryegrass (annual & perennial), smartweed, spodiopogon (common), oxalis (buttercup, creeping & yellow wood-sorrel), purslane (common), sparrowweed (corn), spurge (prostrate & spotted).

*Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: bluegrass (annual), Poa annua, buttercup, chickweed, henbit, geranium (Carolina), parsley-piert, pinappleweed, shepherdspurse.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product

uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Aerial streaking, clumps, or clumps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

SUGGESTED SPREADER SETTINGS					
These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.					
50 lbs. covers 12,500 sq. ft. at the 17 1/2 lbs. product/acre - MEDIUM RATE (40 lbs./1,000 sq. ft.)					
50 lbs. covers 9,000 sq. ft. at the 240 lbs. product/acre - HIGH RATE (6.5 lbs./1,000 sq. ft.)					
50 lbs. covers 5,000 sq. ft. at the 365 lbs. product/acre - HEAVY RATE (8.4 lbs./1,000 sq. ft.)					
SPREADER	GROWTH SPEED	WIDTH OF CUTTER, FEET	MEDIUM RATE (40 lbs./1,000 sq. ft.)	HIGH RATE (6.5 lbs./1,000 sq. ft.)	HEAVY RATE (8.4 lbs./1,000 sq. ft.)
Andros ACP-2000, SR-2000 (30" x 4")	3 mph	11' 1/2"	12	M	N 1/2
Andros ACP-1000	3 mph	11' 1/2"	12	M	N 1/2
Scots Rotator R-6, R-6A (30" x 4")	3 mph	11' 1/2"	12	K 1/2	M 1/4
Lilly Mow-Master WTR, WTR 161, 160 (17" x 14" x 10") Pattern R-A	4.5 mph	40' 1/2"	5	6	7 1/2
Vicon (35" x 40")	4.5 mph	36' 1/2"	36	42	57 1/2

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs, and domestic water supplies.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION.

Product of USA

Manufactured by:
Free Flow Fertilizer
P.O. Box 119
Mammoth, OH 43053

Net Wt. 50 Lbs. (22.68 kg)

Covers up to
12,500 Sq. Ft.



3 18655 50917 5

IFA23DM5

L6 (BLKP1929B)

5/5

NEXT

LABEL



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

MAR 13 2007

Ms. Dawn L. Walters
Product Registration
The Andersons Lawn Fert. Div., Inc
Fert. PO Box 120
Maumee, OH 43537

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Alternate Brand Name "Intermountain Farmers Association Crabgrass &
Spurge Preventer with Slow Release Nitrogen"
EPA Reg. No. 9198-120
Application Dated January 11, 2007

Dear: Ms. Walters,

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 01/11/07 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
Washington, DC 20460

Registration
Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 9198-120	2. EPA Product Manager Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Dimension Herbicide II	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow Fert. PO Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Alternate Brand Names per PR Notice 98-10 : Intermountain Farmers Association Crabgrass & Spurge Preventer with Slow Release Nitrogen

This notification is consistent with the guidance of PR Notice 98-10 and the requirements of EPA's regulations at 40 CFR 156.10 and 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 98-10 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	
		5. Location of Label Directions <input type="checkbox"/> On Label	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Dawn L. Walters		Title Regulatory Administrator	Telephone No. (Include Area Code) 334-262-2766
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature Dawn L. Walters		3. Title Regulatory Administrator	
4. Typed Name Dawn L. Walters		5. Date 7/11/07	



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

January 11, 2007

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U. S. Environment Protection Agency
1200 Pennsylvania Ave., NW
Washington, D.C. 20460-001

Subject: NOTIFICATION - Additional Brand Names

**RE: The Andersons Dimension Herbicide II
EPA Reg. No. 9198-120**

Per PR Notice 98-10, enclosed are two EPA Forms 8570-1, applications notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand names are:

**Intermountain Farmers Association Crabgrass & Spurge Preventer with Slow Release Nitrogen
Intermountain Farmers Association Crabgrass & Spurge Preventer with Nitrogen**

One copy of each final printed label is enclosed.

Enclosed are self-addressed cards indicating the acceptance or rejection of these notifications. Please complete the cards and return via mail.

If there are any questions, please contact me.

Sincerely,
The Andersons Lawn Fertilizer Division, Inc.

Dawn L. Walters
Regulatory Administrator
Phone 419-351-3717
Email dawn_walters@andersonsinc.com

enclosure

3/4



Crabgrass & Spurge Preventer with Slow Release Nitrogen

23-3-3

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

GUARANTEED ANALYSIS

Total Nitrogen (N).....	23.00%
1.18% Ammoniacal Nitrogen	
21.82% Urea Nitrogen*	
Available Phosphate (P ₂ O ₅).....	3.00%
Potash (K ₂ O).....	3.00%
Sulfur (S).....	1.71%
1.71% Free Sulfur (S)	

Plant nutrients derived from ammonium phosphate, urea, polymer coated sulfur coated urea and potassium chloride. Chlorine (Cl) Max. 2.54% 5.75% Slowly available nitrogen from polymer coated sulfur coated urea.

F292

ACTIVE INGREDIENT:

Dithiopyr: 3,5-pyridinedicarboxylic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester	0.103%
INERT INGREDIENTS:	99.897%
Total.....	100.000%

EPA Reg. No. 9198-120
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
(Underlined letter is first letter used in run code on end of bag.)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Cause eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If eyes: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including crayfish and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to individual areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 11.0 pounds per 1,000 square foot (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm newly established ornamentals when used according to label directions.

USE PRECAUTIONS

Turfgrass Safety: This product may be used on aerated, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well established, or on those that have been weakened by weather, pest, disease, chemical, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses

Bentgrass, creeping (*Agrostis perennis*)
Bluegrass, Kentucky (*Poa pratensis*)
Fescue, fine (*Festuca rubra*)
Fescue, tall (*Festuca ovina*)
Ryegrass, perennial (*Lolium perenne*)

Warm-Season Grasses

Bahia grass (*Passatum notatum*)
Bermudagrass (*Cynodon dactylon*)
Buffalograss (*Buchloe dactyloides*)
Carpentergrass (*Paropogon affine*)
Centipedegrass (*Eriochloa ciliaris*)
Kleingrass (*Pennisetum clandestinum*)
St. Augustinegrass (*Stenotaphrum secundatum*)
Zoysiagrass (*Zoysa japonica*)

*Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Soaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (*Agrostis tenuis*) varieties.

**Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Allanta', 'Banner', 'Beauty', 'Bigant', 'CF-2', 'Enjoy', 'HF-80', 'Highlight', 'Ivala', 'Jamestown', 'Kokoi', 'Majesta', 'Mary', 'Pentlawn', 'Tanjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

Re-seeding, Overseeding, or Sprigging: Re-seeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When re-seeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (silt seeders) designed to give good seed to soil contact.

Other Use Precautions: The early postemergence crabgrass control described below is limited. Post-emergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core, spike, or hydro-aeration, and verticuring, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns and ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use chippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until soil has settled.

APPLICATION DIRECTIONS

Control of Crabgrass

Preemergence and Early Postemergence Control: This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Post-emergence crabgrass control will be obtained only when this product is applied prior to the falling of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 11 lbs. per 1,000 sq. ft. per application, and no more than 33 lbs. per 1,000 sq. ft. per year.

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence or crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence or crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1: Recommended Application Use Rate* (Lbs. of this product/1,000 sq. ft.)			
PROGRAM	1	2	3
USE (turfgrass cut)	Preemergence (high)	Preemergence (low)	Preemergence (low)
REGION		Postemergence (high)	Postemergence (low)
North	2.6	4.0	5.5
Transition	4.0	5.5	5.5
South	5.5	8.4	11.0*
Coastal South	8.4	11.0*	11.0*

*DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Post-emergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas.
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY, & RI.
South: AL, AR, AZ, CA, GA, IA, MS, NC, NM, NV, OK, SC, TN, & TX.
Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

**May require split or sequential applications for full control.

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control: For best results, apply this product within a few days after mowing and before moving again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barnyardgrass, bluegrass (annual, roughstalk), crowfootgrass, foxtail (yellow & green), goosegrass, kyllburggrass, ryegrass (annual & perennial), smartgrass, sparrowgrass (common), oxalis (heart-shaped, creeping & yellow wood-scented), purslane (common), speedwell (common), spurge (prostrate & spotted).

*Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: bluegrass (annual) - *Poa annua*, bittercrass, chickweed, henbit, geranium (Carolina), juncoweed, pineappleweed, shepherdspurse.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product

uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. A, 3rd streaking, skips, or overlaps during application. Check equipment for uniformity to ensure equipment is functioning properly and applying uniform distribution of granules.

SUGGESTED SPREADER SETTINGS (These suggested, A, S, settings are not intended to replace label directions.) Please calibrate your spreader before applying. Output: 50 lbs. covers 12,500 sq. ft. at the 174 lbs. product/acre - MEDIUM RATE (40 lbs./1,000 sq. ft.) 50 lbs. covers 9,000 sq. ft. at the 240 lbs. product/acre - HIGH RATE (55 lbs./1,000 sq. ft.) 50 lbs. covers 5,500 sq. ft. at the 366 lbs. product/acre - HEAVY RATE (84 lbs./1,000 sq. ft.)					
SPREADER	GROUND SPEED	WHEEL HOF COEFFICIENT	MEDIUM RATE 0.18 lbs./A	HIGH RATE 0.25 lbs./A	HEAVY RATE 0.36 lbs./A
Andersens AcolPro 200... SR-200 (pins 4)	3 mph	1.0	L 1/2	M	H 1/2
Andersens AcolPro 1000	3 mph	1.1	L 1/2	M	H 1/2
Solis P-Series R4, R-BA (pins 4)	3 mph	1.1	J 3/4	K 1/2	M 1/4
LadyMaid WTR WTR 1250 (PFO #450) P-Series 4	4.5 mph	40 test	5	6	7 1/2
Vicon (33 Series)	4.5 mph	36 test	36	42	57 1/2

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs, and domestic water supplies.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permitted, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION.

Product of USA

Manufactured by:
Froo Fertilizer
P.O. Box 119
Mauern, OH 43537

Net Wt. 50 Lbs. (22.68 kg)

Covers up to
12,500 Sq. Ft.



IFA235DM5
L6 (BLKP1929B)

Handwritten mark: 2/5