

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

JAN 0 8 2008

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Ms. Dawn Walters Product Registration The Andersons Lawn Fert. Div., Inc PO Box 119 Maumee, OH 43537

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request Alternate Brand Name"The Andersons Golf Products Contec DG Dispersing Granule Technology with 0.103% Dimension Turf Herbicide'

EPA Reg. No. 9198-120

Application Dated December 11, 2007

Dear Ms Walters:

The Agency is in receipt of your 12/11/07 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped Notification and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader

Registration Division (7505P) Office of Pesticide Programs

Please read instructions on reverse before con. ing form.			Form Approv			OMB No. 2	070-006	O. Approval expires 2-28-95	
United States Environmental Protecti Washington, DC 20			- -			Registra Amendn Other		OPP Identifier Number	
	4	pplication	n for Pesti	cide - Sec	tion	l			
1. Company/Product Number 9198-120			2. EF Mille	PA Product Man	ager		3. Pr	oposed Classification	
4. Company/Product (Name) The Andersons Dimension Herb	icide II	79	PM# 23						
5. Name and Address of Applica	ant (Include ZIP Cod	e)	6. E	xpedited Rev	reiw.	In accordar	nce with	FIFRA Section 3(c)(3)	
The Andersons Lawn Fert PO Box 119 Maumee, OH 43537		Free Flow F	ert. (b)(i)	, my product	is simi	ilar or identi	cal in co	emposition and labeling	
Check if this is	a new address		Pro	duct Name					
			Section	- 11					
Amendment - Explain be Resubmission in respons Notification - Explain bel	se to Agency letter d	lated		Final printe Agency let "Me Too"	ter date Applica	ition.	to		
Explanation: Use additional provided in the state of the labeling or the confidential further understand that if this notification of FIFRA and I may be subject to the labeling or the confidential further understand that if this notification of FIFRA and I may be subject to the labeling or the confidential further understand that if this notification of FIFRA and I may be subject.	per PR Notice 98-10 : A guidance of PR Notice 9 al statement of formula o ion is not consistent with	Andersons Golf 98-10 and the re f this product. I the guidance o	Products® Contect equirements of EPA understand that it if f PR Notice 98-10 a	DG® Dispersing G 's regulations at 40 is a violation of 18 and the requiremen) CFR 15 U.S.C. 5	56.10 and 40 CF Sec. 1001 to will	R 152.46 a	and no other changes have been any false statement to EPA. I	
			Section	- 111					
1. Material This Product Will Be	Packaged In:								
Yes No	Yes Yes No No No ner		Water Solubl Yes No If "Yes"	e Packaging No. per	2. Type of Container Metal Plastic Glass Paper				
be submitted	nit Packaging wgt.	container	Package wgt	containe	ır		Other (Specify)	
3. Location of Net Contents Info Label Cont 6. Manner in Which Label is Aff	tainer	4. Size(s) Ret	raph	Othe	<u> </u>	Cetion of Lab		ons	
		Paper Stence		···					
			Section						
1. Contact Point IComplete itel	ms directly below fo	r identificatio	n of individual to	o be contacted,	if nec	essary, to pro	cess this	application.)	
Name Dawn L. Walters			Domilatani Administratas			Telephor 419-897	ne No. (Include Area Code) -6790		
l certify that the statement acknowledge that any k both under applicable law	nowlinglly false or m		all attachments		•			6. Cate Application F.coeived (Stamped)	
2. Signature Dawn Kllalter			3. Title Regulatory Adm	ninistrator					
4. Typed Name Dawn L. Walters			5. Date	11107					



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

December 11, 2007

Document Processing Desk (NOTIF)
Office of Pesticide Programs - (7504P)
U. S. Environment Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Subject:

NOTIFICATION - Additional Brand Name

RE:

The Andersons Dimension Herbicide II

EPA Reg. No. 9198-120

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

Andersons Golf Products[®] Contec DG[®] Dispersing Granule Technology with 0.103% Dimension[®] Turf Herbicide

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

Dawn L. Walters

Regulatory Administrator

Dawn & Walter

Phone 419-897-6790 Fax 419-891-2745

Email dawn_walters@andersonsinc.com

Enclosure







Dispersing Granule Technology with 0.103% DIMENSION® Turf Herbicide

24-0-10

This product is a herbicide that provides control of listed annual arasses and broadleaf weeds in established lawns and ornamental turfs.

	ACTIVE INC			
24.00%				
	Dithiopyr: 3.			
	2-(dilluoror			
	(tritluorome			
	INERT INGI			
10.00%	Total			
6.15%	EPA Reg. No. 9			
	EPA Est. No. 9			
6.15% Free Sulfur (S) Plant nutrients derived from methylene ureas, ureaform, potassium				
	(Underlined lett			
ureaform				
	6.15%			

GREDIENT: .5-pyridinedicarbothioic acid. omethyl)-4-(2-methylpropyl)-6-nethyl)-S,S-dimethyl ester . . . 0.103% REDIENTS: 99.897% 9198-OH-1M 9198-OH-2B 9198-AL-001A etter is first letter used in run code on end of bag)

Visit aur website: AndersonsGoliProducts.com

Exercise # 5GN at 50元 平安工业。

Net Wt 50 Lbs (22:68 kg)

Information regarding the contents and levels of metals in this product is available on the internet at

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye initiation. Harmful il inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with scap and water after handling. Remove contaminated dollring and wash before reuse. Prolonged or trequently repeated skin contact while handling this material may cause ellergic reaction in some individuals.

If in eyes: Hold eye open and rinse slowly and pently with water to minutes. Remove contacts, if present, after the first 5
minutes, then continue instinut. Call a poson control center or docto

If inhaled: Move person to fresh etc. If person is not breathing call in immune: wove person to use a a ... u person is not use a ling, can 1911 or an ambulance, then give artificial respiration, preferably imouth-to-mouth it cossible. Call a poison control center or doctor to

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing, Rose skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poisor strol center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including systems and shrimp. Use with care when applying to turf areas adjacent to any body of water. Oriti and runoff from treated turf areas appeared on jury or whose; in the art further aquatic sites. Do not apply directly to water, or to areas where surface water is present or to interfed a ease below the mean high water mark. Do not apply when weather conditions favor drift from theated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a yiolation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for cial seed production, or for research purposes

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 11.0 pounds por 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

GENERAL INFORMATION

This product is a herbickle that provides control of catograss and other armual grasses and broadleaf weeks in established turns and ornamental turfs, including golf course lainways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions

HISE ORECALITIONS

Turfgrass Safety: This product may be used on seeded, sodded, or sprigged lawns and ornamental turis that are well-established. The grass must have developed a good root system and a uniform stand. and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on towns and ornamental turfs that are ot well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress may increase the chances for furl injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turigrass species that have been determined to be tolerant to confications of this product. When applied as directed under the conditions described, the following established turkgrasses are tolerant to this product

rsiatusmed unigrasses are toleram to tirs Cool-Season Grasses Bentgrass, creeping (Agrostis palustris) Bluegrass, Kentucky (Poa pratensis) Fescue, fine'' (Festuca rubra) Fescue, tall (Festuca arundinacea) Ryegrass, perennial (Lotum perennel

Warm-Season Grasses Warm-Season Grasses Bahiagrass (Paspakim notatim) Bermudagrass (Cynodon dach fon) Bu'llalograss''' (Buchtoe dach/loide: Carpelgrass (Axonopus affinis) Centipedegrass (Eremochloa ophiuroides) Kikuyugrass (Pennisetum clandestinum) St. Augustinegrass (Stenotaphrum secundat Zoysiagrass (Zoysia japonica)

*Use of this product on certain varieties of creeping bentgrass, such as "Cohansey". "Carmen", "Seaside", and "Washington" may result in undestrable turtgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass

"I ke of this product on certain varieties of fine fescue may result in udesigable but injury. The following fine feacue varieties have bee found to be sensitive to this product: "Atlanta, "Barner, "Beauty, "Blgar," CF-2; "Enjoy, "HF-93, "Highfight, "Ivalo, "Lamestown, "Roler, Maghanta, "Mary, Perntlawn, Tamara, "Tagarat," Waldoof. and Walring. Not all varieties of fine tescue have been tested

"DO NOT use this product on seed no buffalowass in the spring of the first year of establishment until the turigrass is fully green and has established new roots.

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and tarf management practices before making widespread use.

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turigrasses. When producting which we seatched the order of the production state of the control of

Other Use Precautions: The early posterrengence crabgrass control described below is limited. Posterrengence applications are only effective on crabgrass if applied before crabgrass has displayed is little lad or trist tiler.

For best results, cultural practices that disturb the soil, such as core-spike-, or hydro-aerification, and verticulting, should be done before application of this product. DO NOT apply this product until the grass

Apply this product directly to established lawns or ornamental turbs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use dippings from treated furt for mulching around vegetables or truthness. Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

Control of Crabgrass Preemergence and Early Posternergence Control: This product Prevenegance and Early Posternergance Control: This product provides "prevenergence" crotel of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early posternergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often diffact to see the very small, early stages of crabgrass in well-established lawns and ornamental birds. nergence crabgrass control will be obtained only when this product is applied prior to the tilering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the tawn or turf. So the practical benefit of this products partial manufacture in the Covariant partial and the control of proceed to shirtly preemengence crabgress products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for catalgrass control in the spring, summer, or fall. DO NOT apply more than 11 lbs. per 1,000 sq. it, per application, and no more than 33 lbs nor 1,000 so, ft, our year.

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate conesponding to one of the three control programs fisted in Table 1 below, depending on the user's location, the turkgrass moving height, peon, operatory on the uses a occoron, the using as more groups, and whether the use is considered to be premieragene or early postemengence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat departing on rewarfer, weed pressure, urkgrass competitiveness, and the user's location within a recion.

Use Program 1 for preemergence control at sites where the turfgrass to cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) furfigrass is cut low (e.g., golf tarways), and (b) furfigrass maintenance or weed control has been conducted during the previous year. This programprovides 4-6 months of preemergence crabgrass control. This program may also be used for early posternergence control at site. where turigrass is cut high (e.g., homeowner law

Use Program 3 for preemergence control at sites where (a) tu. y: s is cut low (e.g., golf tanways), and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control.

This program may also be used for early postemeroence control at sites where turigrass is cut low (e.g., golf lairways;

Subsequent, sequential pre- and or posternergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be still across two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1: Recommended Application Use Rates' (Lbx. of this product/1,000 sq. ft.)						
PROGRAM	i	2	1			
USE (turigrass cut)	Preamergence (high)	Preemergence (forr)	Precesse grace (low)			
REGION		Postemergence (high)	Postenergence (low)			
Noth	2.5	4,0	5.5			
Transition	1.0	5.5	3,4			
South	5,5	8,4	11.0**			
Coastal South	5.4	11.9**	1:.0"			

*DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Recions include areas

Norm: All other areas.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT,

southern areas of II., IN, OH, & coastal areas of C1, NY, & RI., South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NY, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

"May require split or sequential applications for full control

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program
3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for improved Control: For best results, apply this product within a lew days after moving and delay moving again for a lew days after the application. When treated lawn or ornamental turl areas are watered or receive significant rantal within a lew days after application of this product, improved weed control may result Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or ss (at the 'Program 2' rates) the following weeds when suppress (all the "Program 2" rates) the following weeds when applied prior to the inengence: harmycrigness, thougass (arrusal, coughstally, cow/bodyrass, footal (rystow & green)**, goosegrass, foliated (reserving, smurghass, segredate) (common), caudia (futherum, deeping & yedow woodsomel). prustane (common), speedvell (com), syunge (mostrata & spotted). "Also controlled at the "Program 2" rates.

Fell Applications: Used as directed for late summer or early tall use, for crabgrass control through the early part of the next spring, this pro "act will also control fat the "Program 3" rates) or suppress In pr. 12(the ascontrol is the intogram's laws) or suppressing "in "in "y yam 2" rates the indowing w., 36 when applied prior to their emeritance hiteograss, annual). "Out armus, biserciess, chickened, henbit geranium (Carolina), par "ay-pie", prinapplewed, shepher(Surse, uns.).

A. PLICATION SOUPMENT AND INSTRUCTIC. IS Apply this product with drop or rotary-type spreaders design: 1 to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufative risk directions for adjusting the spreader settings such

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. STORAGE: Keep container closed and away from food, feedstuffs, and domestic water supplies.
PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper

disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Region

guarance.
CONTAINER DISPOSAL: Completely empty bag into application
equipment. Then dispose of empty bag in a sanitary landitif or by
incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

IMPOHIANT: HEAD BEFORE USE.

Read the entire Decisions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY: Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OF FITNESS FOR A
PARTICLE AR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSCOUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice.

To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:
THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATEJURISDICTION.

Product of USA Manufactured by: The Andersons Lown Fertilizer Division Inc. P.O. Box 119, Maumee, OH 43537

www.andersonsgolfproducts.com
Andorsens Cof Products and Contro DC are registered trademarks of The
Andersons, Inc. Dimension is a registered trademark of Don Agrosciences LLC.

AGC2424WDGDM5

(BLKP1929B) - 1.7

SUGGESTED SPREADER SETTINGS SUGGESTED SPREADER SETTINGS.

These approaches generate change are on the inducted in replace contration.

50 bits, comes a 7.000 cm. Section of the contration of the contrat

	GROUND SPEED		SPREADER SETTINGS				
SPREADER		WIDTH OF COVERAGE	UGHT RATE 0,13 fb, aVA	LOW RATE 0.18 lb. spA	REORUM RATE 0.25 lb. af A	HIGH RATE 0.30 lb, MA	
Andersons Actualite 2002 SR-2000 (come 6)	3 mpts	1º feyt	K 1/4	ń	812	012	
Andersons Acculing 1000	3 meh	It feet	8.174	H	#12	012	
Socs8-8, R-8A (core 5)	i mph	11 less	J 1/4	K 1/2	M12	F12	
Lefy Vosels with with the 1250 (CTO at 450 poor Patrick IB-C	4,5 mph	34 10 25	434	534	£12	/34	
Vican (03 Senes)	45 mgh	20 feet	16	20	24	31	



