

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

FEB 2 2 2008

Ms. Dawn Walters Product Registration The Andersons Lawn Fert. Div., Inc PO Box 119 Maumee, OH 43537

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request Alternate Brand Name "The Andersons Turf Products Fertilizer with

0.103% Dimension Turf Herbicide"

EPA Reg. No. 9198-120

Application Dated January 28, 2008

Dear Ms Walters:

The Agency is in receipt of your 01/28/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader

Registration Division (7505P)

Office of Pesticide Programs

24

Please read instructions	on reverse before comp.	⊿ng form.		Form App	roved.	JMB No. 2	070-006	D. Approval expires 2-28-95	
United States Environmental Protection Agency Washington, DC 20460						Registra Amendn Other		OPP Identifier Number	
		Application	n for Pestici	de - Sect	ion I			· <u></u>	
1. Company/Product Nur 9198-120	nber		2. EPA Kraft	Product Mana	ger		3. Pr	oposed Classification	
4. Company/Product (Name) The Andersons Dimension® Herbicide II			РМ# 23						
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow I PO Box 119 Maumee, OH 43537 Check if this is a new address			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name						
			Section -						
Amendment - Exp Resubmission in Notification - Exp	response to Agency lette	r dated			er date pplica	tion.	to		
This notification is consistent made to the labeling or the co further understand that if this	d Name per PR Notice 98-10 : with the guidance of PR Notice onfidential statement of formula notification is not consistent who be subject to enforcement acti	e 98-10 and the re a of this product. I ith the guidance o	equirements of EPA's re understand that it is a f PR Notice 98-10 and under sections 12 and	egulations at 40 (violation of 18 U the requirements 14 of FIFRA.	CFR 15 I.S.C. S	6.10 and 40 Cf ec. 1001 to will	FR 152.46 a fully make a		
			Section - I	11					
1. Material This Product Will Be Packaged In: Child-Resistant Packaging Yes No * Certification must be submitted Unit Packaging Yes No If "Yes" Unit Packaging wgt. No. per container			No lf "Yes" No. per			Metal Plastic Glass Paper	Metal Plastic Glass		
Label Container 6. Manner in Which Label is Affixed to Product Label			tail Container	S. Location of Label Directions On Label Other					
		Stenc	glued iled						
	· · · · · · · · · · · · · · · · · · ·		Section - I					 	
	lete items directly below	for identification		e contacted,	if nece	ssary, to pro			
Name Dawn L. Walters			Dogulaton, Administrator			•	shone No. (Include Area Code) -897-6790		
•	tatements I have made on it any knowlingliy false on able law.		l all attachments th				•	6. Date Application Received (Stamped:	
2. Signature Dawn & Walters			3. Title Regulatory Administrator						
4. Typed Name Dawn L. Walters			5. Date 1/28/09						



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

January 29, 2008

Document Processing Desk (NOTIF) Office of Pesticide Programs - (7504P) U. S. Environment Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

Subject:

NOTIFICATION - Additional Brand Name

RE:

The Andersons Dimension® Herbicide II

EPA Reg. No. 9198-120

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons® Turf Products Fertilizer with 0.103% Dimension® Turf Herbicide

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

Dawn L. Walters

Regulatory Administrator

Dawn & Walters

Phone 419-897-6790

Fax

419-891-2745

Email dawn walters@andersonsinc.com

Enclosure



Fertilizer with 0.103% Dimension® Turf Herbicide

19-0-0

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turks

GUARANTEED ANALYSIS

Total Nitrogen (N)	19.00%
•	

Plant nutrients derived from urea and polymer coated sulfur coated

*4.75% Slowly available nitrogen from polymer coated sulfur coated urea.

ACTIVE INGREDIENT:

Dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-	
(2-methylpropyl)-6-(trilluoromethyl)-S.S-dimethyl ester	0.103%
(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester	99.897%
Total	100.000%

EPA Reg. No. 9198-120 EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

(Underlined letter is first letter used in run code on end of bar.)

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye igritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove nated clothing and wash before rouse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to Iresh air, if person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treat-

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic ** fish and hig..ly .o...c to other aquatic organisms including bysters and strimp. Use with mare vien applying, a furl awas adjacent to any body of water. Dutt and sund from treated furl may adversely affect aquatic organisms in adjacent equations also. Do not apply directly to mater, or to an us where surface water is present, or to intertidal areas below the mean high water man. No no. apply when weather conditions favor drift from treated areas. On not contamine a water . A. in disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on till being grown fir sales to their commercial use as sod, or for commercial seed production, or for research purpoles.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 11.0 pounds per 1,000 square feel (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau

GENERAL INFORMATION

This product is a herbicide that provides control of crabgiass and other annual grasses and broadleaf weeds in established lawns and ornamental turks, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turigrass Safety: This product may be used an seeded, sodded, or sprigged lawns and arramental turis that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather- pest- disease- chemical- or mechanical-related stress, may increase the chances for turl injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turlgrasses are tolerant to this product:

Cool-Season Grasses

Bentgrass, creeping* (Agrosts palustris) Blucgrass, Kentucky (Poa pratensis) Fescue, fine** (Festuca rubra) Fescue, tall (Festuca arundinacea) Ryegrass, perennial (Lolium pere

Warm-Season Grasses Bahlagrass (Paspalum notatum) Bennudagrass (Cynodon dactylon) Buffalograss*** (Buchloe dactyloides) Carpetgrass (Azonopus altinis) Centipodegrass (Eremochton ophiuroides) Kikuyugrass (Pennisetum clandestinum) St. Augustinegrass (Stenolophium) sacundatient) Zoyslagrass (Zoysla japonica)

"Use of this product on certain varieties of creeping bentgrass, such as "Cohansey". 'Carmen', Scaside', and 'Washington' may result in undestrable turtgrass Injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (Agrostis tenuis) varieties

"Use of this product on certain varieties of fine feacue may result in undesirable turf injury. The following fine leacue varieties have been found to be sensitive to this product: 'Alfanta', 'Banner', 'Beauty', 'Bilgant', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf' and

""DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turforase is fully green and has established new roots

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turl management practices before

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of destrable turigrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, trigation and fertilization should be followed. For best results. use mechanical or power seeding equipment (slit seeders) designed to give good seed

Other Use Precautions: The early posteriorgence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before craborass has displayed its fifth leaf or first tiller

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro serification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turks only. DO NOT apply to flowers venerables shrubs or trees

DO NOT use dippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

Preemergence and Early Postemergence Control: This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turis. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabarass, which generally corresponds to the on you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabarass.

Application Frequency and Timing: This product may be applied as a single application, as a splt application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 11 lbs. per 1,000 sq. (t. per application, and no more than 33 lbs. per 1,000 sq. (t. per year,

Spring Applications: For single applications made in the spring of early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turigrass height, and whether the use is considered to be preemergence or early

posteniergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turigrass competitiveness, and the user's

Use Program 1 for preemergence control at sites where the turigrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turigrass is cut low (e.g., golf Gravays), and (b) turigrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence chapters control. This program may also be used for early postemergence control at eiter where furigrass.

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turl maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. is program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where

Where split tertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1: Recommended Application Use Rates* (Lbs. of this product/1,000 sq. ft.)						
PROGRAM	'	2	3			
USE (turigrass cut) REGION	Preemorgence (Nigh)	Preemergence (low) Postemergence (high)	Preemergence (low) Pestemergence (low)			
North	2.8	4.0	5.5			
Transition	4.0	5.5	8.4			
South	5.5	8.4	11.0**			
Coastal South	8.4	11,0**	11.0**			

*DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections shave. Regions include great listed below

North: All other areas

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL. IN, OH, & constal areas of CT, NY, & RI.
South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal greas of AL, GA, LA, MS, NC, SC, & TX.

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control

Tips for Improved Control: For best results, apply this product within a lew days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide imp

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will aboc control of the Program 3' rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barryandgrass, bitsegrass (annual, rough-stalk), crowlooprass, foxfall (yellow & green), goosegrass, kikuyugrass, byegrass (annual) & perennial), smutgrass, lespedeza (common), exalis (buttercup, creeping & yellow wood-sorrel), pursiane (common), epecdwell (corn), spurge (prostrate & epotted). "Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence; bluegrass (annual) - Pop annua, biltercress, chickweed, henbit, geranium (Carolina), parsley-piert, pineappleweed, shepherdspurse.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or organization tall furfiness. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overtaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

SUGGESTED SPREADER SETTINGS

These suggested spreader settings are not intended to replace calibration.

Prease cultrate your spreade: before applying product.

50 lbs. covers 17,800 sq. ft. (0.41 acre) at the 122 lbs. product acre - LIGHT RATE (2.8 lbs.1,000 sq. ft.) 50 lbs, covers 12,500 sq. ft. (0.29 acre) at the 174 lbs, product acre - LOW RATE (4.0 lbs/1,000 sq. ft.) 50 lbs. covers 9,000 sq. ft. (0.21 acre) at the 240 lbs. product/acre - MEDIUM RATE (5.5 lbs/1,000 sq. ft.) 50 lbs. covers 5,900 sq. ft. (0,14 acre) at the 366 lbs. product/acre - HIGH RATE (8.4 lbs/1,000 sq. ft.)

	[SPREADER SETTINGS				
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LIGHT RATE 0.13 lbs. al/A	LOW RATE 0.18 lbs. al/A	MEDIUM RATE 0.25 lbs. al/A	HIGH RATE 0.38 lbs. aVA	
Andersons AccuPro 2000, SR-2000 (cone 4)	3 mph	11 feet	K 1/2	L 1/2	u	N 1/2	
Scotts Rotaries R-8, R-8A (cone 4)	3 mph	11 leet	J	J 3/4	K 1/2	M 16E	
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rom) Pattern II-A	4,5 mph	40 feet	4 1/2	5	5	7 1/2	
Vicon (03 Series)	4,5 mph	35 feet	28	36	42	57.5	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs, and domestic

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environme Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary land!!!l or by incineration, or. if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Magulacturer makes NO OTHER EXPRESS OR IMPLIED. WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAHAS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted

THIS CIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS: YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATEJURISDICTION TO STATEJURISDICTION.

Manufactured by: The Andersons Lawn Fertilizer Division, Inc. P.O. Box 119 Maumee, OH 43537

www.andersonslawnproducts.com

The Andersons logo is a registered trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences LLC.

SGN 215 Net Wt. 50 Lbs. (22.68 kg)

AR (APTRI KPMBRNR)



[&]quot;May require split or sequential applications for full control.