

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

NOV 7 2008

Ms Clair Engelbert The Andersons, Inc P.O.BOX 119 Maumee, Ohio 43537

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request Alternate Brand Name "The Andersons Turf Products Fertilizer

with Dimension Turf Herbicide"

EPA Reg. No. 9198-120

Application Dated September 16, 2008

Dear: Registrant,

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 09/16/20008 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

Manager of the treatment in a				· 5 A		0140				
Please read instructions on p	Environmenta	Inited States		Form Ap		Regi	strati Indme	on	O. Approval expires 2-2 OPP Identifier Number	
		Application	for Pestici	de - Sec	tion	l			1	
1. Company/Product Numbe 9198-120	r	·	2. EPA Miller	Product Man	neger			3. Pro	posed Classification	
4. Company/Product (Name) The Andersons Dimension Herbicide II				PM# 23						
5. Name and Address of App The Andersons Lawn F PO Box 119 Maumee, OH 43537			rt. (b)(i), r to: EPA		is simi	ilar or i	FOL	ific/	FIFRA Section 3(c)(3	
			Section -					"		
Resubmission in resp Notification - Explain Explanation: Use addition Notification of Alternate Brand Na This notification is consistent with made to the labeling or the confident further understand that if this notification of FIFRA and I may be significant to the second secon	nal page(s) if necessar me per PR Notice 98-10: the guidance of PR Notice ential statement of formula ication is not consistent with	y. (For section I Andersons Golf Pro 98-10 and the requ of this product. I ur th the guidance of P	ducts® Fertilizer with , , , , , , , , , , , , , , , , , , ,	egulations at 40 violation of 18 the requiremer	Applica plain be erbicide 0 CFR 15 U.S.C. S	160w. 56.10 and	l to willful	ly make a	ny false statement to EPA.	
			Section -	11					· · · · · · · · · · · · · · · · · · ·	
1. Material This Product Will	Be Packaged In:						_			
child-Resistant Packaging Yes No No			Yes No			2. Ty	P	Netal Yastic Glass		
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per containe	er			Paper Other (S	pecify)	
3. Location of Net Contents Label C 6. Manner in Which Label is	ontainer	4. Size(s) Retai	15 - 2,000 lbs.	V Othe	-	. 1	of Label Label	Direction	ns	
· · · · · · · · · · · · · · · · · · ·	· · · · · · · · · · · · · · · · · · ·	Stencile	Section - I	V						
1. Contact Point (Complete	items directly below	for identification			if nec	esserv	ta proc	ase this	anglication 1	
Name Clair Engelbert		. T	roduct Develop				Т		e No. (include Area Cod	
I certify that the state I acknowledge that an both under applicable	y knowlinglly false or	Certificati this form and a misleading state	ll attachments th	nereto are tru nishable by t	ie, acci	ırate ar impriso	nment o	or CC	6. Date Application Received (Stamped)	

Product Development Technitian

4. Typed Name

5. Date

Clair Engelbert

09/16/08



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

09/16/08

Document Processing Desk (NOTIF)
Office of Pesticide Programs - (7504P)
U. S. Environment Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Subject:

NOTIFICATION - Additional Brand Name

RE:

The Andersons Dimension Herbicide II

EPA Reg. No. 9198-120

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons Turf Products Fertilizer with Dimension Turf Herbicide

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely.

The Andersons Lawn Fertilizer Division, Inc.

Clair Engelbert

Product Development Technician

Phone: 419-891-2780

Fax: 419-891-2745

Email: clair_engelbert@andersonsinc.com

Enclosure



Total Nitrogen (N).

Turf Products

Fertilizer with DIMENSION TO THE PROPERTY OF T

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs

GUARANTEED ANALYSIS

18.00% Urea Nitrogen* Soluble Potash (K ₂ O).	4.00%
Plant nutrients derived from urea, potassium chloride and coated sulfur coated urea. Chlorine (CI) Max	polymer 3.41%
*3.60% Slowly available nitrogen from polymer coated sul coated urea.	lfur F292
ACTIVE INGREDIENT: Dithiopy: 3,5-pyridinedicarbothicic acid, 2-(difluoromethyl)-4- (2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester INGREDIENTS: Total	0.103% 99.897% 100.000%

EPA Reg. No. 9198-120 EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A (Underlined letter is first letter used in run code on end of bag.)

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove confaminated dothing and wash before rouse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals

If in eyes: 1 lold eye open and tinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue missing. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

It swallowed: Call a poison control center or doctor for peatment addice. Have person sip a glass of water if able to sive "c". Do not induce vorniting miless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treat-

Have the product container or label with you with nealing a puison control center or doctor or going for transment. You may also contact 1.800 57-8951 for energency medical treatment shortestion.

EN TIRCTIMENTAL HAZARDS " "

This product is toxic to fish and highly loxic to other equatic organisms including cysters and shrimp. Use with care when applying to furl areas adjacent to any body of water. Drill and nunoil from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean birth water mark. On not aunity when weather conditions taxos chill from treated areas. Do not contaminate water when disposing of equipment washwaters

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on furl being grown for sale or other commercial use as sod, or for commercial seed production, or for research our poses

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 11.0 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course lairways, roughs, and tee boxes. This product will not harm nearby established omamentals when used according to label directions.

USE PRECAUTIONS

Turigrass Safety: This product may be used on seeded, sockled, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good roof system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for furl injury.

This product should only be applied to lawns and ornamental truts that are composed of the following turigrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turigrasses are tolerant to this product:

ConkSeason Grasses

Benigrass, greeping (Agrostis palustris) Bluegrass, Kentucky (Poa pratensis) Fescue, line" (Festuca rubre) Fescue, tall (Festuca arundinacea) Ryegrass, perennial (Lolium perenn

Warm-Season Grasses Bahiagrass (Paspalum notatum coning as a (Fastalum Indiaum)
Bermudaglass (Oxnofon dectyloides)
Bullalograss ("Buchloe dectyloides)
Carpetgrass (Axonopus alfines)
Callipodegrass (Exemochloe ophuroides)
Kituyugrass (Penasetum clandestirum) St. Augustinegrass (Stenotaphrum secundatum) Zoysiagrass (Zoysia japonica)

"Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turtgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentarass (Agrostis tenuis) varieties

"He of this product on certain varieties of fine fescue may result in undesirable furf injury. The following line lescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jarnestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waktort' and 'Waldina' Not all varieties of line tescue have been tested

""DO NOT use this product on seedling bullalograss in the spring of the lirst year of

Note: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and furl management practices before making widespread use.

Reseading, Overseeding, or Sprigging: Reseading, overseeding, or sprigging of neared unity, or was assumed to the region of this product may inhibit be stablishment of describble furgrasses. When research or oversecting, proper cultural practices such as soft cliffully value, in gradient and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed

Other Use Precautions: The early postemergence crabgrass control described below is limited. Posternergence applications are only effective on crabgrass if applied before crabgrass has displayed its filth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core, spike, or bydre aerification, and verticulting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turks only. DO NOT apply to licwers, vegetables, shrubs, or frees

DO NOT use clippings from treated furt for mulching around vegetables or fruit trees. Keep people and pols off treated areas until dust has settled.

APPLICATION DIRECTIONS

Control of Craborass Preemergence and Early Postemergence Control: This product provides "preemer gence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and

It can also provide "early posternergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turts. Posternergence crabgrass control will be obtained only when this product is applied prior to the titlering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turl. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 11 lbs. per 1,000 sq. ft. per application, and no more than 33 lbs. per 1,000 sq. ft. per year

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turigrass mowing height, and whether the use is considered to be preemergence or early posternergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turtgrass competitiveness, and the user's

Use Program 1 for preemergence control at sites where the turigrass is cut-high (e.g., homeowner tawns). This program provides 3-5 months of preemergence crathgrass control. This program should not be used for early postermergence crathgrass control.

Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) trutorass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgiass control. This program may also be used for early postemergence control at sites where turigrass is cul high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turigrass is cut low (e.g., golf fairways), and (b) furl maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early posternergence control at sites where turigrass is cut low (e.g., golf lairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1: Recommended Application Use Rates* (Lbs. of this product/1,000 sq. ft.)						
PROGRAM	1	2	3			
USE (turigrass cut) REGION	Preemergence (high)	Prestantigence (low) - Posternorgence (high)	Preemergence (low) Postemergence (low)			
North	2.8	40	5.5			
Transition	40	55 -	8.4			
South	5.5	e 4	11.0**			
Coastal South	8.4	11.0**	11 0**			

"DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.
South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

"May require split or sequential applications for full control.

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabginss through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season; long control.

Tins for improved Control: For best results, apply this product within a few days after moving and delay moving again for a lew days after the application. When treated favn or ornamental turl areas are watered or receive significant rainfall within a few days after application of this proched, improved weed control may result. Use of splt (half-ared) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Sorting Applications: Used as directed for crabonass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barnyardgrass, bluegrass (annual, alk), crowlootgrass, loxtail (yellow & green)*, goosegrass, (annual & perennial), smutgrass, lespedeza (common), oxalis (buttercup, creeping & yellow woodsorrel), purstane (common), speedwell (corn), spurge (prostrate & spotted) Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for late summer or early fall use, for craburass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: bluegrass (annual) - Poa annua, bittercress, chickweed, henbil, geranium (Carolina), parsley-piert, pineappleweed, shepherdspurse.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or over-laps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

SUGGESTED SPREADER SETTINGS

These suggested spreader settings are not intended to replace calibration Please calibrate your soveader before applying product

50 lbs. covers 17,800 sq: IL (0.41 acre) at the 122 lbs. product/acre - LIGHT RATE (2.8 lbs/1,000 sq. ft.) 50 lbs. covers 12,500 sq. ft. (0.29 scre) at the 174 lbs. productiscre - LOW RATE (4.0 lbs/1,000 sq. ft.) 50 lbs. covers 5,000 sq. ft. (0.21 scre) at the 240 lbs. productiscre - HEDNIM RATE (6.5 lbs/1,000 sq. ft.) 50 lbs. covers 5,500 sq. ft. (0.14 scre) at the 366 lbs. productiscre - HGH RATE (6.4 lbs/1,000 sq. ft.)

			SPREADER SETTINGS					
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LIGHT RATE 0.13 lbs, al/A	LOW RATE 0.18 (bs. aVA	MEDIUM RATE 0.25 (bs. al/A	HIGH RATE 0.38 lbs. aVA		
Andersons AccuPro 2000, SR-2000 (cone 3)	3 mph	11 (ee)	K 1/2	L 1/2	M 1/2	0		
Scotts Rotaries R-8, R-8A (come 3)	3 mph	11 feet	13/4	J 3/4	K 1/2	M 1/4		
Lety Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern 11.48	45 mph	34 feet	5 1/4	6	6 3/4	8		
Vicon (03 Series)	4.5 mph	24 (99)	25	30	36	48		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

STORAGE: Keep container closed and away from food, feedstuffs, and domestic

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess nesticide is a violation of Federal Law. If these wastes cannot be disposed of by pesitions as a wastern or received raw. It ties wastes carried be disposed or by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment Nonretifiable container. Do not reuse or retill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary tandfill or by incineration, or, it allowed by State or local authorities, by burning. It burned, stay out of

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

WARRANTY OF MARCH SING UNITATION OF LOBELT 1: MANUFACTURE WARRANTY OF MERCHANTAGE AND A CONFIDENCE OF THE MERCHANTAGE WARRANTY OF MERCHANTAGE LITY OF FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manulacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to return of purchase price or replacement of product, at Buyor's choice. To the eitent of purchase price or replacement of product, at Buyor's choice. To the eitent consistent with a pricable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of USA

Manufactured by: The Andersons Lawn Fertilizer Division, Inc. P.O. Box 119

www.andersonslawnproducts.com

The Andersons logo is a registered tradernark of The Andersons, Inc.; Dimension is a

SGN 215 Net Wt. 50 Lbs. (22.68 kg)

NOTIFICATION MOV



