3/8/2010



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

8 2010

MAR

Ms. Debbie Ziehr Product Registration The Andersons Lawn Fert. Div., Inc PO Box 119 Maumee, OH 43537

SUBJECT: Application for Pesticide Notification (PRN 98-10) Request Alternate Brand Name "The Andersons Turf Products Fertilizer with 0.103% Dimension Turf Herbicide 13-0-5" EPA Reg. No. 9198-120 Application Dated December 24, 2009

Dear Ms. Ziehr:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 12/24/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

			C	
Environmenta Weshi	United States I Protection Agen ington, DC 20460	cy	Registratio Amendmei ✓ Other	
	Application for Pe			
. Company/Product Number 9198-120		2. EPA Product Mana Miller	3. Proposed Classification	
Company/Product (Name) The Andersons Dithiopyr Herbicide II		РМ# 23		
. Name and Address of Applicant (Include ZIP Co The Andersons Lawn Fert. Div., Inc. J/b/a Free Flow Fert. P.O. Box 119 Maumee, OH 43537	(i t	b)(i), my product is o: EPA Reg. No	similar or identical	with FIFRA Section 3(c)(3) in composition and labeling
	Sectio	Product Name		
Amendment - Explain below. Resubmission in response to Agency letter Notification - Explain below. xplanation: Use additional page(s) if necessar otification of Alternate Brand Name per PR Notice 98-10: his notification is consistent with the guidance of PR Notice ade to the labeling or the confidential statement of formula inther understand that if this notification is not consistent with	 Y. (For section I and Section The Andersons Turf Products F 98-10 and the requirements of I of this product: I understand that 	Agency lette "Me Too" Ap Other - Expla on II.) Fertilizer with 0.103% Dim EPA's regulations at 40 C at it is a violation of 18 U.	ension Turf Herbicide 13-0 FR 156.10 and 40 CFR 15 S.C. Sec. 1001 to willfully	52.46 and no other changes have been make any false statement to EPA, 1
olation of FIFRA and I may be subject to enforcement actio	Sectio			
hild-Resistant Packaging Yes Vo No Certification must Unit Packaging Ves Vo If "Yes" Unit Packaging Ves Vo No	Ve ✓ No No. per If "Yes"	No. per	✓ Pla GI ✓ Pa	tainer stal astic ess per
e submitted	· /	1		her (Specify)
e submitted	4. Size(s) Retail Container 50 lbs 2,00		Location of Label D	
Label	50 lbs 2,00		r . 1	
e submitted Location of Net Contents Information Label V Container	50 lbs 2,00	00 lbs. ✓ Other	On Label	irections
e submitted Location of Net Contents Information Label	50 lbs 2,00	00 lbs. [√] Other n - IV	On Label	irections
e submitted Location of Net Contents Information Label ✓ Container Manner in Which Label is Affixed to Product . Contact Point (Complete items directly below for ame	50 lbs 2,00 Lithograph Paper glued Stenciled Section for identification of individu. Title	00 lbs. [√] Other n - IV	On Label Pocket necessary, to proces Tele	irections
e submitted Location of Net Contents Information Label ✓ Container Manner in Which Label is Affixed to Product . Contact Point (Complete items directly below for ame	50 lbs 2,00 Lithograph Paper glued Stenciled Section for identification of individu. Title Regulatory Certification this form and all attachme	00 lbs. √ Other n - IV al to be contacted, if Administrator ents thereto are true,	On Label Pocket necessary, to process tele 419 accurate and complet	s this application.) phone Nç.º((îrcîtúde Area Code) 891-6671 6. Date Application Recêived °. (Stêmped)
Label ✓ Container . Location of Net Contents Information Label ✓ Container . Manner in Which Label is Affixed to Product . Contact Point (Complete items directly below for lame Debbie Ziehr I certify that the statements I have made on I acknowledge that any knowlinglly false or	50 lbs 2,00 Lithograph Paper glued Stenciled Section for identification of individue Title Regulatory Certification this form and all attachme misleading statement may 3. Title	00 lbs. √ Other n - IV al to be contacted, if Administrator ents thereto are true,	On Label Pocket necessary, to proces Tele c of the accurate and comple accurate and comple	irections s this application.) phone Nບູ.•(ໂກຕໂມde Area Code) -891-6671 6.ເວີຂະອ Application ເຂດ ເຮດ ເຮດ ເຮດ ເຮດ ເຮດ ເຮດ ເຮດ ເຮດ ເຮດ ເຮ



The Andersons, Inc. P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

December 24, 2009

d 2

Document Processing Desk (NOTIF) Office of Pesticide Programs - (7504P) U. S. Environment Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

Subject: **NOTIFICATION - Additional Brand Name**

RE: The Andersons Dithiopyr Herbicide II EPA Reg. No. 9198-120

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons Turf Products Fertilizer with 0.103% Dimension Turf Herbicide 13-0-5

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

Lebbie Ziele

Debbie Ziehr **Regulatory Administrator** Phone #419-891-6671 Fax #419-891-2745 Email: Debbie ziehr@andersonsinc.com

Enclosure

υσιοι ιιιζο

ιςυυς

3



Fertilizer with 0.103% Dimension[®] Turf Herbicide 13-0-5

Contains NutriSphere-N[®]

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs

NOTIFICATION

GUARANTEED ANALYSIS Total Nitrogen (N)	2010
Soluble Potash (K ₂ O).	5.00%
trients derived from urea, potassium chloride.	
Chlorine (CI) Max	4.24%
*2.60% nitrogen stabilized with maleic-itaconic co-polymer	
	F292
ACTIVE INGREDIENT:	
Dithiopyr: 3.5-pyridinedicarbothioic acid; 2-(dituoromethyl)-4- (2-methytpropyl)-6-(nifluoromethyl)-5,5-dimethyl ester	0.103% 99.897%

EPA Reg. No. 9198-120 EPA Est. No. 9198-0H-1M, 9198-0H-2B, 9198-AL-001A

(Underlined letter is first letter used in run codo on end of bag.)

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation. Harmful if inhated. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water atter handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

tled: Move person to fresh air. It person is not breathing, call 911 or an Jance, then give antificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treat ment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information

ENVIRONMENTAL HAZARDS

This product is toxic to lish and highly toxic to other aquatic organisms including cysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runolf from treated turl may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drill from treated areas. Do not contaminate water when disposing of equipment washwaters

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod. or for commercial seed production, or for research purposes,

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 11.0 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year, and is prohibited from use in Nassau and Suffolk Counties

GENERAL INFORMATION

This product is a herbicide that provides control of cratyrass and other annual grasses and broadleaf weeds in established lawins and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established on a mentals when used according to label directions.

USE PRECAUTIONS Turigrass Salety: This product may be used on seeded, soddled, or sprigged lawns and ornamental turis that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two movings following its seed-ing, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornantental turts that are not will established, or on those that have been weakened by weather, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turl injury.

This product should only be applied to lawns and ornamental turts that are composed of the following turbarass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established tudorasses are tolerant to this product:

CoolSesson Brasses Benitgrass, Crooping' (Agrostis pelustris) Bluegrass, Kantucky (Poe pratensis) Fescue, Inti' (Festuce andrea) Fescue, Inti (Festuce andrea) Ryegrass, perenniai (Lolium perenne)	Warn-Season Grasses Barliagrass ("Arepatium notatum) Bermudagrass ("Arendon dectylon) Buttagrass" ("(Buthine acutylondes) Caretigrass (Kanonpos allina) Centipedegrass (Eremoshibe aphrucades) Katuyugrass (Permissum dendestimum) St. Augustinegrass (Stenotaphrum secundas Zoysiagrass (Zoyaia, jepomos)
--	--

'Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen'. 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (Aprostis tenuis) varieties.

"Use of this product on certain variaties of fine (escue may result in undesirable turt injury. The following fine lescue varieties have been found to be sensitive to this product: 'Allanta', Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jameslown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Wakbrt' and 'Waldina'. Not all varieties of line fescue have been tested

***DO NOT use this product on seedling buffatograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: It is recommended that use of this product be made on a trial basis initially to determine its sately under local use conditions and turl management practices before making widespread use

Researing, Overseeding, or Springing: Researing, overseeding, or springing of treated areas within 4 months after application of this product may inhibit the establish ment of desirable turgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass it applied before crabgrass has displayed its fifth leaf or first tiller.

For best results cultural practices that disturb the soil such as core-, spike- or hydro aerification, and verticuting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turts only _DO NOT apply s, vegetables, shrubs, or trees

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled

APPLICATION DIRECTIONS Control of Crabgrass

Preemergence and Early Postemergence Control: This product provides "preemer

gence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established tawns and ornamental turbs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early staties of crationass in well-established laws and ornamental turls. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the fawn or turt. So the practical benefit of this product's additional, early post-emergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass

Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for craborass control in the spring, summer, or fall. DO NOT apply more than 11 lbs. per 1,000 sq. ft. per application, and no more than 33 lbs. per 1,009 sq. II, per year

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turkgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control

provided by this product is decidy related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region

Use Program 1 for preemergence control at sites where the turgenss is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence graburass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turigrass is cut low (e.g., golf (airways), and (b) lurigrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control This program may also be used for early postemergence control at sites where turigrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at siles where (a) turtgrass is cut low (e.g., golf fairways), and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control This program may also be used for early postemergence control at sites where turigrass is cut low (e.g., golt fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired

Where solid fartilizer applications are recommended, the rates in Table 1 may be solid across two applications made 6-10 weeks apart and prior to crabyrass emergence.

PROGRAM	1	2	3 Preemangence (low)	
USE (turfgrass cut)	Preemergenca (high)	Preemengence (iow)		
REGION		Postemergence (high)	Postemergence (low)	
North	2.8	4.0	55	
Transition	4.0	5.5	64	
South	5.5	8.4	11.0**	
Coastal South	8.4	11 0**	11.0**	

"DO NOT apply more than 11 lbs, per application, and no more than 33 lbs, per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below

North: All other areas

nustum)

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX. Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

"May require split or sequential applications for full control

Fall Applications: This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control

Tips for Improved Control: For best results, apply this product within a lew days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turl areas are watered or receive significant raintall within a tew days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the ministo control tarime i rougiani o rates) is appressi tai merioritoriai e assistine I folowing weeks when applied pior to their emergenco: Item mardinas, bluegrass (anuel, roughtalik), nowlootigrass, toxiai (yellow & green), goosegrass, kikuyagrass, ryograss (annual & peneniai), smulgiass, lespedoz (common), oxalis (butteroup, creeping & vellow woodsorreli), purstane (common), speechvell (com), spurge (prostrate & spottech) "Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring. this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: bluegrass (annual) - Poa annua, bittercress, chickweed, henbit, geranium (Carolina), parsley piert, pineappleweed, shepherdspurse.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental jurf area. A more uniform application can be made by spreading hall of the required amount over the area and then applying the remaining hall at a right angle to the previous direction. Avoid streaking, skips, or over-laps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

	ed spinates si e cut order you 0.41 acre) at th 0.29 acre) at th	he 174 lbs. prod	itervied to repla se anplying sco uct/acre - LIGH uct/acre - LOW	dauti TRATE (2.8 lbs RATE (4.0 lbs	1,000 sq. ft.)
	4	J J	SPREADER SETTINGS		
SPREADER		W.DTH OF COTERAGE	E'GHT HATE 0.13 lbs. LVA	1 LOW 1 RATE 1 0.1a lbs.	MEDIUM RATE 0.25 lbs. ai/A
A: Jersons Accurrós:000, SR-2000 (cone 4)	3 mph	12 leel	К 1/2	L 1/2	м
Scolls R-8, R-8A (cone 4)	3 mph	12 feel	1 3/4	J 1/2	K 1/4
Lesco 020093 (pattern 0.5)	3 mph	10 leel	E 1/2	G	1
Lesco 705698 (pattern 0 5)) 3ron ,	- 1991 J	13	16	20
لي بي WTR, WFR, HR, 1250 ب (PTO al 450 rpm) Pallern 11-8	4 5 .ap.i	. છે મધ્ય નૈ	5	6	6 1/2
Vicon (03 Series)	45 mph	36 1691	25	30	36

STORAGE AND DISPOSAL Do not contaminate water, lood, or feed by storage and disposal

STORAGE: Keep container closed and away from lood, feedstuffs, and domestic water supplies

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Nonrelillable container. Do not rouse or refit! this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary tandlill or by incineration, or, it allowed by State or local authorities, by burning. If burned, stay out of smoke,

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liabitity before using this product. It terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to retund of purchase price or replacement of product, at Ruyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted. LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of USA

Manufactured by: The Andersons Lawn Fertilizer Division, Inc. PO Box 119 Maumen OH 43537

www.andersonslawnproducts.com

The Andersons logo is a registered trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences LLC; NutrSphere is a registered trademar Specially Feithizer Hodours, LLC. hark of

SGN 215 Net Wt. 50 Lbs. (22.68 kg)

