



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

April 20, 2017

Debbie Ziehr  
Regulatory Administrator  
The Andersons, Inc.  
1947 Briarfield Blvd.  
Maume, OH 43537

Subject: Notification per PRN 98-10 – Alternate Brand Name  
Product Name: The Andersons Dimension Herbicide II  
EPA Registration Number: 9198-120  
Application Date: February 16, 2017  
Decision Number: 526938

Dear Ms Ziehr:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact Kathryn Montague at 703-305-1243 or via email at [montague.kathryn@epa.gov](mailto:montague.kathryn@epa.gov).

Sincerely,

A handwritten signature in blue ink that reads "Kathryn Montague" with a stylized flourish at the end.

Kathryn Montague, Product Manger 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

# The Andersons Professional Turf™ Products Fertilizer with Dimension® Herbicide 0-0-7

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

GUARANTEED ANALYSIS	
Soluble Potash (K <sub>2</sub> O) .....	7.00%
Plant nutrients derived from potassium chloride.	
Chlorine (Cl) Max .....	5.91%
	<b>F292</b>
<b>ACTIVE INGREDIENT:</b> Dihyopyr, 3,5-pyridinedicarboic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-5-(trifluoromethyl)-S,S-dimethyl ester .....	0.103%
<b>INERT INGREDIENTS:</b> Total .....	99.897%
	100.000%

EPA Reg. No. 9198-120  
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A  
(Underlined letter is first letter used in run code on end of bag.)

**FLORIDA APPLICATIONS:**  
We recommend that you follow the Florida Green Industries BMP's at [http://www.flaes.org/pdf/BMP\\_Book\\_Final.pdf](http://www.flaes.org/pdf/BMP_Book_Final.pdf)  
We recommend that you follow the Florida Golf Course BMP's at [http://www.flaes.org/pdf/golf/BMP\\_Golf\\_Course.pdf](http://www.flaes.org/pdf/golf/BMP_Golf_Course.pdf)

## NOTIFICATION

9198-120

The package is certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated: 04/20/2017

## KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

### FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

**New York Use Restrictions:** In New York State this product may only be used by professional applicators and at no more than 11.0 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

### GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

### USE PRECAUTIONS

**Turfgrass Safety:** This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather, pest, disease, chemical, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

#### Cool-Season Grasses

Bentgrass, creeping\* (*Agrostis palustris*)  
Bluegrass, Kentucky (*Poa pratensis*)  
Fescue, fine\*\* (*Festuca rubra*)  
Fescue, tall (*Festuca arundinacea*)  
Ryegrass, perennial (*Lolium perenne*)

#### Warm-Season Grasses

Bahiagrass (*Paspalum notatum*)  
Bermudagrass (*Cynodon dactylon*)  
Buffalograss\*\*\* (*Buchloe dactyloides*)  
Carpentergrass (*Axonopus affinis*)  
Centipedeagrass (*Eremochloa ophiuroides*)  
Kikuyugrass (*Pennisetum clandestinum*)  
St. Augustinegrass (*Stenotaphrum secundatum*)  
Zoysiagrass (*Zoysia japonica*)

\*Use of this product on certain varieties of creeping bentgrass, such as 'Cohansen', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (*Agrostis tenuis*) varieties.

\*\*Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlantia', 'Banner', 'Beauty', 'Bigard', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Mejenta', 'Mary', 'Pennlawn', 'Tamara', 'Tajana', 'Waldorf' and 'Waldonia'. Not all varieties of fine fescue have been tested.

\*\*\*DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

**Note:** It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

**Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (silt seeders) designed to give good seed to soil contact.

**Other Use Precautions:** The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aeration, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep people and pets off treated areas until dust has settled.

### APPLICATION DIRECTIONS

**Control of Crabgrass**  
**Preemergence and Early Postemergence Control:** This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

**Application Frequency and Timing:** This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 11 lbs. per 1,000 sq. ft. per application, and no more than 33 lbs. per 1,000 sq. ft. per year.

**Spring Applications:** For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

PROGRAM	1		3
	Preemergence (high)	Preemergence (low)	
USE (turfgrass cut)		Preemergence (low)	
			Postemergence (low)
REGION		Postemergence (high)	
			Postemergence (low)
North	2.8	4.0	5.5
Transition	4.0	5.5	8.4
South	5.5	8.4	11.0**
Coastal South	8.4	11.0**	11.0**

\*DO NOT apply more than 11 lbs. per application, and no more than 33 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI

South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

\*\*May require split or sequential applications for full control.

**Fall Applications:** This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

**Tips for Improved Control:** For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

### CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

**Spring Applications:** Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: barnyardgrass, bluegrass (annual, roughstalk), crowfootgrass, foxtail (yellow & green), goosegrass, kikuyugrass, ryegrass (annual & perennial), smutgrass, lespedeza (common), oxalis (buttercup, creeping & yellow woodsores), purslane (common), speedwell (corn), spurge (prostrate & spotted). \*Also controlled at the "Program 2" rates.

**Fall Applications:** Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: bluegrass (annual) - *Poa annua*, bittercress, chickweed, henbit, geranium (Carolina), parsley-piert, pineappleweed, shepherdspurse.

### APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

### SUGGESTED SPREADER SETTINGS

These suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

50 lbs. covers 17,800 sq. ft. (0.41 acre) at the 122 lbs. product/acre - LIGHT RATE (2.8 lbs/1,000 sq. ft.)  
50 lbs. covers 12,500 sq. ft. (0.29 acre) at the 174 lbs. product/acre - LOW RATE (4.0 lbs/1,000 sq. ft.)  
50 lbs. covers 9,000 sq. ft. (0.21 acre) at the 240 lbs. product/acre - MEDIUM RATE (5.5 lbs/1,000 sq. ft.)  
50 lbs. covers 5,900 sq. ft. (0.14 acre) at the 366 lbs. product/acre - HIGH RATE (8.4 lbs/1,000 sq. ft.)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS			
			LIGHT RATE 0.13 lbs. ai/A	LOW RATE 0.18 lbs. ai/A	MEDIUM RATE 0.25 lbs. ai/A	HIGH RATE 0.38 lbs. ai/A
Andersons Model 2000 (cone 4); Andersons 2000 SR (cone 4)	3 mph	12 feet	K 1/2	L 1/2	N 1/2	Q
Scotts R-R, R-8A (cone 4)	3 mph	12 feet	J	K 1/2	M 1/2	P 1/2
Lesco 020093 (pattern 0.25)	3 mph	11 feet	E 1/2	G	I	L
Lesco 705698 (pattern 0.25)	3 mph	11 feet	13	16	20	26
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern III-B	4.5 mph	32 feet	5	6	7	8 3/4
Vicon (03 Series)	4.5 mph	38 feet	23	27	34	50

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**STORAGE:** Keep container closed and away from food, feedstuffs, and domestic water supplies.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

### IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability.

### WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

### LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of USA

Manufactured by:  
The Andersons Lawn Fertilizer Division, Inc.  
P.O. Box 119  
Maumee, OH 43537  
[www.andersonspro.com](http://www.andersonspro.com)

Dimension is a registered trademark of Dow AgroSciences LLC.

**SGN 215**  
**Net Wt. 50 Lbs. (22.68 kg)**



**APT7DM5.1**