9198-121 12-15-2004 Please read instructions on reverse before completing form Form Approved, OMB No. 2070-0060, Approval expire OPP Identifier Number Registration United States **Environmental Protection Agency Amendment** Washington, DC 20460 Other Application for Pesticide - Section I 1. Company/Product Number 2. EPA Product Manager 3. Proposed Classification 9198-121 Miller Restricted None 4. Company/Product (Name) PM# The Andersons Dithiopyr Herbicide III 23 5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) The Andersons Lawn Fertilizer Division, Inc. (b)(i), my product is similar or identical in composition and labeling d/b/a Free Flow Fertilizer to: PO Box 119 EPA Reg. No. Maumee, OH 43537 Check if this is a new address **Product Name** Section - II Amendment - Explain below. Final printed labels in repsonse to Agency letter dated NOTIFICATION "Me Too" Application. Resubmission in response to Agency letter dated ... Notification - Explain below. Other - Explain below. DEC 1 5 2004 Explanation: Use additional page(s) if necessary. (For section I and Section II.) Notification of Minor Label changes per PR Notice 98-10 , addition of optional marketing claims, see cover letter for details This notification is consistent with the provisions of PR Notices 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any take statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. Section - III 1. Material This Product Will Be Packaged In: Child-Resistant Packaging Unit Packaging Water Soluble Packaging 2. Type of Container Yes Yes Metal Yes Plastic No No Nο Glass Paper If "Yes" No. per If "Yes" No. per * Certification must Unit Packaging wgt. container Package wgt container Other (Specify) be submitted 3. Location of Net Contents Information 4. Size(s) Retail Container 5. Location of Label Directions On Label Container Label 6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Section - IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Telephone No. (Include Area Code) Name Regulatory Administrator 334-262-2766 Dawn L. Walters 6. Date Application Certification Received if certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or (Stamped) both under applicable law.

3. Title

5. Date

Regulatory Administrator

11/22/09

Signature

4. Typed Name

Dawn L. Walters

Dun Hillitan

9198-121H Page 1 of 13

(FRONT PANEL) HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

The Andersons Dithiopyr Herbicide III

NOTIFICATION DEC 15 2004

- A Herbicide Plus Fertilizer Combination Product that Provides Control of Crabgrass and Other Weeds in Established Lawns and Ornamental Turfgrasses.
- Premium Granular Fertilizer that Feeds Your Lawn Providing The Nutrients Needed for A Beautiful, Green and Vigorous Lawn.
- Contains Dimension® Herbicide A Herbicide for Crabgrass Control!
- Can Be Applied Up to 4 Weeks Later Than Other Crabgrass Preventers and Controls Crabgrass All Season Long.

Optional for subregistrant:

Controls crabgrass & feeds your lawn

Contains iron for deeper green color

One application controls crabgrass all season long

Stops crabgrass before it starts in established lawns

Contains slow release nitrogen for long-lasting greening

ACTIVE INGREDIENTS:

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(diflu -6-(trifluoromethyl)-S,S- dimethyl ester		0.164%
INERT INGREDIENTS		
	TOTAL	.100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT ____ LBS.

EPA Reg. No. 9198-121 EPA Est. No. 9198-OH-1

Dimension is a registered trademark of Dow AgroSciences LLC

3/15

NET WEIGHT ____ LBS.

The Andersons Dithiopyr Herbicide III

GUARANTEED ANALYSIS	
Total Nitrogen (N)	16%
Available Phosphate (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	4%
Primary Plant Food derived from: Urea, Muriate of Potash, Ammoniated Phospha	tes.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN
CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes	Hold eye open an	nd rinse slowly and	I gently with water	for 15-20 minutes.	Remove
			.44		

contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison

control center or doctor for treatment advice.

If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then

give artificial respiration, preferably mouth-to mouth if possible. Call a poison control

center or doctor for treatment advice.

If swallowed Call a poison control center or doctor for treatment advice. Have person sip a glass

of water if able to swallow. Do not induce vomiting unless told to by a poison control

center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-

20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

4/15

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 7.0 pounds per 1000 square feet (equal to 0.5 pound active ingredient per acre); and is prohibited from use in Nassau and Suffolk Counties.

Use only according to label instructions.

GENERAL INFORMATION

This product is a combination product that provides both crabgrass control and the required nutrients for optimum turfgrass growth all in a single product.

This product contains Dimension, a crabgrass preventer that can be applied up to 4 weeks later than other crabgrass preventers and provide up to 4 months control of crabgrass and other grass and broadleaf weeds prior to their establishment in lawns and ornamental turfgrasses. This product should be applied to lawns and ornamental turfgrasses that are well established (See "Use Precautions" section below for a list of the tolerant turfgrasses). This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

This product should only be applied to lawns and ornamental turfgrasses that are well established. Use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury.

The Andersons Dithiopyr Herbicide

can be applied to the following turfgrasses when applied as directed:

Cool-Season Turfgrasses

- Creeping Bentgrass
- Fine Fescues
- Kentucky Bluegrass
- Perennial Ryegrass
- Tall Fescues

Warm-Season Turfgrasses

- Bahiagrass
- St. Augustinegrass
- Bermudagrass
- Zoysiagrass
- Buffalograss
- Carpetgrass
- Centipedegrass

This product should be applied in the spring or fall prior to the weed seed germination in established lawns and ornamental turfgrasses.

This product can be applied up to 4 weeks later than other crabgrass preventers and still provide season-long weed control of crabgrass. Dimension herbicide provides preemergence control of crabgrass and other weeds when applied prior to their germination, and it also provides postemergence control of crabgrass. The postemergence control provided by this product is limited to crabgrass only and will provide control of crabgrass when applied up to 4 weeks after crabgrass germination. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass.

For control of other grass and broadleaf weeds, this product must be applied as a preemergent treatment. Only crabgrass can be controlled with delayed applications after germination.

Tips for Improved Weed Control: For best results, apply this product within a few days after mowing and delay further mowings after the application of this product until the lawn or ornamental turfgrass has been watered or received significant rainfall to dissolve the fertilizer granules. Improved weed control may be observed when treated lawn or ornamental turfgrass is watered or receives rainfall within a few days after application of this product.

Reseeding, Overseeding, or Sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until 16 weeks after application. Reseeding, overseeding, or sprigging upto 16 weeks after application may prevent the establishment of the newly reseeded, overseeded or sprigged turfgrasses.

(BACK PANEL) HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

5/15

- •DO NOT apply more than 7 pounds of this product per 1,000 square feet of turfgrass area during the year (1,000 square feet area = 100 feet x 10 feet).
- •DO NOT apply to flowers, vegetables, or fruit trees.
- •DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees.
- •DO NOT apply this product to pastures.
- •Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

The Andersons Dithiopyr Herbicide III will provide control of crabgrass and other grasses and broadleaf weeds listed below when applied prior to weed seed germination in established lawns and ornamental turfgrasses.

Weeds Controlled and/or Suppressed by The Andersons Dithiopyr® Herbicide

Grass Weeds:

Annual Bluegrass
Annual Ryegrass
Barnyardgrass
Crabgrass
Crowfootgrass
Goosegrass
Green Foxtail
Kikuyugrass (seedling)
Perennial Ryegrass
Smutgrass (seedling)

Broadleaf Weeds:

Bittercress
Buttercup Oxalis
Carolina Geranium
Chickweed
Common Lespedeza
Common Purslane
Corn Speedwell
Creeping Oxalis
Henbit
Parsley-piert
Pineappleweed
Spotted Spurge
Yellow Woodsorrel

Application Timings:

Yellow Foxtail

This product may be applied either in the spring or in the fall for the control of weeds.

When this product is applied in the fall, a sequential or second application should be made the following spring to provide season-long control of weeds.

Length of Control:

This product will provide up to 4 months of weed control if applied according to label directions.

The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States that this product is being applied.

Please check the map below to determine which "Region" of the U.S. that applies to your particular use area.

Use "Regions" within the United States:



4/15

Recommended Use Rates:

	Recommended Use Rat	tes
	for The Andersons Dithiopyr F	-lerbicide
(Pol	inds of this product to apply per 1,000 s	quare feet of turfgrass)
	Length of V	Weed Control
Region of U.S.	2-3 Months Control	3-4 Months Control
North	1.8	2.5
South	2.7	3.5

SUGGESTED SPREADER SETTINGS

(Available at Final Printing)

Broadcast Spreaders: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

Drop Spreaders: First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

Important: It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. Do not hand spread.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs and domestic water supplies.

DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If party filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL INFORMATION - LANDSCAPE ORNAMENTALS

This product provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces where ornamental plants are grown for aesthetic purposes in ornamental gardens, and residential properties.

Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES

Apply this product with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following table for product recommendations when treating a small area.

DO NOT apply more than 16 pounds per 1,000 square feet per year. Do not apply in Nassau and Suffolk Counties of New York.

Pounds of Product per 1,000 square feet

Ounces of Product per 100 Square Feet 11.2

APPLICATION TIMING AND RECOMMENDATIONS

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS

- Only apply this product to established ornamentals.
- Do not apply this product directly to bare root of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- Do not apply this product on grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing condition. Follow directions given above to determine plant tolerance under your growing conditions prior to large scale use.

COMMON NAME	BOTANICAL NAME	TOLERANT CULTIVARS
Abelia, Dwarf	Abelia X Grandiflora	Nana
Ajuga	Ajuga reptans	Bronze
	Ajunga genevensis	Bronze Beauty
Almond, Flowering	Prunus gladulosa	
Apple*	Malus pumila	
Arbovitae	Thuja occidentalis	Nigra
		Pyramidalis
		Smaragh
		Techny
		Woodwardii
Arborvitae, Golden	Thuja orientalis	
Aster, Chinese	Callistephus chinensis	Dwarf Queen
Ash, Green	Fraxinus pennsylvanica	
Ash, Mountain	Sorbus acucparia	
Ash, Purple	Fraxinus americana	
Azalea	Rhododendron spp.	Brillant
		Buccaneer
		Carror
		Chimes (Belgian)

		Exbury
		Fashion
		Hardijzer Beauty
		Hershey Red
		Higasa
		Hinocrimson
		Holland (Hybrid)
		Marion Lee
		Northern Lights
		Orange Cup
		Orchid Lights
		Snow
		Southern Charm
Azalea, Flame	Rhododendron calendulaceum	
Azalea, Kirishima		· · · · · · · · · · · · · · · · · · ·
Bamboo, Heavenly		
Barberry	Berberis thumbergii	Aurea
		Dwarf Pigmy
		Green
		Kobold _
		Pygmy Red
		Rose Glow
Barberry, Purple		Atropurpurea
Bearberry (common)	Arctostaphylos uva-ursti	Massasschusetts
Birch, River	Betula nigra	
Blueberry*	Vaccinium spp.	Bluecorp
		Blue Jay
		Jersey
		North Blue
		Northland
Bottlebrush	Callistemon citrinum	
Boxwood, Japanese		Japonica
Boxwood, Weller	Buxus sempervirens	
Broom	Cytisus spp.	Moonlight
Broom	Genista pilosa	Vancouver Gold
Bugle Carpet		
Carex, Variegated	Carex	
Camellia	Camellia japonica	Debutante
		Mathotiana Supreme
		Chansonette
	Camellia sasangua	
Cedar, Red	Juniperus virginiana	
Cockscomb, Plumosa	Celosia cristata	Red Kewpie
Coleus	Coleus blumei	, tou tour pro
Copper leaf	Acalypha wilkesiana	
Cotoneaster	Cotonester apiculatus	
Cycads	Cycads revoluta	
Coyotebrush	Baccharis pelularis	
Cyrpress, Bald	Taxodium distichum	
		Clause
Cypress, Italian	Cupressus sempervirens	Glauca
Cypress, Leyland	Cupressocyparis leylandii	—
Cypress, Japanese False	Chamaecyparis obtusa	Gracilis
Daffodil	Narcissus spp.	King Alfred

Daylilly	Hemerocallis spp.	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilson's Yellow
Dogwood	Cornus florida	
Dogwood, American	Cornus sericea	
Douglas Fir	Pseudotsuga menziesii	
Dusty Miller	Senecio cineraria	
Elm, Drake	Ulmus parvifolia	
Euonymus	Euonymus fotunei	Argenteo-variegata Auereo-marginata Colorata Emerald Gaiety Emerald 'N Gold Gold Edge Gold Princess Silver King Tricolor
		Vegetus
Fan Palm, European	Chamaerops humilis	
Fan Palm, Mexican	Washingtonia robusta	
Fescue	Festuca glauca	
Fetterbush	Leucothoe fontiana	Rainbox
Ficus	Ficus retusa	Nitidia
Forsythia	Forsythia X intermedia	Arnold Dwarf Bronxensis Dwarf Lynwood Gold Meadowlark
	Spring Glory	Weeping
Fir Fraser	Abies fraseri	
Gardenia	Gardenia jasminoides	Mystery Radicans
Geranium	Pelargonium X hortorum	
Gum	Eucalyptus citriodora	
Hawthorn	Crataequs spp.	Cockspur White Crimson Cloud Enchantress Jack Evans Washington White
Heather, Twisted	Erica cinerea	Mediterranean Pink
Hibiscus	Hibscus spp.	Brilliant Hula Girl Blue Bird
Hemlock, Canada	Tsuga canadensis	
Holly	llex spp.	Blue Boy Blue Girl Burfodii China Girl Ciompacta Forsteri Hellerie Japanese Northern Beauty
		Needlepoint

10/15

(BACK PANEL) HOMEOWNER USE ONLY (The above is for internal use only and will not appear on the final label copy)

		Nellie R. Stevens
		Savanah
<u> </u>	Ilex X attenuata	
Holly, Chinese	llex cornuta	
Holly, Japanese	llex crenata	
Holly, Yaupon	llex vomitoria	Oleverte Durert
Honeysuckle	Lonicera japonica	Clavey's Dwarf Halliana Tatarian Canadian White Zebelli Red Hosta
Hosta	Hosta spp. Hosta lancifolia	Algo Marginata
Ice Plant	Carpobrotus edulis	
Impatiens	Impatiens wallerana	
Iris	Iris spp.	Wedgewood Dwarf blue
lvy, English	Hedera helix	Bulgaria Thorndale
Jasmine, Asian	Trachelopermum asiaticum	
Juniper	Juniperus spp.	Arcadia Armstrong Bar Harbor
	Juniperus horizontalis	Blue Chip Blue Pacific Blue Rug
	Juniperus chinesis	Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Endora Compacta Fuitlandi Green Gold Tip Hetzi Hughes Manhatten Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blude Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan Tamariscifolia
	Juniperus chinesis	Tamariscifolia
	Juniperus sabina	Weberi Youngstown Yukon Belle

King Palm	Archontophoenix	
Milg Faith	cunninghamiana	
Laurel, Australian	Pittosporum tobira	
Laurel, Mountain	Kalmia latifolia	
Leucothoe	Leucothoe fontanesiana	
Lily, Africian	Agapanthus africanus	Albus
	, igapaining amounas	Peter Pan
Lily, African Blue		
Lily of the Valley	Pieris japonica	Mt. Fire
Lilyturf	Liriope muscari	Evergreen Giant
•	· ·	Lilac Beauty
		Majestic
		Monroe White
		Variegata
Ligustrum, Japanese	Ligustrum japonicum	
Liriope, Green	Liriope spicata	
Maple, Japanese	Acer japonicum	
Maple, Norway	Acer platanoides	
Maple, Red	Acer rubrum	
Maple, Sivler	Acer saccharinum	
Maple, Sugar	Acer saccharum	
Marigold	Tagetes patula	Honeycomb
		Wheeler's Dwarf
		Variegata
Mock Orange*	Philadelphus spp	Golden
		Snowflake Double White
Mopndo Grass	Ophiopogon japonicus	
Moss Rose	Portulaca grandiflora	Sunnyside
Myrtle, Crape	Lagerstoemia indica	Faurei
		Langer
		Muskogee Standard Pink
Myrtle, Wax	Myrica cerifer	Standard Fink
Nadina	Nadina domestica	Composto
INAUINA	ivadina domestica	Compacta Nana
Oleander	Nerium oleander	Hardy Red
Oleander	Neriam oleander	Petite Pink
		Sister Agnes
Oak, Laurel	Quercus laurifolia	
Oak, Pin	Quercus palustris	
Oak, Red	Quercus rubra	
Oak, Southern L	Quercus virginiana	
Oak, Willow	Quercus phellos	
Oteospermum	Osteospermum fruitcosum	Wirligig
Palm, Bangalow	- Colored Colo	
Pampas Grass	Cortadaria selloana	
Paper Flower	Bougainvillea glabra	Barbara Karst
Pachysandra	Pachysandra terminalis	541 5414 13414
Peach*	Prunus persica	
Periwinkle, Dwarf	Vinca minor	
Petunia	Petunia X hybrida	Picoti
Photinia, Red Tip	Photinia X fraseri	1 10011
r nomina, red Tip	T F HOUHIA A HASOH	

Pieris	Pieris japonica	
Pine, Australian	Pinus nigra	
Pine, Japanese Black	Pinus thumbergiana	
Pine, Loblolly	Pinus trambergiana Pinus teada	
Pine, Longleaf	Pinus palustris	
Pine, Mugho	Pinus mugho	
Pine, Scotch	Pinus sylvestris	
Pine, Slash	Pinus eliottii	
Pine, Swiss Mt.	Pinus mugo	
Pine, Virginia	Pinus virginiana	
Pine, White	Pinus strobus	
Pineapple, Guava	Feijoa fellowaiana	
Pittosporum, Japan	1 Oljou lollowalallu	
Potentilla	Potentiall fruticosa	Abbotswood
Privet	Ligustrum japanicum	Golden Vicary Regal Texanum Yellow Tipped Wax
Privet, Glossy	Ligustrum lucidum	
Pyracantha	Pyracantha koidzumii	
Queen Pam	Arecastrum rammanzoffianum	
Quince, Japanese*		
Rhododendron	Rododendron spp	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina	Rhododendron carolinianum	
Rododendron, Catawba	Rhododendron catawbiense	
Rodie Max (Rosebay)	Rhododendron maximum	
Rose	Rosa banksiaae	Luta
Rosemary*	Rosmarinum officinalis	
Rosemary, Bog	Andromeda polifolia	Nana
Salvia	Salvia farinacea	Rhea
Sedum	Sedumm spurium	Dragon Blood Red Red Carpet Yellow
Spiraea	Astilbe X arendsii	Fanall
Spiraea	Spiraea spp.	Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houttee White
Spiraea, Garland	Spiraea spp.	
Spruce, Black Hills	Picea glauca	
Spruce, Colorado Hlue	Picea pungens	
Spruce, Norway	Picea abies	
Spruce, White	Picea glauca	

3/15

(BACK PANEL) HOMEOWNER USE ONLY (The above is for internal use only and will not appear on the final label copy)

Sourwood	Oxydendrum arboreum	
Sweetflag, Grassyy-Le		
Sweetgum	Liquidambar styraciflua	
Sycamore	Platanus occidentalis	
Tree Fern (Tiki Fern)	Asparagus virgatus	
Trumpet Flower, Evening	Gelsemiun sempervirens	
Tulip	Tulip, spp	Apeldoorn
Verbana, Shrub	Lantana sellowiana	
Vervain	Verbena spp.	St. Paul
Viburnum	Viburnum spp.	American Cranberry Bush Arrowood Common Snowball European Cranberry Bush Linden Mohican Wright
Windmill Palm	Trachycarpus fortunei	
Xylosma	Xylosma congestum	
Yaupon	llex vomitoria	Dwarf
Yew	Taxus cuspidatra Taxus X media	Denisiformis

^{*}Ornamental species only: DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS

WEEDS CONTROLLED

When used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established grasses except for crabgrass in early stages of development, and broadleaf weeds, therefore, the area to be treated should be free of weeds prior to application.

GRASSES

barley (Hordeum spp.) barnyardgrass (Echinochloa crus-galli) bluegrass, annual (Poa annua) brome (Bromus spp.) crabgrass, large (Digitaria sanguinalis) crabgrass, smooth (Digitaria ischaemum) crabgrass, Southern (Digitaria ciliaris) crowfootgrass (Dactyloctenium aegyptium) dallisgrass (seedling) (Paspalum dilatatum) goosegrass (Eleusine indica) foxtail, green (Setaria verdi) fotxtail, yellow (Setaria faberi) kikuyugrass (Pennisetum clandestinum) oats, wild (Avena fatua) ryegrass (annual & pernnial) (Lolium spp.) smutgrass (Sporobolus poireti)

BROADLEAF WEEDS

bittercress (Cardamine spp.)

carpetweed (Mollugo verticillata) chickweed (Stellaria spp.) geranium, Carolina (Geranium carolinianum) henbit (Lamium spp.) knotweed, prostrate (Polygonum saviculare) lespedeza, common (Lespedeza striata) marestail (Hippuris vulgaris) medic, black (Medicago lupulina) mustard (Brassica spp.) oxalis, buttercup (Oxalis pes-caprae) pineappleweed (Matricaria matricarioides) pigweed, redroot (Amaranthus retroflexus) parsley-piert (Alchemilla avrensis) pursiane, common (Portulaca oleracea) rocker, London (Sisymbrium irio) shepherdspurse (Capsella bursa-pastoris) speedwell, corn (Veronica arvensis) spurge, garden (Euphorbia hirta) spurge, prostrate (Euphorbia humistrata) spurge, spotted (Euphorbia maculata) woodsorrel, creeping (Oxalis corniculata) woodsorrel, yellow (Oxalis stricta)

14/15

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured by: The Andersons, Lawn Fertilizer Division d/b/a Free Flow Fertilizer P.O. Box 119 Maumee, OH 43537

Optional for subregistrant:

	Recommended Application Timing Guide for Southern States Premium Lawn Products
SSC Product	EarlyeSpring Late Spring > Summer > Fall
Lawn Starter	
Early Spring Lawn Food with Crabgrass Control	
Crabgrass Preventer	
Late Spring Lawn Food with Weed Control	
Broadleaf Weed Control	
Summer Lawn Food with Insect Control	
Lawn Food	
Fall Lawn Food	

Date Printed	11/16/04 notification	
Date Approved	5/8/03	
Supersedes	5/26/98	
Label Changes	Notification - subregistrant optional	
Filename	9198-121h	
Label Code	RVnot111604	



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

November 23, 2004

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7505C)
US Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, DC 20460-0001

RE:

Notification of Minor Label Change

SUBJECT:

The Andersons Dithiopyr Herbicide III

EPA Reg. No. 9198-121

Per PR Notice 98-10 enclosed is an EPA Form 8570-1, an application notifying the Agency of a revision to the homeowner version of the labeling for the subject registration. One copy of the revised labeling is also enclosed with the additions underlined and highlighted in yellow.

We have made the following change by notification:

1. Added optional marketing statements:

Controls crabgrass & feeds your lawn
Contains iron for deeper green color
One application controls crabgrass all season long
Stops crabgrass before it starts in established lawns
Contains slow release nitrogen for long-lasting greening

2. Added optional marketing program as noted on page 13 of label

Enclosed is a card indicating the acceptance or rejection of this notification. Please complete the card and return via mail. If there are any questions or comments regarding this notification, please feel free to contact me.

Sincerely.

The Andersons Lawn Fertilizer Division, Inc.

Daion Studitun

Dawn L. Walters

Regulatory Administrator

Phone 334-262-2766

Fax 334-265-5703

Email dawn_walters@andersonsinc.com

enclosures