



## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

APR 29 2004

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Ms. Dawn L. Walters  
The Andersons Lawn Fert. Div. d/ba Free Flow Fert.  
P.O. Box 119  
Maumee, OH 43537

Dear Ms Walters:

Subject: The Andersons Fertilizer with 0.21% Barricade® Herbicide  
EPA Registration No. 9198-123  
Application and Letter Dated March 19, 2003, Request  
to Amend Registration of Subject Pesticide Product as  
Described in Letter; Application Dated March 24, 2004  
and letter Dated March 25, 2004, Resubmission of Label  
with Revisions for Commercial and Homeowner Labeling

The proposed label amendments as reflected on the labeling  
submitted with your application and letter dated March 24 and  
March 25, 2004, respectively, have been reviewed and found to be  
acceptable under the Federal Insecticide, Fungicide and  
Rodenticide Act, as amended, provided that you:

1. Revise the "Overseeding/Reseeding Table" on page 5 by  
adding under the column "This Product Rate (lb/1,000  
sq.ft.)" the same rates as given in the "Maximum Annual  
Rates Table" (Homeowner Version).
2. Revise the "Spreader Settings" section by listing a  
column for each rate given in the "Maximum Annual Rates  
Table" (Homeowner Version). The rates given are not those  
in the Table. Those rates must be reflected on page  
6 for both "Broadcast Spreader Settings" and "Drop  
Spreader Settings" (Homeowner Version).
3. On page 6, "Landscape Ornamentals", "Use Rates" must be  
revised to reflect those listed in the "Maximum Annual  
Rates Table" on page 4 (Homeowner Version).
4. Submit one copy of the final printed labeling prior to  
shipping under these revised labels.

If these conditions are not complied with, this  
registration will be subject to cancellation in accordance with

2 8 14

FIFRA, section 6(e). Your release for shipment of the product under the revised labeling constitutes acceptance of these conditions.

Stamped copies of the accepted labels, the commercial and the homeowner label, for this product is enclosed for your records.

Sincerely yours,

151

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division  
(7505C)

Enclosures (2 Stamped Labels)

3 8 14

(Front Panel)

# The Andersons Fertilizer with 0.21% Barricade® Herbicide

Active Ingredient:

Prodiamine* .....	0.21%
Other Ingredients: .....	99.79%
Total	100.00%

\*(CAS No. 29091-21-2)

Barricade – trademark of Syngenta Crop Protection

For preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennial & wildflower plantings

*Optional:*

Prevents crabgrass, foxtail, barnyardgrass, goosegrass and other grassy weeds while feeding your lawn with a premium blend of fertilizer that provides quick and extended greening  
 Contains Poly-S® controlled release nitrogen for extended greening  
 Season long control of crabgrass & grassy weeds

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel for additional Precautionary Statements and First Aid

EPA Reg. No. 9198-123

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code

Net Weight  
Covers

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
 APR 29 2004  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act  
 as amended, for the pesticide  
 registered under EPA Reg. No.  
9198-123

4 7 14

(Back Panel)

## The Andersons Fertilizer with 0.21% Barricade® Herbicide

### GUARANTEED ANALYSIS

Total Nitrogen (N) .....	XXX.XX%
Available Phosphorus (K <sub>2</sub> O) .....	XXX.XX%
Soluble Potash (P <sub>2</sub> O <sub>5</sub> ) .....	XXX.XX%
Plant Nutrients Derived From:	

*For products being sold into states which require metals information:*  
 Information regarding the contents and levels of metals in this product is available on the internet at:  
<http://www.regulatory-info-ap.com/>.

*For products containing the micronutrient iron:*  
 Notice: this product contains the micronutrient iron to help prevent iron deficiency. Iron may stain concrete surfaces and should not be applied on water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash this product from concrete surfaces with water.

### PRECAUTIONARY STATEMENTS

#### CAUTION

#### Hazards to Humans and Domestic Animals

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### FIRST AID

- If in eyes            Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
  
- If swallowed        Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
  
- If on skin or clothing    Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
  
- If inhaled            Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

#### Environmental Hazards

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

(Back Panel)

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

*For final printed labels which do include landscape ornamental uses:*

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

*For final printed labels which do not include landscape ornamental uses:*

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

## GENERAL INFORMATION

The Andersons Fertilizer with 0.21% Barricade Herbicide is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in turfgrasses and landscapes. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

## APPLICATION DIRECTIONS


Apply uniformly through suitable, calibrated application equipment.

## ESTABLISHED TURF

This product is a preemergence herbicide that, when properly applied as directed, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

## CRABGRASS GERMINATION DATES



Approx. Date

- 1 - After May 30
- 2 - After May 10
- 3 - After April 20
- 4 - After March 20
- 5 - Jan. 1 to March 20

(Back Panel)

**Use Restrictions and Precautions**

The following precautions apply to the use of this product in turf grasses and lawns:

- Application of this product may thin emerged annual bluegrass and newly over seeded grasses.
- Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer.
- Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- Do not apply to newly set sod until the following year.
- Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
- Do not graze or feed livestock forage cut from areas treated with this product.
- Do not apply this product through any type of irrigation system unless otherwise instructed by this label. Do not apply aerially.
- Do not apply to golf course putting greens.

**Rates of Application**

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds.

This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

**Maximum Annual Rates**

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

**Maximum Annual Rates Table**

<b>Maximum Application Rate of this product per Calendar Year by Turf Grass Species<sup>1</sup></b>	
<b>Turf Species</b>	<b>lbs. product/ 1000 sq. ft.</b>
Bermudagrass <sup>2</sup> , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	12.3
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	8.2
Creeping Red Fescue	6.1
Creeping Bentgrass	5.4

<sup>1</sup>These are the maximum rates per calendar year by species limitations.

<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 4.2 LB./1000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Do not apply more than 12.3 lbs./1,000 sq. ft. of this product per calendar year. Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates table.

**Weeds Controlled**

When used at the recommended rates this product will control:

(Back Panel)

Barnyardgrass	Cupgrass, Woolly	Lambsquarter, Common	Signalgrass, Broadleaf
Bluegrass, Annual ( <i>Poa annua</i> )	Foxtails, Annual	Lovegrass	Speedwell, Persian
Carpetweed	Goosegrass <sup>1</sup>	Oxalis	Sprangletop
Chickweed, Common	Henbit	Panicum	Spurge, Prostrate
Chickweed, Mouseear (from seed)	Itchgrass	(Texas, Fall, Browntop)	Witchgrass
Crabgrass (Large, Smooth)	Johnsongrass (from seed)	Pigweed	Woodsorrel, Yellow (from seed)
Crowfootgrass	Junglerice	Purslane, Common	
	Knotweed	Pusley, Florida	
	Kochia	Rescuegrass <sup>3</sup>	
		Shepherd's Purse <sup>2</sup>	

<sup>1</sup> In many areas a single application of 5.4 to 12.3 lbs./1,000 sq.ft. of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 5.4 to 8.2 lbs./1,000 sq. ft. followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rates Table**. Do not exceed the maximum rate for turf grass species listed in the **Maximum Annual Rates Table** above.

<sup>2</sup> Applications for this weed should be made in late summer, fall, or winter prior to germination. Do not exceed the maximum application rates specified in the Maximum Annual Rates table.

<sup>3</sup> Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum application rates specified in the Maximum Annual Rates table. All applications must be made prior to germination of the weed seeds.

**Overseeding/Reseeding**

This product will inhibit the germination of turf species if overseed/reseeded too soon after application. The interval between application and overseeding/reseeding is dependent on the application rate, soil temperature, and soil moisture. A shorter interval is possible if growing conditions are warm and wet, while a longer interval is needed if conditions are cool and dry.

Use the Overseeding/Reseeding Intervals Table below as a guide when choosing the appropriate overseeding/reseeding interval for each situation.

Overseeding/Reseeding Intervals Table:

This Product Rate (lb/1,000 sq. ft.)	Overseeding/Reseeding Interval (months) Based Upon Seeding Method and Predominant Soil Conditions for the Season			
	Broadcast Seeding		Drill Seeding	
	Wet and Warm	Cool and Dry	Wet and Warm	Cool and Dry
5.7	4	6	3	4
10.9	8	12	6	8
16.4	12	18	9	12

**SPREADER SETTINGS**

*{available at final printing – will vary among subregistrants and market availability of equipment}*

**Broadcast Spreader Settings**

For one pass application: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of wheels will ensure complete coverage.

(Back Panel)

Spreader	7.2 lb. / 1000sq. ft.	8.1 lb. / 1000 sq. ft.	10.9 lb. / 1000 sq. ft.	16.4 lb. / 1000 sq. ft.
aa	X	X	X	X
bb	X	X	X	X
cc	X	X	X	X

**Drop Spreader Settings**

For one pass application: First apply 2 border strips along the edge of each end of the lawn. Then apply products along the length of the lawn, overlapping wheels slightly and shutting off the spreader before turning as you reach the edge of the border strips.

Spreader	7.2 lb. / 1000sq. ft.	8.1 lb. / 1000 sq. ft.	10.9 lb. / 1000 sq. ft.	16.4 lb. / 1000 sq. ft.
aa	X	X	X	X
bb	X	X	X	X
cc	X	X	X	X

It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. Do not apply by hand or with a hand held spreader.

**LANDSCAPE ORNAMENTALS**

This product may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

**Use Rates**

Apply this product at 7.2 to 16.4 lbs./1,000 sq. ft in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 16.4 lbs./1,000 sq. ft..

**Application Timing and Information**

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

**Established Landscape Ornamentals**

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Overhead irrigation or rainfall soon after application will wash granules off plant foliage and activate the product in the soil.

**Newly-Transplanted Landscape Ornamentals**

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before



(Back Panel)

applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

### Tolerant Ornamental Species

This product will not harm trees, shrubs, vines, and flowers listed below. The species listed below in Table 1 are tolerant to this product. This product is also approved for application, except in California, to the species in Table 2, below. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of the ornamental species listed below.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product. Avoid application to plants that are under stress.

**Table 1 - Tolerant Ornamental Species - All States**

Scientific name	Common name
<i>Abies</i> spp.	Fir species** (Balsam, Fraser, Noble, etc.)
<i>Acer palmatum</i>	Japanese Maple
<i>Acer platanoides</i>	Norway Maple
<i>Actinidia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Berberis gladwynensis</i>	Barberry
<i>Berberis julianae</i>	Wintergreen Barberry
<i>Berberis mentorensis</i>	Mentor Barberry
<i>Berberis thunbergii</i>	Japanese Barberry
<i>Berberis verruculosa</i>	Warty Barberry
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisioides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Cleyera japonica</i>	Cleyera
<i>Citrus</i> spp.	Citrus species*
<i>Cornus florida</i>	Flowering Dogwood
<i>Cornus stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>Cotoneaster buxifolius</i>	Cotoneaster
<i>Cotoneaster dammeri</i>	Bearberry Cotoneaster
<i>Cotoneaster microphyllus</i>	Rockspray Cotoneaster
<i>Crataegus</i> spp.	Hawthorne
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Dodonaea viscosa</i>	Hop Bush
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>Euonymus japonica</i>	Japanese Spindle Tree (Evergreen Euonymus)
<i>Euonymus kiautschovicks</i>	Spreading Euonymus

(Back Panel)

<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythia viridissima</i>	Greenstem Forsythia
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine
<i>Gladiolus</i> spp.	Gladiolus species**
<i>Hedera helix</i>	English Ivy
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus Rosa-sinensis</i>	Chinese Hibiscus
<i>Ilex cornuta</i>	Chinese Holly
<i>Ilex crenata</i>	Japanese Holly
<i>Ilex opaca</i>	American Holly
<i>Ilex pernyi</i>	Holly
<i>Ilex vomitoria</i>	Yaupon Holly
<i>Iris</i> spp.	Iris species**
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper
<i>Juniperus conferta</i>	Shore Juniper
<i>Juniperus horizontalis</i>	Creeping Juniper
<i>Juglans</i> spp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum japonicum</i>	Japanese Privet
<i>Ligustrum lucidum</i>	Glossy Privet (Wax-Leaf)
<i>Liriope muscari</i>	Big Blue Lillyturf
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>Lonicera tatarica</i>	Tatarian Honeysuckle
<i>Magnolia</i> spp.	Magnolia species**
<i>Malephora luteola</i>	Ice Plant
<i>Malus</i> spp.	Crabapple*
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus</i> spp.	Narcissus species**
<i>Nerium</i> spp.	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Persea americana</i>	Avocado*
<i>Photinia fraseri</i>	Frasier's Photinia (Redtip)
<i>Picea</i> spp.	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Shrub
<i>Pinus brutia</i>	Calabrian Pine
<i>Pinus canariensis</i>	Canary Island Pine
<i>Pinus elliotii</i>	Slash Pine
<i>Pinus halepensis</i>	Aleppo Pine
<i>Pinus nigra</i>	Austrian Black Pine
<i>Pinus palustris</i>	Longleaf Pine
<i>Pinus radiata</i>	Monterey Pine
<i>Pinus strobus</i>	Eastern White Pine
<i>Pinus sylvestris</i>	Scotch Pine
<i>Pinus taeda</i>	Loblolly Pine
<i>Pinus thunbergiana</i>	Japanese Black Pine
<i>Pinus virginiana</i>	Virginia Pine

(Back Panel)

<i>Pistacia</i> spp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>Pittosporum tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn Scarlet
<i>Pyracantha fortuneana</i>	Firethorn
<i>Pyracantha koidzumii</i>	Firethorn
<i>Pyrus</i> spp.	Bradford Pear spp.
<i>Quercus rubra</i>	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Rhododendron</i>	'Coral Bells'
(including Azalea)	'Formosa'
	'Hino-crimson'
	'PJM'
	'Roseum Elegans'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rumohra adiantiformis</i>	Leatherleaf Fern
<i>Santolina virens</i>	
<i>Sedum album</i>	Stonecrop
<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>Taxus media</i>	Yew
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa</i> spp.	Tulip species
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>Viburnum odoratissimum</i>	Sweet Viburnum
<i>Viburnum plicatum</i>	Japanese Snowball
<i>Viburnum rigidum</i>	Canary Island Viburnum
<i>Viburnum tinus</i>	Laurustinus
<i>Viburnum trilobum</i>	Cranberry Bush
<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Vinca
<i>Vinca minor</i>	Dwarf Periwinkle
<i>Vitis</i> spp.	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet
<i>Yucca filamentosa</i>	Yucca, Adam's Needle

\*Do not use on food producing trees, vines, or plants.

\*\*Not for use on container grown plants.

**Table 2 - Tolerant Ornamental Species/Varieties - All States Except CA**

Scientific name	Common name
<i>Abelia grandiflora</i>	Abelia: Sherwood
<i>Achillea</i> spp.	Yarrow: King Edward
<i>Agapanthus orientalis</i>	

## (Back Panel)

<i>Akebia quinata</i>	Five-Leaf or Chocolate Vine
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia</i> spp.	Aquilegia: Red and Gold
<i>Artemisia</i> spp.	Wormwood; Silver Mound, Castle
<i>Aster</i> spp.	Aster: Bonny Blue, Purple Dome
<i>Aster X frikartii</i>	
<i>Athyrium filix-femina</i>	Lady Fern; Fern Lady
<i>Begonia</i> spp.	Fibrous Begonia: Hardy Grandis
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroides</i>	Snowbank
<i>Bougainvillea</i> spp.	Bougainvillea
<i>Buddleia davidii</i>	Butterfly-Bush (Dwarf Blue); Royal Red
<i>Callistemon citrinus</i>	Crimson Bottlebrush
<i>Campanula carpatica</i>	Tussock Bellflower; (White Clips)
<i>Campis X tagliabuana</i>	Trumpet Creeper, Trumpet Flower; Madame Galen
<i>Ceratostigma plumbaginoides</i>	
<i>Chrysanthemum nipponicum</i>	
<i>Coreopsis</i> spp.	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
	Lucifer
<i>Crocsmia</i> spp.	Cooperi Pink
<i>Delosperma</i> spp.	Larkspur; Blue Elf
<i>Delphinium</i> spp.	Dianthus, Maiden Pinks 'Zing'
<i>Dianthus deltoides</i>	Cheddar Pink
<i>Dianthus gratianopolitanus</i>	Coneflower, Purple; Magnus
<i>Echinacea pupurea</i>	Weeping Forsythia
<i>Forsythia suspensa</i>	Gaillardia, Blanket Flower: 'Goblin'
<i>Gaillardia</i> spp.	
<i>Gaura</i> spp.	
<i>Gentiana daturica</i>	Gentian
<i>Geranium cinereum</i>	Cranesbill
<i>Gypsophila repens</i>	Baby's Breath
<i>Helianthemum</i> spp.	Sunrose
<i>Hemerocallis</i> spp.	Daylily: Aztec Gold, Stella De Oro, Tender Love
<i>Heuchera</i> spp.	Coral Bell; Bridget Bloom
<i>Hibiscus</i> spp.	Mallow; Disco Belle White
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>Hosta sieboldiana</i>	Hosta, 'Searsucker'
<i>Houttuynia cordata</i> var. <i>variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Juniperus davurica</i>	Parsoni
<i>Lagerstromia indica x fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender</i> spp.	Lavender; Munstead
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>Lilium</i> spp.	Lily: Jazz
<i>Liriodendron muscari</i> var. <i>variegata</i>	Liriodendron, Variegated
<i>Liriodendron spicata</i>	Liriodendron, Creeping
<i>Lobelia cardinalis</i>	Cardinal Flower, Indian Pink

(Back Panel)

- |                                 |                                    |
|---------------------------------|------------------------------------|
| <i>Loropetalum chinense</i>     | Burgundy                           |
| <i>Lythrum</i> spp.             | Loosestrife; Modern Pink           |
| <i>Miscanthus sinensis</i>      | Yaku Jima, Silberfeder             |
| <i>Oenothera missouriensis</i>  | Evening Primrose                   |
| <i>Osmanthus heterophyllus</i>  | Osmanthus (False Holly): Gulf Tide |
| <i>Paeonia suffruticosa</i>     | Tree Peony                         |
| <i>Pennisetum setaceum</i>      | Fountain Grass (Dwarf)             |
| <i>Perovskia atriplicifolia</i> |                                    |
| <i>Physostegia virginiana</i>   | Dragonhead, False; Vivid           |
| <i>Quercus shumardii</i>        | Oak, Shumard's Red                 |
| <i>Raphiolepis umbellata</i>    | Yedda Hawthorne                    |
| <i>Rhododendron</i>             | 'Delaware Valley White'            |
| (including azalea)              | 'Flame Creeper'                    |
|                                 | 'Girard Crimson'                   |
|                                 | 'George L. Tabor'                  |
|                                 | Wakelebisu                         |
|                                 | White Gumpo                        |
| <i>Rudbeckia</i> spp.           | Black-Eyed Susan: Goldstrum        |
| <i>Saxifraga</i> spp.           | Saxifrage; Purple Dome             |
| <i>Scabiosa</i> spp.            | Pincushion Flower                  |
| <i>Sedum caudicola</i>          | Stonecrop; Lidakense               |
| <i>Sedum dasyphyllum</i>        | Stonecrop                          |
| <i>Sedum spurium</i>            | Stonecrop; Dragon's Blood          |
| <i>Spiraea bumalda</i>          | Spiraea: Anthony Waterer           |
| <i>Syzyglum paniculatum</i>     | Australian Brushcherry             |
| <i>Teucrium</i> spp.            | Germander                          |
| <i>Thalictrum dipterocarpum</i> | Meadow Rue                         |
| <i>Veronica</i> spp.            | Veronica, Speedwell; Sunny Border  |
| <i>Viburnum suspensum</i>       | Arrowood Viburnum                  |

**STORAGE AND DISPOSAL**

*For homeowner products:*  
Do not contaminate water, food, or feed by storage or disposal.

**Storage:**  
Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

**Disposal:**  
If empty: Do not reuse this container. Place in trash or offer for recycling if available.  
If partly filled: Call your local solid waste agency or 1-800CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**IMPORTANT: READ BEFORE USE:**  
Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER and LIMITATION of LIABILITY:**  
Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY

14 8 14

(Back Panel)

FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:  
THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVamd030804)

Manufactured by:  
The Andersons Lawn Fertilizer Division, Inc.  
d/b/a Free Flow Fertilizer  
PO Box 119  
Maumee, OH 43537

*Optional fertilizer marketing claims:*  
XX% Nitrogen – gives lawns their green color. Fertilizers with controlled-release nitrogen formulas will feed and green your lawn over a longer period of time.  
XX% Phosphate – this nutrient is important for normal root development. Good roots are vital for grass to take up water and nutrients  
XX% Potassium – Also called potash, helps lawns better tolerate extreme weather conditions.  
Contributes to the overall vigor of grass.

Date Printed	3/8/04
Date Approved	
Supersedes	11/5/97
Label changes	Amendment – to match Syngenta's basic label (Barricade F & 65WG)
Filename	9198-123
Label code	RVamd030804