

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
	<b>Application for Pesticide - Section I</b>		
	1. Company/Product Number 9198-124	2. EPA Product Manager Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Barricade Herbicide II	PM# 23	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____  Product Name _____	
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fertilizer Division, Inc. d/b/a Free Flow Fertilizer PO Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address			

**Section - II**

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

**NOTIFICATION**

MAR 28 2003

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of Minor Label change per PR Notice 98-10 : added 4-step program for subregistrant

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Section - III**

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt.    No. per container	If "Yes" Package wgt    No. per container

3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____		

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Dawn L. Walters	Title Regulatory Administrator	Telephone No. (Include Area Code) 334-262-2766
-------------------------	-----------------------------------	---

<p style="text-align: center;"><b>Certification</b></p> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Administrator	
4. Typed Name Dawn L. Walters	5. Date 3/14/03	

2/11

# THE ANDERSONS BARRICADE® HERBICIDE II

Active Ingredient:

Prodiamine\* ..... 0.28%

Inert Ingredients: ..... 99.72%

Total ..... 100.00%

\*CAS No. 29091-21-2

**NOTIFICATION**

**MAR 28 2003**

Guaranteed Fertilizer Analysis:

Nitrogen XX% (N)  
Phosphorous X% (P2O5)  
Potassium X% (K2O)

Nitrogen from: XXXX

For selective preemergence control of grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns and landscape ornamentals (including perennial & wildflower plantings)

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See the following section, Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

Net Weight:

EPA Reg. No. 9198-124  
EPA Est. No.

Barricade® is a Registered Trademark of Syngenta Crop Protection

3/11

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

#### Statements of Practical Treatment (or First Aid)

If in eyes: Flush thoroughly with water for several minutes. Get medical attention if irritation persists.

If on skin: Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists.

If inhaled: Remove victim to fresh air.

#### Environmental Hazards

This product has low solubility in water. At the time of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

##### STORAGE

Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

##### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

##### CONTAINER DISPOSAL

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or; puncture and dispose of in a sanitary landfill or at an incineration facility, or; if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

Paper and plastic containers: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or at an incineration facility, or; if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

Fiber Drums: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residues into application equipment. Dispose of empty liner in a sanitary landfill or at an incineration facility if allowed by state and local authorities. If the drum cannot be reused, dispose of in a sanitary landfill or by incineration if allowed by state and local authorities.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

4/11

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in turf grasses and landscapes. This product controls susceptible weeds by inhibiting weed seeds germination and root development. For most effective weed control this product must be incorporated by 0.5 inch of rainfall, irrigation or shallow mechanical incorporation. Incorporation should take place as soon as possible after application and should not be delayed past 14 days from the date of application.

*For final printed labels which do include ornamental uses:*

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

*For final printed labels which do not include ornamental uses:*

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aurally.

Do not apply to golf course putting greens.


**Application Directions**

Apply uniformly through suitable, calibrated commercial application equipment.

**ESTABLISHED TURF**

This product is a selective preemergence herbicide that when properly applied, will control certain grass and broadleaf weeds in established turf grass and lawns. The species of turf grass determines the maximum amount of material that may be applied during a year.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation within 14 day after application and prior to weed germination. See the map below for approximate crabgrass germination dates.



**Use Precautions**

The following use precautions apply to the use of this product in turf grasses and lawns.

Application of this product may thin emerged annual bluegrass and newly over-seeded grasses.

Do not apply to OVERSEEDED TURF for 60 days after seeding or until after the second mowing, which ever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Do not apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

**Maximum Annual Rates**

This product should not be applied above a maximum annual rate of 536 lb/A.

**Turf Species and Rates**

For season long control this product may be applied as a single application, or if needed followed by a second application for increased length of control. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rate within the listed range for longer control. Longer control is desirable in areas where crabgrass germinates between January 1 and April 20 (See Crabgrass Seed Germination Data Map).

Select maximum use rate based on turf grass species. All applications must be made prior to germination of the target weed seeds. This product will not control established weeds.

**Turf Species and Rates Table:**

MAXIMUM RATE/CALENDAR YEAR Of THE ANDERSONS BARRICADE HERBICIE II by Turf Grass Species <sup>1</sup>		
Turf Species:	Lb. ANDERSONS BARRICADE II /A	lb ai/A
Bermudagrass <sup>2</sup> Bahagrass Centipedegrass Seashore Paspalum Tall Fescue (including turf-type) Zoysia	536	1.5
Buffalograss Kentucky Bluegrass Perennial Ryegrass	357	1.0
Creeping Red Fescue St. Augustinegrass	264	0.75
Creeping Bentgrass	236	0.65

<sup>1</sup> These are the maximum rate per calendar year by species limitations.

<sup>2</sup> May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 186 lb/A. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

6/11

- Do not use more than 536 lb. of this product per calendar year.
- Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type.

**Weeds Controlled**

When used at the recommended rates (see Turf Species and Rates Table) this product will control:

- |  |                                |
|--|--------------------------------|
| Barnyardgrass                          | Carpetweed                     |
| Bluegrass, Annual ( <i>Poa annua</i> ) | Chickweed, Common              |
| Crabgrass (large, smooth)              | Henbit                         |
| Foxtails, Annual                       | Knotweed                       |
| Goosegrass <sup>1</sup>                | Lambsquarter, Common           |
| Johnsongrass (from seed)               | Oxalis                         |
| Panicum                                | Pigweed                        |
| Rescuegrass <sup>3</sup>               | Purslane, Common               |
| Signalgrass, Broadleaf                 | Shepard's purse <sup>2</sup>   |
| Spurge, Prostrate                      | Woodsorrel, Yellow (from seed) |

Do not exceed the maximum annual application rates recommended for the turf species. All applications must be made prior to germination of weed seeds. This product will not control established weeds.

<sup>1</sup> In many areas a single application of 236 to 536 lb/A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, make an initial application of 236 to 357 lb/A followed after 60 to 90 days by a second application of 186 to 300 lb/A, respectively, for season long goosegrass control. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.

<sup>2</sup> Winter weeds are best controlled by applications made prior to germination: later summer, fall or winter timings.

<sup>3</sup> Suppression only.

**Overseeding/Reseeding**

This product will inhibit the germination of turf species if overseed/reseeded too soon after application. The interval between application and overseeding/reseeding is dependent on the application rate, soil temperature, and soil moisture. A shorter interval is possible if growing conditions are warm and wet, while a longer interval is needed if conditions are cool and dry.

Use the Overseeding/Reseeding Intervals Table below as a guide when choosing the appropriate overseeding/reseeding interval for each situation.

Overseeding/Reseeding Intervals Table:

This Product Rate (lb/A)	Overseeding/Reseeding Interval (months) Based Upon Seeding Method and Predominant Soil Conditions for the Season			
	Broadcast Seeding		Drill Seeding	
	Wet and Warm	Cool and Dry	Wet and Warm	Cool and Dry
186	4	6	3	4
357	8	12	6	8
536	12	18	9	12

7/11

---

**LANDSCAPE ORNAMENTALS** (including Perennial and Wildflower Plantings)

---

This product may be applied for residual preemergence weed control in ornamentals.

**USE RATES**

Apply this product at 236 to 536 lbs./A in fall and/or spring. Use the higher rate of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 536 lbs./A.

**APPLICATION TIMING AND INFORMATION**

Apply this product prior to weed seed germination to the soil surface surrounding established tolerant ornamentals listed below. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established.

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, before weed seeds germinate. Acceptable control will be obtained when this product is activated within 14 days after application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of The Andersons Fertilizer with 0.28% Barricade Herbicide, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

**Tolerant Ornamental Species**

This product will not harm established ornamental trees, shrubs, and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of the listed species.

<b>Scientific name</b>	<b>Common name</b>
<i>Acer palmatum</i>	Japanese Maple
<i>A. platanoides</i>	Norway Maple
<i>Actinidia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Berberis gladwynensis</i>	Barberry
<i>Berberis julianae</i>	Wintergreen Barberry
<i>Berberis mentorensis</i>	Mentor Barberry
<i>Berberis thunbergii</i>	Japanese Barberry
<i>Berberis verruculosa</i>	Warty Barberry
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisioides</i>	Feathery Cassia

<i>Ceanothus rigidus</i>	Wild Lilac
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Cleyera japonica</i>	Cleyera
<i>Citrus</i> spp.	Citrus species*
<i>Cornus florida</i>	Flowering Dogwood
<i>Cornus stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>Cotoneaster buxifolius</i>	Cotoneaster
<i>Cotoneaster dammeri</i>	Bearberry Cotoneaster
<i>Cotoneaster microphyllus</i>	Rockspray Cotoneaster
<i>Crataegus</i> spp.	Hawthorne*
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Dodonaea viscosa</i>	Hop Bush
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>Euonymus japonica</i>	Evergreen Euonymus
<i>Euonymus kiautschovicks</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythia viridissima</i>	Greenstem Forsythia
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine
<i>Hedera helix</i>	English Ivy
<i>Hibiscus Rosa-sinensis</i>	Chinese Hibiscus
<i>Ilex cornuta</i>	Chinese Holly
<i>Ilex crenata</i>	Japanese Holly
<i>Ilex opaca</i>	American Holly
<i>Ilex pernyi</i>	Holly
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper
<i>Juniperus conferta</i>	Shore Juniper
<i>Juniperus horizontalis</i>	Creeping Juniper
<i>Juglans</i> spp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crepe Myrtle
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum japonicum</i>	Japanese Privet
<i>Ligustrum lucidum</i>	Glossy Privet (Wax-Leaf)
<i>Liriodendron muscari</i>	Big Blue Lillyturf
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>Lonicera tatarica</i>	Tatarian Honeysuckle
<i>Malephora luteola</i>	Ice Plant
<i>Malus</i> spp.	Crabapple*
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Persea americana</i>	Avocado*
<i>Photinia fraseri</i>	Frasier's Photinia (Redtip)
<i>Picea abies</i>	Norway Spruce
<i>Pieris japonica</i>	Lily-of-the-Valley Shrub
<i>Pinus brutia</i>	Calabrian Pine
<i>Pinus canariensis</i>	Canary Island Pine
<i>Pinus halepensis</i>	Aleppo Pine

<i>Pinus nigra</i>	Austrian Black Pine
<i>Pinus radiata</i>	Monterey Pine
<i>Pinus thunbergiana</i>	Japanese Black Pine
<i>Pinus strobus</i>	Eastern White Pine
<i>Pinus sylvestris</i>	Scotch Pine
<i>Pistacia</i> spp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>Pittosporum tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pyracantha coccinea</i>	Firethorn
<i>Pyracantha fortuneana</i>	Firethorn
<i>Pyracantha koidzumii</i>	Firethorn
<i>Pyrus</i> spp.	Bradford Pear spp.
<i>Quercus rubra</i>	Red Oak
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Rhododendron</i> (including Azalea)	'Coral Bells' 'Formosa' 'Hino-crimson' 'PJM' 'Roseum Elegans'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Santolina virens</i>	
<i>Sedum album</i>	Stonecrop
<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>Taxus media</i>	Yew
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>Viburnum odoratissimum</i>	Sweet Viburnum
<i>Viburnum plicatum</i>	Japanese Snowball
<i>Viburnum rigidum</i>	Canary Island Viburnum
<i>Viburnum tinus</i>	Laurustinus
<i>Viburnum trilobum</i>	Cranberry Bush
<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Vinca minor</i>	Dwarf Periwinkle
<i>Vitis</i> spp.	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet
<i>Yucca filamentosa</i>	Yucca, Adam's Needle

\*Ornamental species only. Do not use on food producing trees or vines.

Manufactured by:  
The Andersons Lawn Fertilizer Division, Inc.  
d/b/a Free Flow Fertilizer  
PO Box 119  
Maumee, OH 43537

**Notification dated 3/14/03:**

*Optional marketing statements for subregistrant:*

**Prevents crabgrass before it starts and kills germinating seeds**

Triple Action Feed works at every level  
Grass Greener, Thicker Lawn  
Roots Stronger Root Systems  
Soil Enriches & Fortifies Soil

100% Satisfaction Bayer {logo} A Better Lawn Guaranteed! Or your money back 100% Satisfaction

Bayer Advanced™ Triple Action Lawn Fertilizer PLUS Crabgrass Preventer<sup>II</sup> stops crabgrass, foxtail, spurge, and oxalis while it nourished your lawn. Its Triple Action formula contains slow-release nitrogen which continuously feeds the grass, roots and soil for up to two full months. And it's guaranteed not to burn. The result is a greener, thicker lawn. You're just one application away from a better lawn... That's the Bayer Better Lawn Guarantee.

Triple Action Feeding works at every level for a Deep-Down Green!  
Why is Triple Action better? By feeding at all levels, your lawn received the primary nutrients it needs to thrive. This ensures thicker, greener grass, stronger root systems, and richer soil – the keys to building a lush lawn that lasts.

- Grass Builds thicker, greener grass and enhances overall appearance.
- Roots Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help your lawn withstand heat-stress and drought.
- Soil Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.

Effective control of crabgrass PLUS other listed broadleaf weeds  
Non burning formula

**4-Step Annual Program**

<u>Apply Early Spring</u> Feb. – April	<u>Apply Late Spring</u> April – June	<u>Apply Summer</u> June – Aug.	<u>Apply Fall</u> Oct. – Nov.
<u>(Diagram of crabgrass preventer product)</u>	<u>(Diagram of weed &amp; feed product)</u>	<u>(Diagram of fertilizer product)</u>	<u>(Diagram of winterizer product)</u>
<u>Stops crabgrass</u>	<u>Kills weeds</u>	<u>Revitalizes lawn</u>	<u>Boosts root development</u>
<u>Feeds lawn</u>	<u>Feeds lawn</u>	<u>Fights heat stress</u>	<u>Improves spring green-up</u>

Bayer Advanced Lawn® Crabgrass Preventer<sup>II</sup> stops crabgrass, foxtail, spurge, oxalis and other listed weeds before they start and kills germinating seeds. These convenient granules provide long-lasting protection and are guaranteed not to burn. You're just one application away from a beautiful, weed-free lawn... That's the Bayer Better Lawn Guarantee

Bayer Advanced Lawn® Crabgrass Preventer<sup>II</sup> stops weeds before they start and kills germinating seeds with just one application. These specially formulation granules provide uniform coverage for improved weed control. Watering after application releases the weed killer and forms a protective barrier to stop new weeds growing from seed. Effective, long-lasting control is provided against crabgrass, foxtail, and spurge as well as other listed grassy and broadleaf weeds.

Questions, Comments, Information? 1-877-XXX-XXXX

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price

Made in the USA



**The Andersons, Inc.**

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

March 14, 2003

Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7505C)  
US Environmental Protection Agency  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, DC 20460-0001

RE: Notification of Minor Label Change

SUBJECT: The Andersons Barricade Herbicide II  
EPA Reg. No. 9198-124

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency of a revision to the labeling for the subject registration. One copy of the revised labeling is enclosed. The changes are underlined and marked in bold, yellow highlight for additions.

We have made the following changes by notification:

1. added 4-step program for subregistrant

Enclosed is a card requesting the acceptance or rejection of this notification. Please complete the card and return via mail.

If there are any questions or comments regarding this notification, please feel free to contact me.

Sincerely,  
The Andersons Lawn Fertilizer Division, Inc.

*Dawn L. Walters*

Dawn L. Walters  
Regulatory Administrator  
Phone 334-262-2766  
Fax 334-265-5703  
Email dawn\_walters@andersonsinc.com

enclosures