

9198-124

03/10/2005

1/10

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 9198-124	2. EPA Product Manager Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Barricade Herbicide II	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fertilizer Division, Inc. d/b/a Free Flow Fertilizer PO Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION MAR 10 2005
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Label Change change per PR Notice 98-10 : see cover letter for details

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label	
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Dawn L. Walters	Title Regulatory Administrator	Telephone No. (include Area Code) 334-262-2700
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment both under applicable law.		6. Date Application Received (Stamped)
2. Signature Dawn Walters	3. Title Regulatory Administrator	
4. Typed Name Dawn L. Walters	5. Date 2/15/05	



The Andersons, Inc.
P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

2/10

Alternate Brand Notification dated 2/15/05 for EPA Reg. No. 9198-124:

This notification is consistent with the provisions of PR Notices 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

The Andersons Lawn Fertilizer Division, Inc.

Dawn Walters

Dawn L. Walters
Regulatory Administrator

NOTIFICATION
MAR 10 2005

3/10

(Front Panel)

The Andersons Fertilizer with 0.28% Barricade[®] Herbicide

Active Ingredient:

Prodiamine*	0.28%
Other Ingredients:	<u>99.72%</u>
Total	100.00%

*(CAS No. 29091-21-2)

Barricade – trademark of Syngenta Crop Protection

For preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennial & wildflower plantings

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements and First Aid

EPA Reg. No. 9198-124

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code

Net Weight

Covers

4/16

(Back Panel)

The Andersons Fertilizer with 0.28% Barricade® Herbicide

GUARANTEED ANALYSIS

Total Nitrogen (N) XXX.XX%
Available Phosphorus (K₂O) XXX.XX%
Soluble Potash (P₂O₅) XXX.XX%
Plant Nutrients Derived From:

For products being sold into states which require metals information:
Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.regulatory-info-ap.com/>.

For products containing the micronutrient iron:
Notice: this product contains the micronutrient iron to help prevent iron deficiency. Iron may stain concrete surfaces and should not be applied on water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash this product from concrete surfaces with water.

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

- If in eyes Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
- If swallowed Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- If on skin or clothing Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
- If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

Environmental Hazards

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas. Drift and runoff from

5/10

(Back Panel)

treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For final printed labels which do include landscape ornamental uses:

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

For final printed labels which do not include landscape ornamental uses:

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

GENERAL INFORMATION

The Andersons Fertilizer with 0.28% Barricade Herbicide is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in turfgrasses and landscapes. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF


This product is a preemergence herbicide that, when properly applied as directed, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

6/16

(Back Panel)

CRABGRASS GERMINATION DATES



Approx. Date

- 1 - After May 30
- 2 - After May 10
- 3 - After April 20
- 4 - After March 20
- 5 - Jan. 1 to March 20

Use Restrictions and Precautions

The following precautions apply to the use of this product in turf grasses and lawns:

- Application of this product may thin emerged annual bluegrass and newly over seeded grasses.
- Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer.
- Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- Do not apply to newly set sod until the following year.
- Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
- Do not graze or feed livestock forage cut from areas treated with this product.
- Do not apply this product through any type of irrigation system unless otherwise instructed by this label.
- Do not apply aerially.
- Do not apply to golf course putting greens.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Annual Rates Table

Maximum Application Rate of this product per Calendar Year by Turf Grass Species ¹

7/10

(Back Panel)

Turf Species	lbs. A/A	lbs. product /A	lbs. product/ 1000 sq. ft.
Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	1.5	536	12.3
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	1.0	357	8.2
Creeping Red Fescue	0.75	264	6.1
Creeping Bentgrass	0.65	236	5.4

¹ These are the maximum rates per calendar year by species limitations.

² May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 186 lb./acre or 4.2 LB./1000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Do not apply more than 536 lbs. per acre (12.3 lbs./1,000 sq. ft.) of this product per calendar year. Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates table.

Weeds Controlled

When used at the recommended rates this product will control:

Barnyardgrass	Cupgrass, Woolly	Lambsquarter, Common	Signalgrass, Broadleaf
Bluegrass, Annual (<i>Poa annua</i>)	Foxtails, Annual	Lovegrass	Speedwell, Persian
Carpetweed	Goosegrass ¹	Oxalis	Sprangletop
Chickweed, Common	Henbit	Panicum	Spurge, Prostrate
Chickweed, Mouseear (from seed)	Itchgrass	(Texas, Fall, Browntop)	Witchgrass
Crabgrass	Johnsongrass	Pigweed	Woodsorrel, Yellow (from seed)
(Large, Smooth)	(from seed)	Purslane, Common	
Crowfootgrass	Junglerice	Pusley, Florida	
	Knotweed	Rescuegrass ³	
	Kochia	Shepherd's Purse ²	

¹ In many areas a single application of 5.4 to 12.3 lbs./1,000 sq.ft. (236 to 536 lbs./A) of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 5.4 to 8.2 lbs./1,000 sq. ft. (236 to 357 lbs./A) followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rates Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rates Table above.

² Applications for this weed should be made in late summer, fall, or winter prior to germination. Do not exceed the maximum application rates specified in the Maximum Annual Rates table.

³ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum application rates specified in the Maximum Annual Rates table. All applications must be made prior to germination of the weed seeds.

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When to Overseed After Application – All States*

8/16

(Back Panel)

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

The Andersons Fertilizer with 0.28% Barricade Herbicide Rate			Interval Before Overseeding		
lbs/1000 sq. ft	lbs./A	ai/A	North	Transition	South
4.1	179	0.50	4 mo	4 mo	4 mo
5.4	236	0.65	5 mo	4 mo	4 mo
6.1	264	0.75	6 mo	5 mo	5 mo
6.6	289	0.81	---	6 mo	6 mo
8.2	357	1.00	---	7 mo	7 mo
9.4	411	1.15	---	---	9 mo
10.6	464	1.30	---	---	10 mo
12.3	536	1.50	---	---	12 mo

*Note: In AZ, CA, NV and TX, the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section " *Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NX and TX only)".

***Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NX and TX only)**

Use on golf courses (excluding golf course putting greens) and lawns when overseeding with perennial ryegrass (minimum seed rate of 350 lbs./A).

How Much and When to Apply

Amount to Apply		When to Apply	Expected Control	Use Precautions
Lbs./Acre	Lbs/1000			
135 - 236	3.1 - 5.4	6 to 8 weeks before ryegrass overseeding Second application 4-8 weeks after overseeding date or when perennial ryegrass roots are in the second inch of soil	One application for 70% or greater control of <i>Poa annua</i> Second application should enhance the residual under normal weather conditions	<ol style="list-style-type: none"> Some seed mortality and temporary reduction in root growth of new seedlings may occur To reduce the potential for seeding mortality, maintain a moist seedbed with light, frequent irrigation. Make no more than 2 applications per year for this use, and do not exceed 302 lbs./A (6.9 lbs/1000) for these two applications for <i>Poa annua</i> control in ryegrass overseeded turf Do not make a second application if any injury to the ryegrass is observed after the first application Do not make a second application unless the product was first applied before overseeding

(Back Panel)

Recommended Spreader Settings

These suggested spreader settings are not intended to replace calibration.
Please calibrate your spreader before applying product.

- ___ lbs. treats ___ ft² at the 236 lbs. product/acre – LIGHT RATE (5.4 lbs./1,000 sq. ft.)
- ___ lbs. treats ___ ft² at the 264 lbs. product/acre – LOW RATE (6.1 lbs./1,000 sq. ft.)
- ___ lbs. treats ___ ft² at the 357 lbs. product/acre – MEDIUM RATE (8.2 lbs./1,000 sq. ft.)
- ___ lbs. treats ___ ft² at the 536 lbs. product/acre – HIGH RATE (12.3 lbs./1,000 sq. ft.)

Spreader	Ground or PTO Speed	Width of Coverage	Spreader Settings			
			Light Rate	Low Rate	Medium Rate	High Rate
aa	X mph	XX ft/	X	X	X	X
bb	X mph	XX ft	X	X	X	X
cc	X mph	XX ft	X	X	X	X

LANDSCAPE ORNAMENTALS

This product may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

Use Rates

Apply this product at 236 to 536 lbs./A (5.4 to 12.3 lbs./1,000 sq. ft) in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 536 lbs./A (12.3 lbs./1,000 sq. ft.).

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Overhead irrigation or rainfall soon after application will wash granules off plant foliage and activate the product in the soil.

Newly-Transplanted Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

10/16

(Back Panel)

Tolerant Ornamental Species

This product will not harm trees, shrubs, vines, and flowers listed below. The species listed below in Table 1 are tolerant to this product. This product is also approved for application, except in California, to the species in Table 2, below. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of the ornamental species listed below.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product. Avoid application to plants that are under stress.

Table 1 - Tolerant Ornamental Species - All States

Scientific name	Common name
<i>Abies</i> spp.	Fir species** (Balsam, Fraser, Noble, etc.)
<i>Acer palmatum</i>	Japanese Maple
<i>Acer platanoides</i>	Norway Maple
<i>Actinidia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Berberis gladwynensis</i>	Barberry
<i>Berberis julianae</i>	Wintergreen Barberry
<i>Berberis mentorensis</i>	Mentor Barberry
<i>Berberis thunbergii</i>	Japanese Barberry
<i>Berberis verruculosa</i>	Warty Barberry
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisioides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Cleyera japonica</i>	Cleyera
<i>Citrus</i> spp.	Citrus species*
<i>Cornus florida</i>	Flowering Dogwood
<i>Cornus stolonifera</i>	American Dogwood
<i>Cortaderia seloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>Cotoneaster buxifolius</i>	Cotoneaster
<i>Cotoneaster dammeri</i>	Bearberry Cotoneaster
<i>Cotoneaster microphyllus</i>	Rockspray Cotoneaster
<i>Crataegus</i> spp.	Hawthorne
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Dodonaea viscosa</i>	Hop Bush
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>Euonymus japonica</i>	Japanese Spindle Tree (Evergreen Euonymus)
<i>Euonymus kiautschovicks</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia

(Back Panel)

Forsythia intermedia
Forsythia viridissima
Gardenia jasminoides
Gladiolus spp.
Hedera helix
Hibiscus
Hibiscus Rosa-sinensis
Ilex cornuta
Ilex crenata
Ilex opaca
Ilex pernyi
Ilex vomitoria
Iris spp.
Jasminium nudiflorum
Juniperus chinensis
Juniperus conferta
Juniperus horizontalis
Juglans spp.
Justicia brandegeana
Lagerstromia indica
Ligustrum amurense
Ligustrum japonicum
Ligustrum lucidum
Liriodendron muscari
Lonicera japonica
Lonicera tatarica
Magnolia spp.
Malephora luteola
Malus spp.
Nandina domestica
Narcissus spp.
Nerium spp.
Olea europaea
Ophiopogon japonicus
Osteospermum fruticosum
Oxydendrum arboreum
Persea americana
Photinia fraseri
Picea spp.
Pieris japonica
Pinus brutia
Pinus canariensis
Pinus elliotii
Pinus halepensis
Pinus nigra
Pinus palustris
Pinus radiata
Pinus strobus
Pinus sylvestris
Pinus taeda
Pinus thunbergiana
Pinus virginiana

Border Forsythia
 Greenstem Forsythia
 Gardenia, Cape-Jasmine
Gladiolus species**
 English Ivy
 Rose of Sharon**
 Chinese Hibiscus
 Chinese Holly
 Japanese Holly
 American Holly
 Holly
 Yaupon Holly
Iris species**
 Winter Jasmine
 Chinese Juniper
 Shore Juniper
 Creeping Juniper
 Walnut*
 Shrimp Plant
 Crape Myrtle
 Amur Privet
 Japanese Privet
 Glossy Privet (Wax-Leaf)
 Big Blue Lillyturf
 Japanese Honeysuckle
 Tatarian Honeysuckle
Magnolia species**
 Ice Plant
 Crabapple*
 Heavenly Bamboo
Narcissus species**
 Oleander
 Olive*
 Mondo Grass
 Trailing African Daisy
 Sourwood
 Avocado*
 Frasier's Photinia (Redtip)
Spruce species** (Colorado Blue, Norway, etc.)
 Lily-of-the-Valley Shrub
 Calabrian Pine
 Canary Island Pine
 Slash Pine
 Aleppo Pine
 Austrian Black Pine
 Longleaf Pine
 Monterey Pine
 Eastern White Pine
 Scotch Pine
 Loblolly Pine
 Japanese Black Pine
 Virginia Pine

(Back Panel)

<i>Pistacia</i> spp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>Pittosporum tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn Scarlet
<i>Pyracantha fortuneana</i>	Firethorn
<i>Pyracantha koidzumii</i>	Firethorn
<i>Pyrus</i> spp.	Bradford Pear spp.
<i>Quercus rubra</i>	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Rhododendron</i> (including Azalea)	'Coral Bells' 'Formosa' 'Hino-crimson' 'PJM' 'Roseum Elegans'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rumohra adiantiformis</i>	Leatherleaf Fern
<i>Santolina virens</i>	
<i>Sedum album</i>	Stonecrop
<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>Taxus media</i>	Yew
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa</i> spp.	Tulip species
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>Viburnum odoratissimum</i>	Sweet Viburnum
<i>Viburnum plicatum</i>	Japanese Snowball
<i>Viburnum rigidum</i>	Canary Island Viburnum
<i>Viburnum tinus</i>	Laurustinus
<i>Viburnum trilobum</i>	Cranberry Bush
<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Vinca
<i>Vinca minor</i>	Dwarf Periwinkle
<i>Vitis</i> spp.	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet
<i>Yucca filamentosa</i>	Yucca, Adam's Needle

*Do not use on food producing trees, vines, or plants.
**Not for use on container grown plants.

Table 2 - Tolerant Ornamental Species/Varieties - All States Except CA

Scientific name	Common name
<i>Abelia grandiflora</i>	Abelia: Sherwood
<i>Achillea</i> spp.	Yarrow: King Edward

(Back Panel)

Agapanthus orientalis
Akebia quinata
Allium cernuum
Anemone hybrida
Aquilegia spp.
Artemisia spp.
Aster spp.
Aster X frikartii
Athyrium filix-femina
Begonia spp.
Bergenia cordifolia
Boltonia asteroides
Bougainvillea spp.
Buddleia davidii
Callistemon citrinus
Campanula carpatica
Campis X tagliabuana
Ceratostigma plumbaginoides
Chrysanthemum nipponicum
Coreopsis spp.

Crocsmia spp.
Delosperma spp.
Delphinium spp.
Dianthus deltooides
Dianthus gratianopolitanus
Echinacea pupurea
Forsythia suspensa
Gaillardia spp.
Gaura spp.
Gentiana dahurica
Geranium cinereum
Gypsophila repens
Helianthemum spp.
Hemerocallis spp.
Heucherella spp.
Hibiscus spp.
Hosta plantaginea
Hosta sieboldiana
Houttuynia cordata var. *variegata*
Hydrangea macrophylla
Inula ensifolia
Iris ensata
Iris siberica
Juniperus davurica
Lagerstromia indica x fauriei
Lantana montevidensis
Lavender spp.
Leontopodium alpinum
Ligustrum sinense
Lilium spp.
Lirope muscari var. *variegata*

Five-Leaf or Chocolate Vine
 Lady's Leek, Nodding Onion
 Japanese Anemone
 Aquilegia: Red and Gold
 Wormwood; Silver Mound, Castle
 Aster: Bonny Blue, Purple Dome

Lady Fern; Fern Lady
 Fibrous Begonia: Hardy Grandis

Snowbank
 Bougainvillea
 Butterfly-Bush (Dwarf Blue); Royal Red
 Crimson Bottlebrush
 Tussock Bellflower; (White Clips)
 Trumpet Creeper, Trumpet Flower; Madame Galen

Coreopsis (Calliopsis): Early Sunrise,
 Moonbeam
 Lucifer
 Cooperi Pink
 Larkspur; Blue Elf
 Dianthus, Maiden Pinks 'Zing'
 Cheddar Pink
 Coneflower, Purple; Magnus
 Weeping Forsythia
 Gaillardia, Blanket Flower: 'Goblin'

Gentian
 Cranesbill
 Baby's Breath
 Sunrose
 Daylily: Aztec Gold, Stella De Oro, Tender Love
 Coral Bell; Bridget Bloom
 Mallow; Disco Belle White
 Hosta, Plantain Lily (Fragrant)
 Hosta, 'Searsucker'

Bigleaf Hydrangea

Sword-Leaved Iris; Jodlesong
 Siberian Iris; Cabernet
 Parsoni
 Crape Myrtle; Tuscarora
 Weeping Lantana
 Lavender; Munstead
 Edelweiss
 Chinese Privet; Variegata
 Lily: Jazz
 Lirope, Variegated

(Back Panel)

<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia cardinalis</i>	Cardinal Flower, Indian Pink
<i>Loropetalum chinense</i>	Burgundy
<i>Lythrum</i> spp.	Loosestrife; Modern Pink
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Oenothera missouriensis</i>	Evening Primrose
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly): Gulf Tide
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	Dragonhead, False; Vivid
<i>Physostegia virginiana</i>	Oak, Shumard's Red
<i>Quercus shumardii</i>	Yedda Hawthorne
<i>Rhaphiolepis umbellata</i>	'Delaware Valley White'
<i>Rhododendron</i>	'Flame Creeper'
(including azalea)	'Girard Crimson'
	'George L. Tabor'
	Wakeiebisu
	White Gumpo
<i>Rudbeckia</i> spp.	Black-Eyed Susan; Goldstrum
<i>Saxifraga</i> spp.	Saxifrage; Purple Dome
<i>Scabiosa</i> spp.	Pincushion Flower
<i>Sedum caudicola</i>	Stonecrop; Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea bumalda</i>	Spiraea: Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Teucrium</i> spp.	Germander
<i>Thalictrum dipterocarpum</i>	Meadow Rue
<i>Veronica</i> spp.	Veronica, Speedwell; Sunny Border
<i>Viburnum suspensum</i>	Arrowood Viburnum

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

15/14

(Back Panel)

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVnot021505)

Manufactured by:

The Andersons Lawn Fertilizer Division, Inc.
d/b/a Free Flow Fertilizer
PO Box 119
Maumee, OH 43537

[Optional marketing information:]

[Terre]

[To help your lawn feel & look like a plush carpet, here are some steps for you to follow.]

[Step 1 In early Spring Stop crabgrass before it takes over your lawn with Terre Fertilizers & Crabgrass preventers or use straight Crabgrass preventer. If spot seeding is required, seed with Terre Seed mixes & use "Tupersan" crabgrass control. Do not use any other crabgrass preventer when seeding.]

[Step 2 In late Spring Control Dandelions and broadleaf weeds with our Terre Weed & Feed, or straight weed controls.]

[Step 3 Early Summer Control your insects with Terre Fertilizer & Insect controls or straight controls such as Merit.]

[Step 4 Fall Overseed needed areas in early fall and feed with one of our fertilizers. Then, in late fall prepare your lawn for Winter dormancy by feeding with our Fall Fertilizer. Fall fertilizer (Nov -Dec.) is the key to a strong weed-free lawn.]

Date Printed	2/15/05
Date Approved	4/29/04
Supersedes	11/5/97
Label changes	Notification - optional marketing
Filename	9198-124
Label code	RVnot021505



The Andersons, Inc.
P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

16/16
NOTIFICATION

MAR 10 2005

February 15, 2005

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7505C)
US Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, DC 20460-0001

RE: Notification of Minor Label Change

SUBJECT: The Andersons Barricade Herbicide II
EPA Reg. No. 9198-124

Per PR Notice 98-10 enclosed is an EPA Form 8570-1, an application notifying the Agency of a revision to the homeowner version of the labeling for the subject registration. One copy of the revised labeling is also enclosed with the additions underlined and marked in blue.

We have made the following change by notification:

1. Added optional marketing statements on last page

Enclosed is a card indicating the acceptance or rejection of this notification. Please complete the card and return via mail. If there are any questions or comments regarding this notification, please feel free to contact me.

Sincerely,
The Andersons Lawn Fertilizer Division, Inc.

Dawn L. Walters

Dawn L. Walters
Regulatory Administrator
Phone 334-262-2766
Fax 334-265-5703
Email dawn_walters@andersonsinc.com

enclosures