



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Debbie Ziehr
The Andersons
P.O. Box 119
Maumee, Ohio 43537

DEC - 9 2010

Subject: Application for Pesticide Notificaion (PRN 98-10)
Request Additional Brand Name: The Andersons Turf Products Fertilizer with
0.28% Prodiamine Herbicide 17-0-0
EPA Reg. No. 9198-124
Application Dated: September 22, 2010

Dear Ms. Ziehr,

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for The Andersons Fertilizer with 0.28% Barricade Herbicide (EPA Reg. No. 9198-124) dated August 18, 2010. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at (703) 305-1243.

Sincerely,

A handwritten signature in cursive script that reads "Kathryn V. Montague".

Kathryn V. Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 9198-124	2. EPA Product Manager Montague	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Fertilizer with 0.28% Barricade Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow Fert. P.O. Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Alternate Brand Name per PR Notice 98-10: The Andersons Turf Products Fertilizer with 0.28% Prodiamine Herbicide 17-0-0

This notification is consistent with the guidance of PR Notice 98-10 and the requirements of EPA's regulations at 40 CFR 156.10 and 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 98-10 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted				<input type="checkbox"/> Glass	<input checked="" type="checkbox"/> Paper
	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container	Other (Specify) _____
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container	4. Size(s) Retail Container 50 lbs. - 2,000 lbs.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label		
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Debbie Ziehr		Title Regulatory Administrator		Telephone No. (Include Area Code) 419-891-6671	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature 			3. Title Regulatory Administrator		
4. Typed Name Debbie Ziehr			5. Date 9/22/10		



The Andersons, Inc.

P.O. Box 119 • Maumee, Ohio 43537 • 419/893/5050

3/6

September 22, 2010

Document Processing Desk (NOTIF)
Office of Pesticide Programs - (7504P)
U. S. Environment Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Subject: NOTIFICATION - Additional Brand Name

RE: **The Andersons Fertilizer with 0.28% Barricade Herbicide
EPA Reg. No. 9198-124**

Per PR Notice 98-10, enclosed is an EPA Form 8570-1, an application notifying the Agency that we wish to market additional products under the subject basic registration. The additional brand name is:

The Andersons Turf Products Fertilizer with 0.28% Prodiamine Herbicide 17-0-0

One copy of the draft label is enclosed.

If there are any questions regarding this notification, please contact me.

Sincerely,

The Andersons Lawn Fertilizer Division, Inc.

A handwritten signature in cursive script that reads "Debbie Ziehr".

Debbie Ziehr
Regulatory Administrator
Phone #419-891-6671
Fax #419-891-2745
Email: Debbie_ziehr@andersonsinc.com


Enclosure



Turf Products

Fertilizer with 0.28% Prodiamine Herbicide 17-0-0

NOTIFICATION
DEC 06 2010



For preemergence control of grass and broadleaf weeds in established lawns and turfgrasses (excluding golf course putting greens).

GUARANTEED ANALYSIS		
Total Nitrogen (N)	17.00% Urea Nitrogen	17.00%
Plant nutrients derived from urea.		
ACTIVE INGREDIENT:		
Prodiamine†		0.28%
OTHER INGREDIENTS:		
Total		99.72%
†(CAS No. 29091-21-2)		100.00%
EPA Reg. No. 9198-124		
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A		
Underlined letter is first letter used in run code on bag.		

Rate
This product may be applied as a single application or in sequential applications must be made prior to germination of the target weeds. This

Maximum use rate selection should be based on turf species. The rate applied.

Maximum
This product is recommended for use on the turf grass species list the following table:

Maximum
1These are the maximum rates per calendar year by species limits
2May be used on newly sprung or plugged Bermudagrass at rate Bermudagrass stolon rooting may be temporarily retarded. Suppr rates used in sprigging situations.

Do not apply more than 536 lbs. per acre (12.3 lbs./1,000 sq. ft.) or Use higher rates of this product to achieve higher levels of fertility application rates specified in the Maximum Annual Rates table.

Weeds Controlled
When used at the recommended rates this product will control: B; Common; Chickweed, Mouseear (from seed); Crabgrass (Large, S Henbit; Itchgrass; Johnsongrass (from seed); Jungletice; Knotweed Browntop); Pigweed; Purslane, Common; Pursley, Florida; Rescue; Sprangletop; Spurge, Prostrate; Witchgrass; Woodsorrel, Yellow (f

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

- FIRST AID**
- If in eyes** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
 - If swallowed** Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
 - If on skin or clothing** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
 - If inhaled** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS
This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

GENERAL INFORMATION
The Andersons Turf Products Fertilizer with 0.28% Prodiamine Herbicide 17-0-0 is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in turfgrasses. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

APPLICATION DIRECTIONS
Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF
This product is a preemergence herbicide that, when properly applied as directed, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

- Use Restrictions and Precautions**
The following precautions apply to the use of this product in turf grasses and lawns:
- Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
 - Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer.
 - Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
 - Do not apply to newly set sod until the following year.
 - Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
 - Disturbing the herbicide barrier with cultural practices such as diking may result in reduced weed control.
 - Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
 - Do not graze or feed livestock forage cut from areas treated with this product.
 - Do not apply this product through any type of irrigation system unless otherwise instructed by this label.
 - Do not apply aerially.
 - Do not apply to golf course putting greens.

Maximum Application Rate of this p	
Turf Species	Rate
Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	5.4 to 12.3
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	5.4 to 12.3
Creeping Red Fescue	5.4 to 12.3
Creeping Bentgrass	5.4 to 12.3

¹In many areas a single application of 5.4 to 12.3 lbs./1,000 sq. ft. heavy goosegrass pressure and/or an extended growing season, 1 5.4 to 8.2 lbs./1,000 sq. ft. (236 to 357 lbs./A) followed after 60 to the Maximum Annual Rates Table. Do not exceed the maximum

²Applications for this weed should be made in late summer, fall, or specified in the Maximum Annual Rates table.

³Suppression only.

Sequential applications may be made so long as the total amount the Maximum Annual Rates table. All applications must be made

When to Apply After Overseeding Turf
Do not apply to overseeded turf within 60 days after seeding or ur is likely if this product is applied before seedling secondary roots :


When to Overseed After Application - All States*
This product will inhibit the germination of turf species if overseed best overseeding/reseeding results.

The Andersons Turf Products Fertilizer with Prodiamine Herbicide 17-0-0 Rate		
lbs./1,000 sq. ft.	lbs./A	ai
4.1	179	0.1
5.4	236	0.1
6.1	264	0.1
6.6	289	0.1
8.2	357	1.1
9.4	411	1.1
10.6	464	1.1
12.3	536	1.1

* Note: In AZ, CA, NV and TX, the overseeding interval can be st ryegrass. See the next section "Poa annua Control in Establisher only".

Poa annua Control in Established Bermudagrass Overseeder
Use on golf courses (excluding golf course putting greens) and la lbs./A).

CRABGRASS GERMINATION DATES



Approx. Date:
 1 - After May 30
 2 - After May 10
 3 - After April 20
 4 - After March 20
 5 - Jan. 1 to March 20

Amount to Apply	
Lbs./Acre	Lbs./1,000
135 - 236	3.1 - 5.4

Products

Prodiamine

NOTIFICATION
 DEC 06 2010

of weeds in established lawns (greens).

S	17.00%
F292	0.28%
	99.72%
	100.00%

Rates of Application
 This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Annual Rates Table

¹These are the maximum rates per calendar year by species limitations.
²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 186 lb./acre or 4.2 lb./1,000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Do not apply more than 536 lbs. per acre (12.3 lbs./1,000 sq. ft.) of this product per calendar year.
 Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates table.

Weeds Controlled

When used at the recommended rates this product will control: Barnyardgrass; Bluegrass, Annual (*Poa annua*); Carpetweed; Chickweed, Common; Chickweed, Mouseear (from seed); Crabgrass (Large, Smooth); Crowfootgrass; Cupgrass, Woolly; Foxtails, Annual; Goosegrass¹; Henbit; Itchgrass; Johsongrass (from seed); Junglerice; Knotweed; Kochia; Lambsquarter, Common; Lovegrass; Oxalis; Panicum (Texas, Fall, Browntop); Pigweed; Purslane, Common; Pusley, Florida; Rescuegrass³; Shepherd's Purse²; Signalgrass, Broadleaf; Speedwell, Persian; Sprangletop; Spurge, Prostrate; Witchgrass; Woodsorrel, Yellow (from seed).

Maximum Application Rate of this product per Calendar Year by Turf Grass Species ¹			
Turf Species	lbs. ai/A	lbs. product /A	lbs. product/ 1,000 sq. ft.
Bermudagrass ² ; Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	1.5	536	12.3
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	1.0	357	8.2
Creeping Red Fescue	0.75	264	6.1
Creeping Bentgrass	0.65	236	5.4

¹In many areas a single application of 5.4 to 12.3 lbs./1,000 sq. ft. (236 to 536 lbs./A) of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 5.4 to 8.2 lbs./1,000 sq. ft. (236 to 357 lbs./A) followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rates Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rates Table above.

²Applications for this weed should be made in late summer, fall, or winter prior to germination. Do not exceed the maximum application rates specified in the Maximum Annual Rates table.

³Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum application rates specified in the Maximum Annual Rates table. All applications must be made prior to germination of the weed seeds.

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When to Overseed After Application - All States*

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

The Andersons Turf Products Fertilizer with Prodiamine Herbicide 17-0-0 Rate			Interval Before Overseeding		
lbs./1,000 sq. ft.	lbs./A	ai/A	North	Transition	South
4.1	179	0.50	4 mo.	4 mo.	4 mo.
5.4	236	0.65	5 mo.	4 mo.	4 mo.
6.1	264	0.75	6 mo.	5 mo.	5 mo.
6.6	289	0.81	--	6 mo.	6 mo.
8.2	357	1.00	--	7 mo.	7 mo.
9.4	411	1.15	--	--	9 mo.
10.6	464	1.30	--	--	10 mo.
12.3	536	1.50	--	--	12 mo.

* Note: In AZ, CA, NV and TX, the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section: *Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV and TX only).

***Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV and TX only)**

Use on golf courses (excluding golf course putting greens) and lawns when overseeding with perennial ryegrass (minimum seed rate of 350 lbs./A).

CHILDREN

PESTICIDES AND DOMESTIC ANIMALS

n. Avoid contact with skin, eyes, or clothing. Avoid frequently repeated skin contact may cause allergic

minutes. Remove contacts, if present, after the first 5 doctor for treatment advice.

person sip a glass of water if able to swallow. Do not or. Do not give anything by mouth to an unconscious

ity of water for 15-20 minutes. Call a poison control

in ambulance, then give artificial respiration, preferably or for treatment advice.

doctor or going for treatment. You may also contact

c to fish. However, at concentrations substantially above areas where surface water is present or to intertidal areas us to aquatic organisms in adjacent sites. Do not contami-

eling.

ercial seed production, or for research purposes.

reemergence herbicide that provides residual control of eds by inhibiting weed seeds germination and root ted by at least 0.5 inch of rainfall or irrigation, or with ys following application.

control certain grass and broadleaf weeds in established year is given for each turf grass species in the Maximum obtained when this product is activated by at least 0.5 inch lication. See the map below for approximate crabgrass

ded grasses. and mowing, whichever is longer. ary roots are in the second inch of soil, not thatch plus

may result in turf injury. i reduced weed control. tgrass, velvet bentgrass, or annual bluegrass (*Poa annua*)

structed by this label.

STORAGE AND DISPOSAL
 Do not contaminate water, food, or and separated from other pesticide approved waste disposal facility. not reuse or refill this container. incineration facility; or, if allowed l

IMPORTANT: READ BEFORE USE
 Read the entire Directions for Use return the unopened product conta Liability:


WARRANTY DISCLAIMER and
 Manufacturer warrants that this p the label when used in strict acco MERCHANTABILITY OR FITNESS LIMIT ANY LIABILITY FOR SPEC of product, at Buyer's choice. TC OR OTHER DAMAGES ARISING

LEGAL RIGHTS:
 THIS LIMITED WARRANTY GIVE JURISDICTION TO STATE/JURIS

50 lbs. treats 50 lbs. treats 50 lbs. treats 50 lbs. treats
SPREADER
Andersons AccuPro 2000, SR-2000 (cone 4)
Scotts R-8A (cone 4)
Lesco 020093 (pattern 0.25)
Lesco 705698 (pattern 0.25)
Lely Models WTR, WFR, HR, 1 (PTO at 450 rpm) Pattern III-B
Vicon (03 Series)

The Andersons logo is a registered

APPLICATION DATES



Approx. Date:

- 1 - After May 30
- 2 - After May 10
- 3 - After April 20
- 4 - After March 20
- 5 - Jan. 1 to March 20

tion
to control weeds germinating throughout the year. All applica-
not control established weeds.

ie of residual weed control provided by this product is related to

Rates
wing table. Do not exceed the maximum yearly rate as given in

Tables Table

eed 186 lb./acre or 4.2 lb./1,000 sq. ft. Newly sprigged or plugged
f Foxtail, Goosegrass and Rescuegrass due to reduced product

t per calendar year.
ontrol for each turf type, but do not exceed the maximum

;; Bluegrass, Annual (*Poa annua*); Carpetweed; Chickweed,
wfootgrass; Cupgrass, Woolly; Foxtails, Annual; Goosegrass¹;
nbsquarter, Common; Lovegrass; Oxalis; Panicum (Texas, Fall,
herd's Purse²; Signalgrass, Broadleaf; Speedwell, Persian;

Calendar Year by Turf Grass Species¹

lbs. ai/A	lbs. product /A	lbs. product/ 1,000 sq. ft.
1.5	536	12.3
1.0	357	8.2
0.75	264	6.1
0.65	236	5.4

3 lbs./A) of this product will control goosegrass. However, under
ve weed control may be obtained by making an initial application of
a second application at doses that would not exceed those given in
f grass species listed in the **Maximum Annual Rates Table** above.

x to germination. Do not exceed the maximum application rates

plied does not exceed the maximum application rates specified in
mination of the weed seeds.

second mowing, whichever is longer. Injury to desirable seedlings
econd inch of soil, not thatch plus soil.

n after application. Follow rates and intervals in the table below for

Interval Before Overseeding		
North	Transition	South
4 mo.	4 mo.	4 mo.
5 mo.	4 mo.	4 mo.
6 mo.	5 mo.	5 mo.
--	6 mo.	6 mo.
--	7 mo.	7 mo.
--	--	9 mo.
--	--	10 mo.
--	--	12 mo.

established bermudagrass that has been overseeded with perennial
grass Overseeded with Perennial Ryegrass (AZ, CA, NV and TX)

ennial Ryegrass (AZ, CA, NV and TX only)
overseeding with perennial ryegrass (minimum seed rate of 350

How Much and When to Apply

Amount to Apply		When to Apply	Expected Control	Use Precautions
Lbs./Acre	Lbs./1,000			
135 - 236	3.1 - 5.4	6 to 8 weeks before ryegrass overseeding Second application 4-8 weeks after over-seeding date or when perennial ryegrass roots are in the second inch of soil	One application for 70% or greater control of <i>Poa annua</i> Second application should enhance the residual under normal weather conditions	1. Some seed mortality and temporary reduction in root growth of new seedlings may occur. 2. To reduce the potential for seeding mortality, maintain a moist seedbed with light, frequent irrigation. 3. Make no more than 2 applications per year for this use, and do not exceed 302 lbs/A (6.9 lbs/1,000) for these two applications for <i>Poa annua</i> control in ryegrass overseeded turf. 4. Do not make a second application if any injury to the ryegrass is observed after the first application. 5. Do not make a second application unless the product was first applied before overseeding.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Storage:** Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Completely empty container into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/ JURISDICTION TO STATE/JURISDICTION.

Recommended Spreader Settings

These suggested spreader settings are not intended to replace calibration.

Please calibrate your spreader before applying product.

- 50 lbs. treats 9,200 sq. ft. (0.21 acre) at the 236 lbs. product/acre - LOW RATE (5.4 lbs/1,000 sq. ft.)
- 50 lbs. treats 8,100 sq. ft. (0.19 acre) at the 264 lbs. product/acre - MEDIUM RATE (6.1 lbs/1,000 sq. ft.)
- 50 lbs. treats 6,000 sq. ft. (0.14 acre) at the 357 lbs. product/acre - HIGH RATE (8.2 lbs/1,000 sq. ft.)
- 50 lbs. treats 4,000 sq. ft. (0.09 acre) at the 536 lbs. product/acre - HEAVY RATE (12.3 lbs/1,000 sq. ft.)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS			
			LOW RATE 0.65 lbs. ai/A	MEDIUM RATE 0.75 lbs. ai/A	HIGHRATE 1.00 lbs. ai/A	HEAVY RATE 1.50 lbs. ai/A
Andersons AccuPro 2000, SR-2000 (cone 4)	3 mph	11 feet	N 1/2	O 1/2	Q	S
Scotts R-8A (cone 4)	3 mph	11 feet	M	M 1/2	O 1/2	Q 1/2
Lesco 020093 (pattern 0.25)	3 mph	10 feet	I	I 1/2	K 1/2	M 1/2
Lesco 705698 (pattern 0.25)	3 mph	10 feet	20	21	25	29
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern III-B	4.5 mph	32 feet	7	7 1/2	8 3/4	10
Vicon (03 Series)	4.5 mph	36 feet	35	38	46	62

Manufactured by:
The Andersons Lawn Fertilizer Division, Inc.
P. O. Box 119, Maumee, OH 43537
www.andersonsturf.com

The Andersons logo is a registered trademark of The Andersons, Inc.

SGN 215
Net Wt. 50 Lbs. (22.68 kg)



0 31865 31588 6
APT17PR50
C10 (APTBLKPMBRNB)