

9198-124

4/26/2012

1/6



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

APR 26 2012

Debbie Ziehr
The Andersons Inc
P O Box 119
Maumee OH 43537

Subject Notifications per PR Notice 98 10 – Add Alternate Brand Names
THE ANDERSONS BARRICADE HERBICIDE II
EPA Reg No 9198 124
Application Dated – April 3 2012

Proposed Alternate Brand Names 17 2 5 Fertilizer with 0 28% Barricade Herbicide
28 0 5 Fertilizer with 0 28% Prodiamine Herbicide


Dear Ms Ziehr


The Agency is in receipt of your Applications for Pesticide Notification under Pesticide Registration Notice (PRN) 98 10 for the subject product

The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98 10 and finds that the actions requested falls within the scope of PRN 98 10 The label submitted with the application has been date stamped Notification and will be placed in our records

If you have any questions regarding this letter please contact Maggie Rudick at (703) 347 0257 or rudick_maggie@epa.gov

Sincerely


Kathryn Montague Product Manager 23
Herbicide Branch
Registration Division (7505P)





United States Environmental Protection Agency Washington DC 20460

Registration Amendment Other (checked)

OPP Identifier Number

Application for Pesticide - Section I

1 Company/Product Number 9198 124
2 EPA Product Manager Montague
3 Proposed Classification None Restricted
4 Company/Product (Name) The Andersons Fertilizer with 0 28 / Barricade Herbicide
5 Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert Div Inc
6 Expedited Review In accordance with FIFRA Section 3(c)(3) (b)(i) my product is similar or identical in composition and labeling to EPA Reg No Product Name

Section II

Amendment Explain below
Resubmission in response to Agency letter dated
Notification Explain below (checked)
Final printed labels in response to Agency letter dated
Me Too Application
Other Explain below

NOTIFICATION APR 26 2012

Explanation Use additional page(s) if necessary (For section I and Section II)
Notification of Alternate Brand Name per PR Notice 98 10 28 0 5 Fertilizer with 0 28 / Prodiamine Herbicide
This notification is consistent with the guidance of PR Notice 98 10 and the requirements of EPA's regulations at 40 CFR 156 10 and 40 CFR 152 46 and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA I further understand that if this notification is not consistent with the guidance of PR Notice 98 10 and the requirements of 40 CFR 156 10 and 40 CFR 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA

Section III

1 Material This Product Will Be Packaged In
Child Resistant Packaging Yes No (checked)
Unit Packaging Yes No (checked)
Water Soluble Packaging Yes No (checked)
2 Type of Container Metal Plastic (checked) Glass Paper Other (Specify)
3 Location of Net Contents Information Label Container (checked)
4 Size(s) Retail Container 50 2 000 lbs
5 Location of Label Directions On Label (checked)
6 Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled (checked) Other

Section IV

1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)
Name Debbie Ziehr Title Regulatory Administrator Telephone No (Include Area Code) 419 891 6671

Certification I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law
2 Signature Debbie Ziehr 3 Title Regulatory Administrator
4 Typed Name Debbie Ziehr 5 Date 4/3/2012
6 Date Application Received (Stamped)



The Andersons Inc
PO Box 119 Maumee Ohio 43537 419/893/5050

April 3 2012

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U S Environment Protection Agency
Room S 4900 One Potomac Yard
2777 South Crystal Drive
Arlington VA 22202 4501

Subject NOTIFICATION Additional Brand Name
RE The Andersons Fertilizer with 0 28% Barricade Herbicide
EPA Reg No 9198 124

Per PR Notice 98 10 enclosed is an EPA Form 8570 1 an application notifying the Agency that we wish to market additional products under the subject basic registration The additional brand names are as follows

- 1) **28 0 5 Fertilizer with 0 28% ~~Barricade~~ Herbicide** *Prodicamine (change per 8/15/12 phone call)*

One copy of each of the draft labels are enclosed

If there are any questions regarding this notification please contact me

Sincerely

The Andersons Lawn Fertilizer Division Inc

Debbie Ziehr
Regulatory Administrator
Phone #419 891 6671
Fax #419 891 2745
Email Debbie_ziehr@andersonsinc.com

Enclosure

28-0-5 Fertilizer with 0.28% Proflaminate Herbicide

NOTIFICATION

APR 26 2012

CRABGRASS



• Contains NutriSphere-N®

For preemergence control of grass and broadleaf weeds in established lawns and turfgrasses (excluding golf course putting greens)

GUARANTEED ANALYSIS

Total Nitrogen (N) 28 00 / Urea N trogen	28 00 /
Soluble Potash (K ₂ O)	5 00 /
Plant nutrients derived from urea and potassium chloride	
5 60 / nitrogen stabilized with maleic itaconic co polymer Chlorine (Cl) Max	4 24 /
	F292
ACTIVE INGREDIENT Proflaminate	0 28 /
OTHER INGREDIENTS Total	99 72 /
†(CAS No 29091 21 2)	100 00 /

EPA Reg No 9198 124
EPA Est No 9198 OH 1M 9198 OH 2B 9198 AL 001A
Underlined letter is first letter used in run code on bag

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye injury (irritation) Harmful if inhaled or absorbed through the skin Avoid contact with skin eyes or clothing Avoid breathing dust Wash thoroughly with soap and water after handling Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals

FIRST AID

- If in eyes** Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contacts if present after the first 5 minutes then continue rinsing Call a poison control center or doctor for treatment advice
- If swallowed** Call a poison control center or doctor for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to by a poison control center or doctor Do not give anything by mouth to an unconscious person
- If on skin or clothing** Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
- If inhaled** Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 800 757 8951 for emergency medical treatment information

ENVIRONMENTAL HAZARDS

This product has low solubility in water At the limit of solubility this product is not toxic to fish However at concentrations substantially above the level of water solubility it may be toxic to fish Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites Do not contaminate water when disposing of equipment washwaters

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

Not for use on turf being grown for sale or other commercial use as sod or for commercial seed production or for research purposes

GENERAL INFORMATION

28 0 5 Fertilizer with 0 28 / Proflaminate Herbicide is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in turfgrasses This product controls susceptible weeds by inhibiting weed seeds germination and root development Most effective weed control will be obtained when this product is activated by at least 0 5 inch of rainfall or irrigation or with shallow (1 to 2 inches) incorporation prior to weed seed germination and within 14 days following application

APPLICATION DIRECTIONS

Apply uniformly through suitable calibrated application equipment

ESTABLISHED TURF

This product is a preemergence herbicide that when properly applied as directed will control certain grass and broadleaf weeds in established turf grasses and lawns The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0 5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application See the map below for approximate crabgrass seed germination dates

Use Restrictions and Precautions

The following precautions apply to the use of this product in turf grass Application of this product may thin emerged annual bluegrass an Do not apply to over seeded turf within 60 days after seeding or u Injury to desirable seedlings is likely if this product is applied befo Do not apply to newly set sod until the following yea Application of this product to turf stressed by drought low fertily Disturbing the herbicide barrier with cultural practices such as disl Do not apply this product to putting greens or areas where dorch are desirable species Do not graze or feed livestock forage cut from areas treated with t Do not apply this product through any type of irrigation system un Do not apply aurally Do not apply to golf course putting greens

This product may be applied as a single application or in sequential must be made prior to germination of the target weeds This product

Maximum use rate selection should be based on turf species The length c

This product is recommended for use on the turf grass species listed following table

Maximum Application Rate of this p	
Turf Species	Maximum Annual Rate
Bermudagrass ² Bahiagrass Centipedegrass Seashore Paspalum St. Augustnegrass Tall Fescue (including turf type) Zoysia	
Buffalograss Kentucky Bluegrass Perennial Ryegrass	
Creeping Red Fescue	
Creeping Bentgrass	

¹These are the maximum rates per calendar year by species limitatio
²May be used on newly sprgged or plugged Bermudagrass at rates Bermudagrass stolon rooting may be temporarily retarded³ Suppress rates used in sprgging situations

Do not apply more than 536 lbs per acre (12 3 lbs/1 000 sq ft) of thi Use higher rates of this product to achieve higher levels of fertility and rates specified in the Maximum Annual Rates table

Weeds Controlled

When used at the recommended rates this product will control Barn Chickweed Mouseear (from seed) Crabgrass (Large Smooth) Crow Johnsongrass (from seed) Junglerice Knotweed Kochia Lambsquart Purslane Common Pusley Florida Rescuegrass³ Shepherd's Purse Witchgrass Woodsorrel Yellow (from seed)

¹In many areas a single application of 5 4 to 12 3 lbs/1 000 sq ft (23 goosegrass pressure and/or an extended growing season most effec lbs/1 000 sq ft (236 to 357 lbs/A) followed after 60 to 90 days by a s Annual Rates Table Do not exceed the maximum rate for turf grass

²Applications for this weed should be made in late summer fall or w specified in the Maximum Annual Rates table

³Suppression only

Sequential applications may be made so long as the total amount of Maximum Annual Rates table All applications must be made prior to

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until likely if this product is applied before seedling secondary roots are in

When to Overseed After Applic tior All States


This product will inhibit the germinaj on of turf species if overseeded t overseeding/reseeding results

8 C 5 Fertilizer with 0 28 / Proflaminate Herbicide Rate		
lbs/1 000 sq ft	lbs/A	at
4 1	179	0
5 4	236	0
6 1	264	0
6	289	0
8 2	357	1
9 4	411	1
10 6	464	1
12 3	536	1

576

NOTIFICATION
APR 26 2012

CRABGRASS GERMINATION DATES



- Approx Date**
 1 After May 30
 2 After May 10
 3 After April 20
 4 After March 20
 5 January 1 to March 20

In AZ CA N
 regrass See the ne
Poa annua Control
 Use on golf courses

grasses

28 00 /
 5 00 /
 4 24 /
 F292

0 28 /
 99 72 /
 100 00 /

Avoid breathing
 in some

first 5

Do not
 conscious

control center or

preferably

tact

ally above the
 il areas below
 taminant water

established turf
 maximum Annual
 nor to weed

Use Restrictions and Precautions

The following precautions apply to the use of this product in turf grasses and lawns
 Application of this product may thin emerged annual bluegrass and newly overseeded grasses
 Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer
 Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil not thatch plus soil
 Do not apply to newly set sod until the following year
 Application of this product to turf stressed by drought low fertility or pest damage may result in turf injury
 Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control
 Do not apply this product to putting greens or areas where dichondra colonial bentgrass velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species
 Do not graze or feed livestock forage cut from areas treated with this product
 Do not apply this product through any type of irrigation system unless otherwise instructed by this label
 Do not apply aerially
 Do not apply to golf course putting greens

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year All applications must be made prior to germination of the target weeds This product will not control established weeds

Maximum use rate selection should be based on turf species The length of time of residual weed control provided by this product is related to the rate applied

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table Do not exceed the maximum yearly rate as given in the following table

Maximum Annual Rates Table

Maximum Application Rate of this product per Calendar Year by Turf Grass Species ¹			
Turf Species	lbs at/A	lbs product /A	lbs. product/ 1 000 sq ft
Bermudagrass ² Bahiagrass Centipedegrass Seashore Paspalum St Augustinegrass Tall Fescue (including turf type) Zoysia	15	536	12.3
Buffalograss Kentucky Bluegrass, Perennial Ryegrass	10	357	8.2
Creeping Red Fescue	0.75	264	6.1
Creeping Bentgrass	0.65	236	5.4

¹These are the maximum rates per calendar year by species limitations

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 186 lbs/acre or 4.2 lbs/1 000 sq ft Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded Suppression or yield of Foxtail Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations

Do not apply more than 536 lbs per acre (12.3 lbs/1 000 sq ft) of this product per calendar year Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type but do not exceed the maximum application rates specified in the Maximum Annual Rates table

Weeds Controlled

When used at the recommended rates this product will control Barnyardgrass Bluegrass Annual (*Poa annua*) Carpetweed Chickweed Common Chickweed Mouseear (from seed) Crabgrass (Large Smooth) Crowfootgrass Cupgrass Woolly Foxtails Annual Goosegrass¹ Henbit Itchgrass Johnsongrass (from seed) Jungle Knotweed Kochia Lambsquarter Common Lovegrass Oxalis Panicum (Texas Fall Browntop) Pigweed Purslane Common Pursley Florida Rescuegrass³ Shepherd's Purse² Signalgrass Broadleaf Speedwell Persian Sprangletop Spurge Prostrate Witchgrass Woodsorrel Yellow (from seed)

¹In many areas a single application of 5.4 to 12.3 lbs/1 000 sq ft (236 to 536 lbs/A) of this product will control goosegrass However under heavy goosegrass pressure and/or an extended growing season most effective weed control may be obtained by making an initial application of 5.4 to 8.2 lbs/1 000 sq ft (236 to 357 lbs/A) followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rates Table** Do not exceed the maximum rate for turf grass species listed in the **Maximum Annual Rates Table** above

²Applications for this weed should be made in late summer fall or winter prior to germination Do not exceed the maximum application rates specified in the Maximum Annual Rates table

³Suppression only

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum application rates specified in the Maximum Annual Rates table All applications must be made prior to germination of the weed seeds

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil not thatch plus soil

When to Overseed After Application All States

This product will inhibit the germination of turf species if overseeded too soon after application Follow rates and intervals in the table below for best overseeding/reseeding results

8 C 5 Fertilizer with 0.28 / Prodiame Herbicide Rate			Interval Before Overseeding		
lbs/1 000 sq ft	lbs/A	at/A	North	Transition	South
4.1	179	0.50	4 mo	4 mo	4 mo
5.4	236	0.65	5 mo	4 mo	4 mo
6.1	264	0.75	6 mo	5 mo	5 mo
6	289	0.81		6 mo	6 mo
8.2	357	1.00		7 mo	7 mo
9.4	411	1.15			9 mo
10.6	464	1.30			10 mo
12.3	536	1.50			12 mo

Amount to Apply		
Lbs/Acre	Lbs/	
135	236	31

- Andersons Model
- Andersons 2000 S
- Scotts R 8A (cone)
- Lesco 020093 (pa)
- Lesco 705698 (pa)
- Lely Models WTR V (PTO at 450 rpm)
- Vicon (03 Series)

STORAGE AND DIS

Do not contaminate separated from other waste disposal facilities this container Offer allowed by state and

IMPORTANT READ

Read the entire Directions return the unopened Liability

WARRANTY DISCL

Manufacturer warrant when used in strict accordance with DIRECTIONS OR FITNESS FOR A SPECIAL CONSEQUENCE TO THE EXTENT CONSISTENT WITH ANY USE


LEGAL RIGHTS

THIS LIMITED WARRANTY IS SUBJECT TO STATE/JURISDICTION

NutriSphere is a registered trademark

6/6

APPLICATION DATES



App	Dat
1	After May 30
2	After May 10
3	After April 20
4	After March 20
5	January 1 to March 20

In AZ CA NV and TX the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section *Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ CA NV and TX only)

Poa annua Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ CA, NV and TX only)
Use on golf courses (excluding golf course putting greens) and lawns when overseeding with perennial ryegrass (minimum seed rate of 350 lbs/A)

How Much and When to Apply

Amount to Apply		When to Apply	Expected Control	Use Precautions
Lbs/Acre	Lbs/1 000			
135 236	31 54	6 to 8 weeks before ryegrass overseeding Second application 4 to 8 weeks after overseeding date or when perennial ryegrass roots are in the second inch of soil	One application for 70% or greater control of <i>Poa annua</i> Second application should enhance the residual under normal weather conditions	1 Some seed mortality and temporary reduction in root growth of new seedlings may occur 2 To reduce the potential for seeding mortality maintain a moist seedbed with light frequent irrigation 3 Make no more than 2 applications per year for this use and do not exceed 302 lbs/A (6.9 lbs/1 000) for these two applications for <i>Poa annua</i> control in ryegrass overseeded turf 4 Do not make a second application if any injury to the ryegrass is observed after the first application 5 Do not make a second application unless the product was first applied before overseeding

ns
reseeded grasses
second mowing whichever is longer
secondary roots are in the second inch of soil not thatch plus soil

age may result in turf injury
ult in reduced weed control
bentgrass velvet bentgrass or annual bluegrass (*Poa annua*)

se instructed by this label

ation
to control weeds germinating throughout the year All applications
rol established weeds

ual weed control provided by this product is related to the rate applied

Rates
ing table Do not exceed the maximum yearly rate as given in the

Tables

Annual Rate by Turf Grass Species ¹		
lbs a/A	lbs product /A	lbs product/ 1 000 sq ft.
15	536	12.3
10	357	8.2
0.75	264	6.1
0.65	236	5.4

186 lbs/acre or 4.2 lbs/1 000 sq ft. Newly sprigged or plugged
tial. Goosegrass and Rescuegrass due to reduced product

r calendar year
of for each turf type but do not exceed the maximum application

uegrass Annual (*Poa annua*) Carpetweed Chickweed Common
upgrass Woolly Foxtails Annual Goosegrass¹ Henbit Itchgrass
Lovegrass Oxalis Panicum (Texas Fall Browntop) Pigweed
ss Broadleaf Speedwell Persian Sprangletop Spurge Prostrate

A) of this product will control goosegrass. However under heavy
ontrol may be obtained by making an initial application of 5.4 to 8.2
ation at doses that would not exceed those given in the **Maximum**
nd in the **Maximum Annual Rates Table** above

germination Do not exceed the maximum application rates

ed does not exceed the maximum application rates specified in the
of the weed seeds

nd mowing whichever is longer Injury to desirable seedlings is
nch of soil not thatch plus soil

r application Follow rates and intervals in the table below for best

Interval Before Overseeding		
North	Inter. Int.	South
4 mo	4 mo	4 mo
5 mo	4 mo	4 mo
6 mo	5 mo	5 mo
	6 mo	6 mo
	7 mo	7 mo
		9 mo
		10 mo
		12 mo

Recommended Spreader Settings
These suggested spreader settings are not intended to replace calibration
Please calibrate your spreader before applying product
50 lbs treats 9 200 sq ft (0.21 acre) at the 236 lbs product/acre **LOW RATE** (5.4 lbs/1 000 sq ft)
50 lbs treats 8 100 sq ft (0.19 acre) at the 264 lbs product/acre **MEDIUM RATE** (6.1 lbs/1 000 sq ft)
50 lbs treats 6 000 sq ft (0.14 acre) at the 357 lbs product/acre **HIGH RATE** (8.2 lbs/1 000 sq ft)
50 lbs treats 4 000 sq ft (0.09 acre) at the 536 lbs product/acre **HEAVY RATE** (12.3 lbs/1 000 sq ft)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS			
			LOW RATE 0.65 lbs a/A	MEDIUM RATE 0.75 lbs a/A	HIGH RATE 1.00 lbs a/A	HEAVY RATE 1.50 lbs a/A
Andersons Model 2000 / AccuPro 2000 (cone 4) Andersons 2000 SR / SR 2000 (cone 4)	3 mph	11 feet	N 1/2	O	Q	S
Scotts R 8A (cone 4)	3 mph	11 feet	M 3/4	N 1/4	P 1/4	Q 3/4
Lesco 020093 (pattern 0.25)	3 mph	10 feet	G	G 1/2	H 1/2	L 1/2
Lesco 705698 (pattern 0.25)	3 mph	10 feet	16	17	19	27
Lely Models WTR WFR HR 1250 (PTO at 450 rpm) Pattern II A	4.5 mph	40 feet	7	7 1/4	8 1/2	10 3/4
Vicon (03 Senes)	4.5 mph	32 feet	32	34	40	48

STORAGE AND DISPOSAL
Do not contaminate water food or feed by storage or disposal. **Storage** Store in original container away from fertilizer feed or food stuffs and separated from other pesticides. **Pesticide Disposal** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal** Completely empty container into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If not available then dispose of empty bag or box in a sanitary landfill or at an incineration facility or if allowed by state and local authorities by burning locally. If burned stay out of smoke.

IMPORTANT READ BEFORE USE
Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable return the unopened product container at once. By using this product user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability.


WARRANTY DISCLAIMER and LIMITATION of LIABILITY
Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product at Buyer's choice. To the extent consistent with applicable law Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS
THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION

Manufactured by
The Andersons Lawn Fertilizer Division Inc
P O Box 119 Maumee OH 43537
www.andersonsturf.com

NutriSphere is a registered trademark of Specialty Fertilizer Products LLC

SGN 215
Net Wt 50 Lbs (22.68 kg)



ASP285NPR5
B12 (APTBLKPBRNB)