

9198-125

09/22/2003

9A 9-3 1/3

Read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 9198-125	2. EPA Product Manager Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) The Andersons Barricade Herbicide III	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div., Inc. d/b/a Free Flow Fert. PO Box 119 Maumee, OH 43537 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

SEP 22 2003

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Alternate Brand Name per PR Notice 98-10 : Tee Time Fertilizer with Barricade Herbicide 14-0-28

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
				<input checked="" type="checkbox"/> Glass	<input type="checkbox"/> Paper
				Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 50 lbs.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Dawn L. Walters		Title Regulatory Administrator		Telephone No. (Include Area Code) 334-262-2766	
<p align="center">Certification</p> <p>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment both under applicable law.</p>					6. Date Application Received (Stamped)
2. Signature <i>Dawn L. Walters</i>		3. Title Regulatory Administrator			
4. Typed Name Dawn L. Walters		5. Date 8/11/03			

Tee Time[®]

Fertilizer with **Barricade[®]** HERBICIDE

14-0-28

QUARANTEED ANALYSIS:
 Total Nitrogen (N) 14%
 3.28% Water Insoluble Nitrogen
 6.05% Urea Nitrogen
 4.66% Slowly Available Water Soluble Nitrogen*
 Soluble Potash (K₂O) 28%
 Total Sulfur (S) 9.52%
 9.52% Combined Sulfur (S)
 Plant nutrients derived from methylene urea, urea and sulfate of potash.
 * Slowly Available Nitrogen from Methylene Urea

ACTIVE INGREDIENT:
 Proflamline (N², N²-Di-g-propyl-2,4-dinitro-6-(trifluoromethyl)-D-phenylenediamine) 0.48%
 Inert Ingredients 99.52%
 Total 100.00%

EPA Est. No. 9198-OH-1 EPA Reg. No. 9196-125

Net Wt. 50 Lbs. (22.68 kg)
 Treats Up To 16,868 Sq. Ft.

38 - (ATTB)KBR5 ATT144BR5

MAXIMUM ANNUAL RATES:
 The product should not be applied above a maximum annual rate of 313 lbs./A or 7.1 lbs./1,000 sq. ft.

Turf Sodden and Heavy Traffic
 Maximum Rate / Calendar Year by Turf Grass Species

Turf Grasses	lb./A	lb./1,000 sq. ft.	lb./A
Bermudagrass, Bahiagrass, Centipedegrass, Seashore Paspalum, Tall Fescue (including Lufkin), Zoysia, Kentucky Bluegrass, Perennial Ryegrass, Creeping Red Fescue, St. Augustine, Creeping Bentgrass	300 104 100	6.8 2.3 2.3	1.8 0.75 0.85

* These are the maximum rates per calendar year by species indicated. They are based on the maximum annual rate of 313 lbs./A or 7.1 lbs./1,000 sq. ft. Newly applied or stopped Bermudagrass sods may be temporarily restricted. Suppression of Poa, Cynodon and Paspalum due to reduced product rates used in spring applications.
 † Do not apply more than 313 lbs./A of this product per calendar year.
 ‡ If making more than one application, do not apply more than 150 lbs. of the product within any 60 day period.
 † Use higher rates of the product to achieve higher levels of fertility and weed control for each turf type.

WEEDS CONTROLLED:
 When used at the recommended rate on Turf Sodden and Heavy Traffic, this product will control: Common Broomrape, Annual Bluegrass (Poa annua), Goosegrass, Annual Fescue, Barnyardgrass, Broadleaf Signalgrass, Johnsongrass (from seed), Paspalum, Carpetgrass, Common Chickweed, Common Lambsquarters, Common Purslane, Herb, Knotweed, Pigweed, Prostrate Spurge, Shearwatergrass, Yellow Woodruff (from seed).

Do not exceed the maximum annual application rate recommended for the turf species. All applications must be made prior to germination of weed seeds. This product will not control established weeds.

- In many areas, a single application of 138 to 313 lbs./A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, apply an initial treatment of 138 to 209 lbs./A followed after 60 days with a second treatment of 108 to 175 lbs./A for season long goosegrass control.
- Suppression only.
- Winter weeds are best controlled by applications made prior to germination: late summer, fall or winter drifts.

Overseeding/Reseeding Intervals Table:

This Product Rate (Lbs./A)	Overseeding/Reseeding Interval (Months) Based Upon Seeding Method and Predominant Soil Conditions for the Season			
	Broadcast Seeding		Drill Seeding	
	Wet & Warm	Cool & Dry	Wet & Warm	Cool & Dry
104	4	4	4	4
300	8	12	8	8
313	12	16	12	12

the professional's partner[®]

NOTIFICATION
 SEP 22 2003



KEEP OFF OF LEAD OF CHILDREN
CAUTION
 For relative pre-emergence control of grass and broadleaf weeds, established turfgrasses.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

STATEMENTS OF PRACTICAL TREATMENT

If in eyes: Flush thoroughly with water for several minutes. Get medical attention if irritation persists.
If on skin: Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists.
If inhaled: Remove victim to fresh air.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or at an incineration facility, or if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in turf grasses and landscapes. This product controls susceptible weeds by inhibiting weed seeds germination and root development. For most effective weed control this product must be incorporated by 1/2 inch of rainfall, irrigation or shallow mechanical incorporation. Incorporation should take place as soon as possible after application and should not be delayed past 14 days from the date of application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

Do not graze or feed livestock forage cut from areas treated with this product.


APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated commercial application equipment.

ESTABLISHED TURF

This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established grass turf. The species of turf grass determines the maximum amount of material that may be applied during a year.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation within 14 days after application and prior to weed germination. See the map below for approximate crabgrass germination dates.

CRABGRASS GERMINATION DATES**Approximate Date**

- After May 30
- After May 10
- After April 20
- After March 20
- January 1 to March 20

USE PRECAUTIONS

The following use precautions apply to the use of this product in turf grasses and lawns:

Application of this product may thin emerged annual bluegrass and newly over-seeded grasses.

Do not apply to OVERSEEDED TURF for 60 days after seeding or until after the second mowing, whichever ever is longer. Injury to desirable seedlings is likely if this product is applied before secondary roots are in the second inch of soil, not thatch plus soil.

Do not apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or past damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as diking may result in reduced weed control.

Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

TURF SPECIES AND RATES

For season long control, this product may be applied as a single application, or if needed, followed by a second application for increased length of control. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rate within the listed range for longer control. Longer control is desirable in areas where crabgrass germinates between January 1 and April 20 (See Crabgrass Seed Germination Data Map).

Select maximum use rate based on turf grass species. All applications must be made prior to germination of the target weed seeds. This product will not control established weeds.

See turf rates as listed on front of bag.

OVERSEEDING/RESEEDING

This product will inhibit the germination of turf species if overseeded/reseeded too soon after application. The interval between application and overseeding/reseeding is dependent on the application rate, soil temperature, and soil moisture. A shorter interval is possible if growing conditions are warm and wet, while a longer interval is needed if conditions are cool and dry.

Use the Overseeding/Reseeding Intervals Table on front of bag as a guide when choosing the appropriate overseeding/reseeding interval for each situation.

SUGGESTED SPREADER SETTINGS

Please consult your Tee Time Spreader Setting Card.

MANUFACTURER'S STATEMENT

This product conforms to the ingredient statement on the label. Manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage and handling of this product.

THIS PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO WARRANTIES, EXPRESS OR IMPLIED. IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND.

Manufactured by
**the professional's
partner®**



Lawn Fertilizer Division
P.O. Box 119
Maumee, Ohio 43537