6	•
roval expires 2-28-95	<u>;</u>
dentifier Number	1
	l
	l
Classification	1
Restricted	
Nestricted	
	l
Section 3(c)(3)	1
on and labeling	ļ
_	
	l
	1
	1
	ĺ
FICATION	
6.0 2002	
2 8 2003	
	ļ
aryl	
en made to the	
y false statement	
be in violation of	ŀ
	ł
ĺ	
,	
ion.)	
ćiude Area Code)	
**	
Application	
Application	
Stamped)	
• •	
•	
•	

≎EPA	υ Environme ntal	United States NVIFONMENTAL Protection Agency Washington, DC 20460		V	Registration Amendment Other		OPP Identifier Number	
		Application	for Pestici	ide - Section	1	<u> </u>		
1. Company/Product Number 9198-146		2. EPA	2. EPA Product Manager Hebert		3. Proposed Classification			
4. Company/Product (Name) The Andersons 8% Granular Sevin			PM#					
5. Name and Address of Ap The Andersons Lawr Fert. PO Box 119 Maumee, OH 43537	n Fert. Div., Inc. d/		(b)(i), n to: EPA I		nilar or identica	t in com	FIFRA Section 3(c)(3) apposition and labeling	
			Section -	16				
Agency le		Final printed labe Agency letter dat "Me Too" Applica	Application.					
Notification - Explain	n below.			Other - Explain be	slow.	. [MAY 2 8 2003	
This notification is consisten labeling or the confidential s to EPA. I further understand FIFRA and I may be subject	statement of formula of thi d that if this notification is	nis product. I und s not consistent w	lerstand that it is a vith the terms of Pl er sections 12 and	a violation of 18 U.S. R Notice 98-10 and d 14 of FIFRA.	C. Sec. 1001 to v	willfully m	ake any false statement	
			Section - I	li				
1. Material This Product Wil	il Be Packaged in:				T			
Child-Resistant Packaging Yes No	Unit Packaging Yes No If "Yes"	No. per	Water Soluble P Yes No If "Yes"	Packaging No. per	P	nteiner fletal flestic iless aper		
* Certification must be submitted	Unit Packaging wgt.	container	Package wgt	container		ther (Sp		
3. Location of Net Contents	Information Container	4. Size(s) Retail	l Container 50 lbs.	5. Lo	Cation of Label D	Direction	8	
6. Manner in Which Label is	Affixed to Product	Lithograp Paper glo Stenciled	ph ued d	Other				
			Section - I	V				
1. Contact Point (Complete	items directly below fo	or identification	of individual to b	e contacted, if nec	essery, to proce	ss this q	pplication.)	
Name Dawn L. Walters			itle Regulatory Admii	nistrator		lephone 4-262-27	No: îlhclude Area Code) 66° ***	
	ements I have made on t ny knowlinglly false or n I aw.		l attachments the			te.	Received (\$tamped)	
2. Signature		I * *	Title		****	•	• ••	
Dawn & walte	المار	R	Regulatory Adminis	strator			••••	
4. Typed Name Dawn L. Walters		5.	Date 5/11	1112		į	• • • •	



8% Granular Insecticide with Carbaryl

Active Ingredient: Carbaryl (1-naphtyl N-methylcarbamate) 8.0% OTHER INGREDIENTS: 92.0% TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing before rause.

User Safety Requirements: Applicators and other handlers must wear long-sleeved shirt and long pants, and shoes plus socks. Keep and wash PPE separately from other laundry, use detergent and hot water. Wash thoroughly with soap and water after handling this product. and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Carbaryl is an N-Methyl Carbamate Insecticide

GENERAL: Contact a physician immediately in all cases of suspected poisoning. Transport patient to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in enimals, contact a veterinarien.

Call a poison control center or doctor for treatment advice.

Las a posson control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce voniting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person, hold eye open and hinse slowly and gentity with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue

Intaing.
Call a poison control center or doctor for treatment advice.
Take off contaminated clothing.
Rinse skin immediately with plenty of water for 15-20 minutes.
Call a poison control center or doctor for treatment advice.

If inhalod

Move person to fresh air. If person is not breathing, call 911 or an ambutance, then give artificial

- in person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.

- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ANTIDOTE STATEMENT: ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

NOTE TO PHYSICIAN: Carbaryl is a N-Methyl carbamate that inhibits cholinesterase. Alropine is antidotal. Do not use 2-PAM (pyridine-2-aldoxime methiodide), opiates or cholinesterase inhibiting drugs. Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning equipment or by disposal of wastes.

DIRECTIONS FOR USE

ξ.

It is a violation of Federal Law to use this product in any manner inconsistent with its labeling. Read entire label before using this product.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, of for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or

LAWNS, RECREATIONAL, AND ORNAMENTAL TURF AREAS: For control of turforass pests, apply granules uniformly to the turtgrass. For optimum control, follow specific directions for proper watering or irrigation after application. Do not allow public use of treated area during application.

Apply 1.7 to 2.4 lbs. per 1000 sq. ft. for the following pests:

Ants, Armyworms, Centipedes, Chingers, Chinch Bugs, Crickets, Cutworms, Darkling Ground Beetles, Earwigs, Essex Skipper, European Chafer, Fall Armyworms, Fiery Skipper, Fleas, Grasshoppers, Green June Beetles, Green June Beetle Grubs, Lawn Moths (Sod Webworm). Leathoppers, Lucerne Moth Larvae, Millipedes, Sowbugs, Spittlebugs, Springtails, Ticks, Yellowstriped Armyworms.

For general insect control, apply granules uniformly when grass is dry. For most effective controi, lawn should be watered lightly after application to remove chemical from granule. Repeat applications may be made as needed to control new infestations.

Apply 2.4 lbs. per 1000 sq. ft. for the following pests:

White Grubs (such as Japanese Beetle Grubs, Chafer Beetle Grubs, Phyliophaga spp. Grubs, and Annual White grubs), Bluegrass billibug larvae, European crane fly larvae.

For white grub control, applications should be made when grubs are feeding near the soil surface, usually during late March through May or July to early September, or as recommended by local agricultural extension agents. For most effective control, turf should be watered thoroughly after treatment to remove the chemical from the granules. Repeat applications may be made as needed to control new infestations.

Nuisance Pest Control in and Around Flower Beds and Omemental Plantings and Around Buildings:

For control of Ants, Armyworms, Ticks, Centipedes, Crickets, Cutworms, Earwigs, Firebrats, Silverfish, Fleas, Millipedes, Sowbugs, Scorpions, and Spiders: Apply granules uniformly in a 6 foot band around the building. Apply granules at the rate of 4.6 ounces per 120 square feet (6 x 20 foot band) to shrub and flower beds, foundations, ornamental plantings, and lawn or soil areas immediately adjacent to the building. For most effective control, treated areas should be lightly watered after application to remove the chemical from the granules. Receal treatments as necessary to control new infestations.

imported Fire Ant Control:

For use as a mound treatment to control Imported Fire Ants, apply 1.2 to 2.4 ounces by weight over and around each mound. Irrigate mound thoroughly immediately after treatment to remove the chemical from the granules and flush it down into the mound. A minimum of 1 galion of water should be applied to each mound, but do not use more than 2 gallions of water as flooding of the mound may result. Apply water gently to avoid disturbing ants and causing migration before the chemical can be effective. For best results, apply when ants are least active (65° to 80° F), or in early morning or late evening hours. Treat new mounds as they

Control of ticks which transmit Lyme Disease:

For control of juvenile and adult Ixodes spp. ticks (Deer Tick, Bear Tick, and Black Legged Tick). Apply granules at the rate of 2.0 to 2.4 lbs. of formulated product per 1000 sq. ft.

Begin applications in late spring or early summer (May/June). Uniformly apply granules to entire lawn, perimeter wooded areas, and property boundaries including shrub beds, ornamental plantings, and wooded areas where exposure to ticks may occur. For optimum results, irrigate treated areas after application to remove chemical from the granules.

STORAGE AND DISPOSAL

STORAGE: Store unused product in original container only, in cool, dry area out of the reach of children and animals. Keep container closed and away from food, feedstuffs and dom water supplies. Do not store in areas where temperatures frequently exceed 100° F. PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard in trash.

Suggested Spreader Settings Suggested spreader settings are not intended to replace calibration

Suggessed syreamer security are full released to replace cateriation.

10 Institute of the product, 400 Research caterials suggested to product, 50 Institute of the LOW RATE, 57 No.75000 Sq. Ft. 50 Institute 20,000 sq. ft. (v.57 oct?) Web 5.52 Inst. Affiliates at the LOW RATE, 24 debut 1,000,60 Ft. 55 Institute attitute (GOH RATE, 24 debut 1,000,60 Ft.)

	GROUND	WIDTH OF	SPREADER SETTINGS	
The Andersons AccuPro/ SR-2000, AccuPro 1000 (cone 6)	3 mpff	COVERAGE	LOW CATE	HIGH RATE
The Andersons SS-2 Drap	-		3	4
Scotts Rotaries R-6, R-6A (Cone 6)	2 mph	11 feet	H 3/4	ł 1/2
Lety Modet W (PTO at 450 rpm)	4.5 mph	в Иди	41 HLC	4 1/2 #I-C
Vicon (03 Series)	4.5 mph	20 Post	12 3/4	14 3/4

IMPORTANT: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of terrain. Calculate the size of the area to be treated and do nt over or under apply. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary, in order to assure correct rate of application. Close spreader when filling, stopping or turning, do not hand spread.

Broadcast spreaders

For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

Drop spreaders

First apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

To determine square footage of application area, follow calculation chart.

__length __width __Total Sq. Ft.

Non-lawn areas

IMPORTANT: READ REFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MER-CHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCI-DENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. USE contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS: YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION

> Manufactured by: The Andersons Lawn Fertilizer Division, Inc. P. O. Box 119, Maumee, OH 43537

EPA Reg. No. 9198-146

EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A Underlined letter is first letter used in run code on bag.

The Andersons logo & Professional Turf are trademarks of The Andersons

Net Wt. 50 Lbs. (22,68 kg) Covers up to 29,400 Sq. Ft.

E3 (BLKP1929B)

MAY 2 8 2003